

## Synthesis of 4-Hydroxyquinolines from the Baylis-Hillman Adducts of *o*-Nitrobenzaldehydes

Jae Nyoun Kim,\* Ka Young Lee, Heui-Suk Ham, Hyoungh Rae Kim,<sup>†</sup> and Eung K. Ryu<sup>†</sup>

Department of Chemistry and Institute of Basic Science, Chonnam National University, Kwangju 500-757, Korea

<sup>†</sup>Korea Research Institute of Chemical Technology, P.O. Box 107, Yusong, Taejeon 305-600, Korea

Received December 13, 2000


**Keywords :** 4-Hydroxyquinolines, Baylis-Hillman adducts, *o*-Nitrobenzaldehydes, Photoreaction.

The Baylis-Hillman reaction is one of the most powerful carbon-carbon bond-forming methods in organic synthesis.<sup>1</sup> The Baylis-Hillman adducts, which are allylic alcohol derivatives, can be formed most often by the reaction of activated vinyls and carbonyl compounds.<sup>1</sup> Besides the usefulness of these Baylis-Hillman adducts themselves, further derivatization with various nucleophilic reagents toward synthetically useful compounds has been studied in depth by us and other groups.<sup>2</sup> There were some reported papers on the formation of heterocyclic compounds including quinolines from the Baylis-Hillman adducts.<sup>3</sup>

Quinolines and their derivatives occur in numerous natural products.<sup>4</sup> Many quinolines display interesting physiological activities and have found attractive applications as pharmaceuticals and agrochemicals as well as being general synthetic building blocks.<sup>4b</sup> Although many synthetic methods have been developed for the preparation of quinolines,<sup>5</sup> due to their great importance, the development of novel synthetic methods remains an active research area.<sup>6</sup>

Recently, we have reported on the synthesis of 4-hydroxy-3-ethoxycarbonylquinoline *N*-oxide derivatives from the Baylis-Hillman adducts of 2-nitrobenzaldehydes in acidic conditions.<sup>3a</sup> As a continuous work, we intended to examine the possibility of transforming the Baylis-Hillman adducts of 2-nitrobenzaldehydes into the quinoline *N*-oxides by the photochemical method.

A solution of **1a** in ethanol (0.16 M solution) was irradiated with 250 nm wavelength in a quartz reaction vessel.<sup>7</sup>



Scheme 1

We could isolate quinoline *N*-oxide **3a** in trace amount from the reaction. Instead the deoxygenated 4-hydroxy-3-ethoxycarbonylquinoline **2a** (39%) was obtained as the major product.<sup>8</sup> The reaction might proceed *via* the *aci*-nitro compound **I**, which was generated through the benzylic hydrogen abstraction by the excited *ortho*-nitro group.<sup>9</sup> **I** was converted to the quinoline *N*-oxide **3a** as described in our previous paper.<sup>3a</sup> Elimination of oxygen atom of **3a** and the fate of oxygen atom is uncertain at this point (*vide infra*). In other cases **1b–d** and the nitrile analogue **1e**, we could obtain the similar results as in Table 1.<sup>8</sup>

In the reaction mixture we could observe the quinoline *N*-




Table 1. Photoreaction of the Baylis-Hillman Adducts **1**

| entry | B-H adducts ( <b>1</b> ) | conditions  | products ( <b>2</b> ) | yields (%)      |
|-------|--------------------------|---|-----------------------|-----------------|
| a     |                          | EtOH/250 nm<br>EtOH/350 nm<br>THF/250 nm <sup>a</sup> |                       | 39<br>25<br>30  |
| b     |                          | EtOH/250 nm   |                       | 23              |
| c     |                          | EtOH/250 nm   |                       | 30              |
| d     |                          | EtOH/250 nm   |                       | 26              |
| e     |                          | EtOH/250 nm   |                       | 36              |
| f     |                          | EtOH/250 nm   |                       | 26 <sup>b</sup> |

<sup>a</sup>0.03 M concentration. <sup>b</sup>benzoic acid (25%) was obtained.

oxides in trace amounts (< 5%), which might indicate the intermediacy of quinoline *N*-oxides for the formation of quinolines **2**. Moreover, irradiation of pure **3a** (250 nm, 20 h) in ethanol produced **2a** in 58% isolated yield. Such deoxygenation of quinoline *N*-oxides to quinolines indeed occurred very slowly in ethanol solution without UV irradiation.

Change of the reaction conditions such as wavelength, solvent, or concentration did not improve the yields (entry a in Table 1). Heating **1a** in ethanol (70–80 °C, 48 h) or diphenyl ether (200 °C, 48 h) without UV irradiation did not give any quinoline **2a** nor quinoline *N*-oxide.

Photochemical rearrangements and fragmentations of *o*-nitrobenzyl compounds were well-known.<sup>9</sup> Transfer of oxygen atom of nitro group to the benzylic position and concomitant removal of the alkoxy, carboalkoxy, or phosphate group occurs. In these respects the *o*-nitrobenzyloxy moiety was studied and used as a photochemically labile protecting group.<sup>9</sup> In order to examine the possibility of the Baylis-Hillman adducts as a photochemical protecting group, we performed the photochemical reaction of *O*-benzoyl analogue **1f** in the same reaction conditions. From the reaction we could obtain **2a** (26%) and benzoic acid (25%) as expected. However, to our disappointment, starting material **1f** was recovered in 64% yield.

**Acknowledgment.** This work was supported by Korea Research Foundation Grant (KRF-2000-015-DP0275). The support of the Korea Basic Science Institute (Kwangju branch) is also acknowledged.

### References and Notes

- (a) Ciganek, E. *Organic Reactions*; John Wiley & Sons: New York, 1997; Vol. 51, pp 201–350. (b) Drewes, S. E.; Roos, G. H. P. *Tetrahedron* **1988**, *44*, 4653. (c) Basavaiah, D.; Rao, P. D.; Hyma, R. S. *Tetrahedron* **1996**, *52*, 8001.
- (a) Kim, H. S.; Kim, T. Y.; Lee, K. Y.; Chung, Y. M.; Lee, H. J.; Kim, J. N. *Tetrahedron Lett.* **2000**, *41*, 2613. (b) Lee, H. J.; Seong, M. R.; Kim, J. N. *Tetrahedron Lett.* **1998**, *39*, 6223. (c) Lee, H. J.; Kim, H. S.; Kim, J. N. *Tetrahedron Lett.* **1999**, *40*, 4363, and references cited therein.
- (a) Kim, J. N.; Lee, K. Y.; Kim, H. S.; Kim, T. Y. *Org. Lett.* **2000**, *2*, 343. (b) FAMILONI, O. B.; KAYE, P. T.; KLAAS, P. J. *J. Chem. Soc., Chem. Commun.* **1998**, 2563. (c) BODE, M. L.; KAYE, P. T. *J. Chem. Soc., Perkin Trans. 1* **1993**, 1809. (d) BODE, M. L.; KAYE, P. T. *J. Chem. Soc., Perkin Trans. 1* **1990**, 2612.
- (a) Michael, J. P. *Nat. Prod. Rep.* **1997**, *14*, 605. (b) Balasubramanian, M.; Keay, J. G. In *Comprehensive Heterocyclic Chemistry II*; Katritzky, A. R., Rees, C. W., Scriven, E. F. V., Eds.; Pergamon Press: Oxford, 1996; Vol. 5, pp 245–300.
- Jones, G. In *Comprehensive Heterocyclic Chemistry II*; Katritzky, A. R., Rees, C. W., Scriven, E. F. V., Eds.; Pergamon Press: Oxford, 1996; Vol. 5, pp 167–243.
- (a) Ranu, B. C.; Hajra, A.; Jana, U. *Tetrahedron Lett.* **2000**, *41*, 531. (b) Cho, C. S.; Oh, B. H.; Shim, S. C. *Tetrahedron Lett.* **1999**, *40*, 1499. (c) Cacchi, S.; Fabrizi, G.; Marinelli, F. *Synlett* **1999**, 401. (d) Suginome, M.; Fukuda, T.; Ito, Y. *Org. Lett.* **1999**, *1*, 1977. (e) Katritzky, A. R.; Arend, M. J. *Org. Chem.* **1998**, *63*, 9989.
- For the photolysis experiments a Raynot photochemical reactor (model RPR-2080, the Southern N. E. Ultraviolet Co.) was used. The built-in monochromatic UV light sources (RUL-250 nm UV lamp) was positioned approximately 17 cm from the reaction quartz tube.
- Typical procedure for the synthesis of **2a** and some selected spectroscopic data: A stirred solution of **1a** (400 mg, 1.59 mmol) in ethanol (10 mL) was irradiated (250 nm) for 30 h. After removal of ethanol, column chromatographic purification (CH<sub>2</sub>Cl<sub>2</sub>/EtOH, 14 : 1) gave **2a** as a white solid, 135 mg (39%); mp 267–268 °C (lit.<sup>3a</sup> 269–270 °C); IR (KBr) 3434, 3169, 2982, 2904, 1706, 1623, 1529, 1476, 1380, 1292, 1202, 1141, 766 cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.29 (t, *J* = 7.1 Hz, 3H), 4.23 (q, *J* = 7.1 Hz, 2H), 7.42 (t, *J* = 8.1 Hz, 1H), 7.63 (d, *J* = 8.1 Hz, 1H), 7.71 (t, *J* = 8.1 Hz, 1H), 8.17 (d, *J* = 8.1 Hz, 1H), 8.56 (s, 1H), 12.41 (br s, 1H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 14.52, 59.77, 109.94, 118.98, 124.90, 125.82, 127.44, 132.61, 139.15, 145.11, 165.00, 173.68.  
**2e**: 36%, mp 303–305 °C (dec.) (lit.<sup>10</sup> 301 °C); IR (KBr) 3174, 2959, 2873, 2223, 1618, 1561, 1535, 1352, 760 cm<sup>-1</sup>; <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 7.52 (td, *J* = 7.6 and 1.2 Hz, 1H), 7.61 (d, *J* = 8.1 Hz, 1H), 7.79 (td, *J* = 7.1 and 1.4 Hz, 1H), 8.25 (dd, *J* = 8.2 and 1.3 Hz, 1H), 8.53 (s, 1H); <sup>13</sup>C NMR (CD<sub>3</sub>OD) δ 94.55, 115.71, 118.99, 125.30, 125.44, 126.01, 133.74, 139.59, 146.43, 176.50; MS (70 eV) *m/z* (rel intensity) 63 (45), 76 (33), 114 (41), 115 (49), 142 (93), 170 (M<sup>+</sup>, 100).
- (a) Woodrell, C. D.; Kehayova, P. D.; Jain, A. *Org. Lett.* **1999**, *1*, 619. (b) Pirrung, M. C.; Lee, Y. R.; Park, K.; Springer, J. B. *J. Org. Chem.* **1999**, *64*, 5042. (c) Pirrung, M. C.; Shuey, S. W. *J. Org. Chem.* **1994**, *59*, 3890. (d) Walker, J. W.; Reid, G. P.; McCray, J. A.; Trentham, D. R. *J. Am. Chem. Soc.* **1998**, *110*, 7170. (e) Givens, R. S.; Matuszewski, B. *J. Am. Chem. Soc.* **1984**, *106*, 6860. (f) Amit, B.; Zehavi, U.; Patchornik, A. *J. Org. Chem.* **1974**, *39*, 192. (g) Givens, R. S.; Athey, P. S.; Kueper, L. W., III; Matuszewski, B.; Xue, J.-y. *J. Am. Chem. Soc.* **1992**, *114*, 8708. (h) Givens, R. S.; Athey, P. S.; Matuszewski, B.; Kueper, L. W.; Xue, J.-y.; Fister, T. *J. Am. Chem. Soc.* **1993**, *115*, 6001. (i) Pillai, V. N. R. *Synthesis* **1980**, 1. (j) Givens, R. S.; Kueper, L. W. III. *Chem. Rev.* **1993**, *93*, 55.
- Bredereck, H.; Effenberger, F.; Botsch, H.; Rehn, H. *Chem. Ber.* **1965**, *98*, 1081.