

Improving the Regioselectivity in the Cyclic Iminocarbonate Rearrangement: Enantioselective Synthesis of the Paclitaxel Side Chain

Gae Y. Cho, Kyung Mi An, and Soo Y. Ko*

Department of Chemistry and Division of Molecular Life Sciences, Ewha Womans University, Seoul 120-750, Korea

Received January 18, 2001

Keywords : Cyclic iminocarbonate rearrangement, Enantioselective synthesis, Paclitaxel side chain.

β -Amino- α -hydroxy acids represent an important class of compounds with wide biological activities.¹ In particular, *N*-benzoyl-*syn*-phenylisoserine, the C-13 side chain of Paclitaxel, has been the target of numerous synthetic efforts as the interest in and the paucity of the anti-cancer agent has prompted a wide search for alternative sources for the parent drug and its fragments.²⁻⁹ A key in the side chain synthesis is the stereocontrol of the C-2/C-3 vicinal carbons, and various strategies have been devised to address the relative (*syn*) as well as absolute (*2R,3S*) stereocontrol. These include asymmetric 1,2-induction,³ use of chiral auxiliary,⁴ enzymatic resolution,⁵ and those employing powerful asymmetric oxidation processes such as catalytic asymmetric epoxidation,⁶ dihydroxylation⁷ and aminohydroxylation⁸ (followed by stereospecific functional group transformations as necessary).


Employing Sharpless's asymmetric dihydroxylation (AD) as the key step, one faces two issues in the synthesis of the Paclitaxel side chain: the regiocontrol and the stereocontrol, particularly in a *relative* sense (*i.e.*, diastereocontrol). Thus, a (*2R,3S*)-cinnamate diol, available *via* the AD, needs to be converted to *N*-benzoyl-(*2R,3S*)-phenylisoserine with a substitution of *N*-benzoyl group *regioselectively* at the C-3 and with *net-retention of configuration*. A reaction sequence involving an *N*-benzoyl substitution with inversion of configuration at C-3, a perhaps more straightforward synthetic plan that would require (*2R,3R*)-cinnamate diol starting material, is not applicable as anti(*erythro*)-diols are not directly accessible *via* AD process.^{10,11} While the reported syntheses solved the issue of regiocontrol by taking advantage of the fact that the C-3 was activated as benzylic, this could cause the stereocontrol to be less than complete during the necessary double inversion (net-retention) *N*-substitution steps due to the participation of *S_N1*-type reaction pathways. Thus, one AD-based synthesis endured an incomplete (6 : 1) regioselection,^{7a} while another suffered from a modest (4 : 1) diastereoselection.^{7b}

Our solution to this double-faced problem is to use the cyclic iminocarbonate rearrangement, which we reported recently (Scheme 1).¹² The rearrangement converts *syn*-diols (AD products) to protected *syn*-amino alcohols in a one-pot operation; the configurations at both carbinol carbons are retained. Thus, ethyl *trans*-cinnamate (**1**) was dihydroxylated using AD-mix- α to give the (*2R,3S*)-cinnamate diol (**2**, 85% yield, 96% ee¹³), which was subjected to the cyclic iminocarbonate rearrangement protocol. While the original procedure

(using tetrabutylammonium bromide as the nucleophile) had been initially reported to produce preferentially the C-2 nitrogen isomer (**4a**, the "wrong" regioisomer for the synthesis of the side-chain, 1:2.7 regioselectivity),^{12a} subsequent works led to a discovery that using LiI nucleophile in acetonitrile/C₂H₄Cl₂ reversed the regioselection of the rearrangement toward the desired C-3(benzylic)-nitrogen isomer (**3a**, 3 : 1 regioselectivity).^{12b} Further optimizations of the reaction conditions have now resulted in a great improvement in regioselectivity (>20 : 1 for the desired isomer **3a**, 82% combined yield) when the rearrangement was performed with MgI₂ as the nucleophile in acetonitrile/C₂H₄Cl₂ at room temp. No diastereomeric products were observed and the enantiomeric purity of the product was confirmed to be as high as that of the starting diol **2**.¹³ The stereochemical integrity, therefore, was fully maintained during the rearrangement. The high diastereoselectivity is presumably due to the fact that the rearrangement (**2** \rightarrow **3a**) involves an intramolecular (cyclic) nucleophilic substitution step and the *syn*-diastereomer, as obtained, is also thermodynamically more stable. Therefore, an *S_N1*-type reaction pathway, if present at all, would also yield the net retention diastereomer preferentially.¹⁴ For an additional advantage, the desired regioisomer **3a** conveniently crystallized out from the crude product (hexane-ethyl acetate).

The rearranged product, ethyl 1-benzoyl-5-(*S*)-phenyloxazolidin-2-one-4-(*R*)-carboxylate (**3a**), is only a short distance away from the Paclitaxel side chain. It has been reported that various deprotection methods under basic or acidic solvolytic conditions usually remove the *N*-benzoyl group first.^{12a} Nevertheless, we sought to cleave selectively the carbonyl group of the oxazolidinone ring, while leaving the *N*-benzoyl group intact, as this would lead to a very atom-economic route to the Paclitaxel side chain. Although our endeavor was met with a partial success, we have devised an efficient synthetic process for our target compound, nonetheless.

When the *N*-benzoyl-oxazolidinone **3a** was treated with Cs₂CO₃ in anhydrous methanol, a mixture of products was obtained, consisting of the oxazolidinone ring-opened product **5** and the *N*-debenzoylated compound **6** in a 1 : 2 ratio (100% combined yield). *N*-Benzoyl-(*2R,3S*)-phenylisoserine methyl ester (**5**), the methyl ester of the Paclitaxel side chain, crystallized out from hexane-ethyl acetate (28% yield from **3a**, 97% ee). The *N*-debenzoylated compound, 5-(*S*)-phenyl-



Scheme 1

oxazolidin-2-one-4-(*R*)-carboxylate methyl ester (**6**), recovered from the mother liquor, could be re-*N*-benzoylated to give compound **3b** (BzCl, DMAP, 93% yield), which may be recycled to furnish additional amounts of the product **5**.

Alternatively, the oxazolidinone **3a** was subjected to exhaustive hydrolytic conditions (10% HCl, reflux for 7 days), and without isolation of the resulting phenylisoserine, the crude hydrolyzed product was *N*-benzoylated under Schotten-Baumann conditions to yield *N*-benzoyl-(2*R*,3*S*)-phenylisoserine (**7**) in 77% (from **3a**). The stereochemical integrity was sustained through the harsh and prolonged reaction conditions.¹⁵

In conclusion, we have achieved a very efficient synthesis of the Paclitaxel side chain, *N*-benzoyl-(2*R*,3*S*)-phenylisoserine, in a three-step sequence employing as key steps the asymmetric dihydroxylation and the regio- and diastereoselective cyclic iminocarbonate rearrangement processes.

Experimental procedure for the MgI₂-catalyzed cyclic iminocarbonate rearrangement. To a solution of (2*R*,3*S*)-cinnamate diol ethyl ester (213 mg, 1.0 mmol) in dichloroethane (15 mL), dibutyltin oxide (305 mg, 1.2 mmol) was added. The mixture was heated to reflux for 4 hr with a concomitant removal of water using a Dean-Stark trap. Benzoyl isothiocyanate (0.187 mL, 1.4 mmol) and triethylamine (0.142 mL, 1.0 mmol) were added and the mixture was heated to reflux for further 2 hr. After it was cooled to rt, the mixture was diluted with MeCN (30 mL) and MgI₂ (304 mg, 1.0 mmol) was added. It was then stirred at rt overnight. Aqueous extractive work-up (EtOAc - aq. NaHCO₃, *dil.* HCl, then brine) yielded the crude product (277 mg, 82%, >20 : 1 regioisomeric mixture **3a** : **4a**). Crystallization from hexane-EtOAc produced pure ethyl 1-benzoyl-5-(*S*)-phenyloxazolidin-2-one-4-(*R*)-carboxylate (**3a**, 75%).¹⁶

Acknowledgment. This work was supported by the KRF (2000-015-DP0271). G.Y.C. is grateful to the Brain Korea 21 project for a graduate fellowship.

References and Notes

- (a) Cole, D. C. *Tetrahedron* **1994**, *50*, 9517. (b) Cardillo, G.; Tomasini, C. *Chem. Soc. Rev.* **1996**, *25*, 117. (c) *Enantioselective Synthesis of β-Amino Acids*; Juaristi, E. Ed.; VCH Publishers: New York, 1996.
- Reviews: (a) Boa, A. N.; Jenkins, P. R.; Lawrence, N. J. *Contemporary Org. Synth.* **1995**, *47*. (b) Nicolaou, K. C.; Dai, W.-M.; Guy, R. K. *Angew. Chem. Int. Ed. Engl.* **1994**, *33*, 15. (c) Boge, T. C.; Georg, G. I. in reference 1(c), Chapter 1.
- (a) Kang, S. H.; Kim, C. M.; Youn, J.-H. *Tetrahedron Lett.* **1999**, *40*, 3581. (b) Tomasini, C.; Vecchione, A. *Org. Lett.* **1999**, *1*, 2153. (c) Cardillo, G.; Gentilucci, L.; Tolomelli, A.; Tomasini, C. *J. Org. Chem.* **1998**, *63*, 2351. (d) Nocioni, A. M.; Papa, C.; Tomasini, C. *Tetrahedron Lett.* **1999**, *40*, 8453. (e) Dondoni, A.; Perrone, D.; Semola, T. *Synthesis* **1995**, 181. (f) Lee, K.-Y.; Kim, Y.-H.; Park, M.-S.; Ham, W.-H. *Tetrahedron Lett.* **1998**, *39*, 8129. (g) Denis, J.-N.; Correa, A.; Greene, A. E. *J. Org. Chem.* **1991**, *56*, 6939. (h) Hanessian, S.; Sanceau, J. Y. *Can. J. Chem.* **1996**, *74*, 621. (i) Palomo, C.; Cossio, F. P.; Aizpuru, J. M.; Mielgo, A.; Aurrekoetxea, N. *Tetrahedron Lett.* **1990**, *31*, 6429. (j) Davis, F. A.; Reddy, T.; Reddy, R. E. *J. Org. Chem.* **1992**, *57*, 6387.
- (a) Georg, G. I.; Mashava, P. M.; Akgün, E.; Misteard, M. W. *Tetrahedron Lett.* **1991**, *32*, 3151. (b) Kanazawa, A. M.; Denis, J.-N.; Greene, A. E. *J. Org. Chem.* **1994**, *59*, 1238. (c) Kanazawa, A. M.; Denis, J.-N.; Greene, A. E. *J. Chem. Soc., Chem. Commun.* **1994**, 2591. (d) Swindell, C. S.; Tao, M. *J. Org. Chem.* **1993**, *58*, 5889. (e) Hattori, K.; Yamamoto, H. *Tetrahedron* **1994**, *50*, 2785. (f) Hattori, K.; Miyata, M.; Yamamoto, H. *J. Am. Chem. Soc.* **1993**, *115*, 1151. (g) Bunnage, M. E.; Davies, S. G.; Goodwin, C. J. *J. Chem. Soc., Perkin Trans. 1* **1994**, 2385. (h) Bunnage, M. E.; Davies, S. G.; Goodwin, C. J. *J. Chem. Soc., Perkin Trans. 1* **1993**, 1375. (i) Commerçon, A.; Bézard, D.; Bernard, F.; Bourzat, J. D. *Tetrahedron Lett.* **1992**, *33*, 5185. (j) Farina, V.; Hauck, S. I.; Walker, D. G. *Synlett* **1992**, 761. (k) Ojima, I.; Habus, I.; Zucco, M.; Park, Y. H.; Sun, C. M.; Brigaud, T. *Tetrahedron* **1992**, *48*, 6985. (l) Ojima, I.; Habus, I.; Zhao, M. *J. Org. Chem.* **1991**, *56*, 1681. (m) Mukai, C.; Kim, I. J.; Furu, E.; Hanaoka, M. *Tetrahedron* **1993**, *49*, 8323. (n) Jost, S.; Gimbert, Y.; Greene, A. E.; Fotiadu, F. *J. Org. Chem.* **1997**, *62*, 6672. (o) Ha, H.-J.; Park, G.-S.; Ahn, Y.-G.; Lee, G.-S. *Bioorg. Med. Chem. Lett.* **1998**, *8*, 1619.
- (a) Kaywer, M. M.; Mihoviolvic, M. D.; Kearns, J.; Feicht, A.; Stewart, J. D. *J. Org. Chem.* **1999**, *64*, 6603. (b) Kearns, J.; Kaywer, M. M. *Tetrahedron Lett.* **1994**, *35*, 2845. (c) Patel, R. N.; banerjee, A.; Howell, J. M.; McNamee, C. G.; Brozowski, D.; Mirfakhrae, D.; Nanduri, V.; Thottathil, J. K.; Szarka, L. *J. Tetrahedron Asymmetry* **1993**, *4*, 2069. (d) Lee, D.; Kim, M.-J. *Tetrahedron Lett.* **1998**, *39*, 2163. (e) Brieva, R.; Crich, J. Z.; Sih, C. J. *J. Org. Chem.* **1993**, *58*, 1068. (f) Wuts, P. G. M.; Gu, R. L.; Northuis, N. M. *Tetrahedron Asymmetry* **2000**, *11*, 2117. (g) Gou, D.-

- M.; Liu, Y.-C.; Chen, C.-S. *J. Org. Chem.* **1993**, *58*, 1287.
(h) Barco, A.; Benetti, S.; Risi, C. D.; Pollini, G. P.; Romagnoli, R.; Zanirato, V. *Tetrahedron Lett.* **1994**, *35*, 9289.
6. (a) Denis, J. N.; Greene, A. E.; Serra, A. A.; Luche, M. J. *J. Org. Chem.* **1986**, *51*, 46. (b) Deng, L.; Jacobsen, E. N. *J. Org. Chem.* **1992**, *57*, 4320. (c) Jacobsen, E. N.; Furukawa, Y.; Martinez, L. E. *Tetrahedron* **1994**, *50*, 4323. (d) Righi, G.; Rumboldt, G. *J. Org. Chem.* **1996**, *61*, 3557. (e) Bonini, C.; Righi, G. *J. Chem. Soc., Chem. Commun.* **1994**, 2767.
7. (a) Wang, Z. M.; Kolb, H. C.; Sharpless, K. B. *J. Org. Chem.* **1994**, *59*, 5104. (b) Koskinen, A. M. P.; Karvinen, E. K.; Siirilä, J. P. *J. Chem. Soc., Chem. Commun.* **1994**, 21. (c) Denis, J. N.; Correa, A.; Greene, A. E. *J. Org. Chem.* **1990**, *55*, 1957. (d) Song, C. E.; Lee, S. W.; Roh, E. J.; Lee, S. *Tetrahedron Asymmetry* **1998**, *9*, 983.
8. (a) Li, G.; Sharpless, K. B. *Acta Chemica Scandinavica* **1996**, *50*, 649. (b) Bruncko, M.; Schlingloff, G.; Sharpless, K. B. *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 1483. (c) Song, C. E.; Oh, C. R.; Roh, E. J.; Lee, S.; Choi, J. H. *Tetrahedron Asymmetry* **1999**, *10*, 671.
9. Kobayashi, S.; Ishitani, H.; Ueno, M. *J. Am. Chem. Soc.* **1998**, *120*, 431.
10. (a) Ko, S. Y.; Malik, M. *Tetrahedron Lett.* **1993**, *34*, 4675. (b) Ko, S. Y.; Malik, M.; Dickinson, A. F. *J. Org. Chem.* **1994**, *59*, 2570. (c) Ko, S. Y.; Lerpiniere, J. *Tetrahedron Lett.* **1995**, *36*, 2101.
11. cf. Wang, L.; Sharpless, K. B. *J. Am. Chem. Soc.* **1992**, *114*, 7568.
12. (a) Cho, G. Y.; Ko, S. Y. *J. Org. Chem.* **1999**, *64*, 8745. (b) Cho, G. Y.; Park, J. N.; Ko, S. Y. *Tetrahedron Lett.* **2000**, *41*, 1789. (c) Park, J. N.; Ko, S. Y.; Koh, H. Y. *Tetrahedron Lett.* **2000**, *41*, 5553.
13. The enantiomeric purity was determined by chiral HPLC (Pirkle column; hexane-*i*-PrOH gradient eluent).
14. When an *anti*(*erythro*)-diol was subjected to the MgI₂-catalyzed rearrangement protocol, the corresponding *syn*(*trans*)-diastereomeric oxazolidinone was obtained, via an apparent *inversion* of configuration at one carbinol carbon. This observation implies a thermodynamic control of stereoselectivity (as opposed to stereospecificity).
15. [α]_D -39.4 (*c* 0.83, EtOH); Lit.^{7a} [α]_D 35.5 (*c* 1.07, EtOH). Anal. Calcd for C₁₆H₁₄NO₄: C, 67.36; H, 5.30; N, 4.91. Found: C, 67.01; H, 5.36; N, 4.91.
16. A full characterization of this compound has been reported elsewhere (see reference 12a).
-