

Solid-Phase Synthesis of Benzopyran Derivatives via Highly Efficient Epoxidation Using Two-Phase Solvents

Young-Dae Gong and Sung-eun Yoo*


The Bio-organic Science Division, Korea Research Institute of Chemical Technology,
The Center for Molecular Design and Synthesis, P.O. Box 107, Yusong, Daedeog Sciencetown, Daejeon 305-606, Korea
Received May 14, 2001

Keywords : Solid-phase synthesis, Combinatorial, Epoxidation, Two-phase solvents, Benzopyranes.

Solid-phase organic synthesis has emerged as a powerful technique in generating combinatorial libraries of small organic molecules useful for drug discovery.¹ Heterocyclic compounds provide scaffolds on which pharmacophores can be arranged to yield potent and selective drugs. A variety of heterocycles have been synthesized on solid support.² In our research program for the development of potassium channel activators, we needed to develop synthetic strategies and chemistries applicable in a combinatorial approach to the various benzopyran derivatives.³ Herein, we would like to report our findings about a new epoxidation condition of the chromene derivatives **3** on a solid-phase where the normal reaction condition failed to produce the desired epoxide products. We then report a successful application of the epoxides **4** to generate the 3-hydroxy-4-amino substituted benzopyran library (Scheme 1).

We selected the Wang resin **1** as a polymer support, as the hydroxy group of the Wang resin is useful in the introduction of 6-amino-chromenes **6** through the carbamate linker which also serves as an efficient protection group for the amino group against the subsequent oxidation and alkylation reactions.⁴ The benzopyran derivatives **6** were finally liberated from the resin by trifluoroacetic acid (TFA).

The carbamate resin **3** was synthesized in a two-step procedure starting from Wang resin as follows (Scheme 1).




Scheme 1. Reagents and conditions: (a) *p*-nitrophenyl chloroformate, pyridine, CH₂Cl₂; (b) 6-amino-2,2-dimethyl chromene **7**, DIPEA, DMA; (c) *m*-CPBA, CHCl₃ : sat. aqueous NaHCO₃ (9 : 1); (d) 2 equiv. Mg(ClO₄)₂, 4 equiv. R₃R₄NH, CH₃CN; (e) TFA : CH₂Cl₂ (1 : 3), **4**, **5**, and **6** are racemates.

In the first step, the 4-nitrophenyl carbonate resin **2** was prepared by adding pyridine in CH₂Cl₂ to the Wang resin **1** in the presence of *p*-nitrophenyl chloroformate in CH₂Cl₂.⁵ The reaction of carbonate resin **2** with 6-amino-2,2-dimethyl chromene and *N,N*-diisopropylethylamine (DIPEA) in *N,N*-dimethylacetamide (DMA) afforded the carbamate resin **3** and the progress of the reaction was verified by the complete disappearance of the carbonate peak at 1760 cm⁻¹ in the IR spectrum.


With the carbamate **3a** in hand, we examined the epoxidation of **3a** under normal oxidation conditions with *m*-chloroperbenzoic acid (*m*-CPBA) in dichloromethane.⁶ However, under this condition we found that the *m*-chlorobenzoic acid added adduct resin **8a** was mostly formed (Scheme 2). This is probably due to the presence of excessive *m*-chlorobenzoic acid which attacked the initially formed epoxide resin **4a** preferentially. In order to resolve this problem we tried to search for other oxidation conditions.

We examined various oxidants such as oxone, dimethyl dioxirane, hydrogen peroxide, *t*-butyl hydrogen peroxide and sodium perchlorate⁶ in dichloromethane and failed to obtain the desired epoxide resin **4a**. The treatment of the polymer bound olefine **3a** with these oxidants in water or acetone also failed to produce **4a**. The failure is most likely due to the poor resin swelling by polar solvents such as water or acetone.

Therefore, we decided to scrutinize the *m*-CPBA condition more carefully, particularly the solvent system. After examining various solvent systems and two-phase



Scheme 2. Reagents and conditions: (a) *m*-CPBA, CH₂Cl₂; (b) 20% TFA, CH₂Cl₂.



Scheme 3. Reagents and conditions: (a) *m*-CPBA (2.2 eq), CHCl₃ : sat. aqueous NaHCO₃ (9 : 1).

solvent systems, we found that the two-phase solvent system comprised of chloroform and saturated aqueous NaHCO₃ was quite satisfactory. Under this condition the desired epoxide resin **4a** was obtained in good yield without the formation of **8a**. We assumed that the success of this reaction was due to the basic aqueous solution's ability to remove excess *m*-chlorobenzoic acid quite effectively. This explanation was well supported by the fact that the use of bases such as triethylamine, DIPEA, NaOH, and NaHCO₃ without water⁶ did not produce the desired epoxidation product. The progress and yield of this *m*-CPBA epoxidation reaction were monitored by measuring the amount of **9a** and the desired products **6a** which were released from the resin **8a** and **5a** by treating the resin with 25% TFA for 3 hrs, respectively.

In a typical epoxidation procedure, the chromene resin **3a** (1.0 g, 1.1 mmol), dissolved in 20 mL of chloroform, was added to *m*-CPBA (455.6 mg, 2.4 mmol) and the mixture was stirred at 0 °C. After 5 min., 2 mL of saturated aqueous NaHCO₃ was added and the mixture was stirred at room temperature for 12 hrs. The resin was filtered and washed with saturated aqueous NaHCO₃ solution, water, water/MeOH, MeOH and dichloromethane. The resin was then dried under a vacuum overnight.

Finally we carried out the ring opening reaction of the polymer bound epoxide **4a** with nine amines to produce the desired benzopyrane products **6** in good overall yields without significant contamination of the by-products (Table 1).

In conclusion, we have demonstrated that the two-phase solvent system for *m*-CPBA oxidation is suitable for the epoxidation of polymer bound chromenes **3** to minimize the

formation of the by-product **8a**. The epoxides **3** can serve as useful intermediates for reacting with various nucleophiles, such as carboxylic acids, alcohols and thiols etc. The hydroxyl compounds **6** can also be used for further combination with acylating agents to preparing diverse chemical libraries for biological evaluation.

Acknowledgment. We are grateful to the Ministry of Science and Technology, and Ministry of Commerce Industry and Energy of Korea for financial support of this research.

References and Notes

- (a) Hermkens, P. H. H.; Ottenheijm, H. C. J.; Rees, D. *Tetrahedron* **1996**, *52*, 4527. (b) Hermkens, P. H. H.; Ottenheijm, H. C. J.; Rees, D. *Tetrahedron* **1997**, *53*, 5643. (c) Gordon, E. M.; Barrett, R. W.; Dower, W. J.; Foder, S. P. A.; Gallop, M. A. *J. Med. Chem.* **1994**, *37*, 1385.
- (a) Thompson, L. A.; Ellman, J. A. *Chem. Rev.* **1996**, *96*, 557. (b) Gong, Y.-D.; Najdi, S.; Olmsted, M. M.; Kurth, M. J. *J. Org. Chem.* **1998**, *63*, 113. (c) Nicolaou, K. C.; Pfefferkorn, J. A.; Roecker, A. J.; Cao, G.-Q.; Barluenga, S.; Mitchell, H. J. *J. Am. Chem. Soc.* **2000**, *122*, 9939. (d) Terrett, N. K.; Gardner, M.; Gordon, D. W.; Kobylecki, R. J.; Steele, J. *Tetrahedron* **1995**, *51*, 8135.
- Jang, I. J.; Yu, K. S.; Shon, J. H.; Bae, K. S.; Cho, J. Y.; Yi, S. Y.; Shin, S. G.; Ryu, K. H.; Cho, Y. B.; Kim, D. K.; Yoo, S.-e. *J. of Clinical Pharm.* **2000**, *40*, 752.
- (a) Kim, S. W.; Hong, C. Y.; Lee, E. J.; Koh, J. S. *Bioorg. Med. Chem. Lett.* **1998**, *8*, 735. (b) Hauske, J. R.; Doff, P. *Tetrahedron Lett.* **1995**, *37*, 3031. (c) Marsh, I. R.; Smith, H.; Bradley, M. *Chem. Commun.* **1996**, 941. (d) Kaljuste, K.; Undén, A. *Tetrahedron Lett.* **1996**, *37*, 3031.
- (a) Rossé, G.; Ouertani, F.; Schröer, H. *J. of Comb. Chem.* **1999**, *1*, 397. (b) Zaragoza, F. *Tetrahedron Lett.* **1995**, *36*, 8677. (c) Ho, C. Y.; Kukla, M. J. *Tetrahedron Lett.* **1997**, *38*, 2799.
- (a) Hetet, C. L.; David, M.; Carreaux, F.; Carboni, B.; Sauleau, A. *Tetrahedron Lett.* **1997**, *38*, 5183. (b) David, P. R. *J. Am. Chem. Soc.* **1996**, *118*, 12246.

Table 1.



	R ₃	R ₄	Yield (%)*		R ₃	R ₄	Yield (%)*
6a	4-Ethylphenyl	H	32	6f	4-Methoxy phenyl	H	33
6b	Phenyl	Et	26	6g	2-Methoxy phenyl	H	27
6c	Benzyl	H	22	6h	2,4-Dimethoxy phenyl	H	29
6d	Allyl	H	20	6i	3-Nitrophenyl	H	25
6e	Benzyl	Me	13				

*Five-step overall yield from Wang resin (the Wang resin loading capacity is 1.1 mM/g)