

## BODIPY Appended Crown Ethers: Selective Fluorescence Changes for Hg<sup>2+</sup> Binding

Hyun Jung Kim, Sang Hoon Kim, Ja Hyung Kim, Eil-Hee Lee,<sup>†</sup> Kwang-Wook Kim,<sup>†</sup> and Jong Seung Kim<sup>\*</sup>

Department of Chemistry, Korea University, Seoul 136-701, Korea. \*E-mail: jongskim@korea.ac.kr

<sup>†</sup>Korea Atomic Energy Research Institute, Daejeon 305-600, Korea

Received July 1, 2008

**Key Words :** BODIPY, Crown ether, Mercury ion, PET, Fluorescence

Recently, the development of fluorescent chemosensors capable of selective recognition and sensing of metal ions is one of the most challenging fields from the vantage of organic and supramolecular chemistry.<sup>1,2</sup> The best effective fluorescence chemosensor must convert the event of metal ion recognition by the ionophore into light signals of the fluorophore with high sensitivity and ease of monitoring.<sup>3,4</sup> In designing sensors, the ionophore linked to the fluorophore should be preliminarily considered because they are responsible for the selectivity and binding efficiency of the whole chemosensors.

There is a growing interest in synthesizing macromolecule appended fluorescence materials showing marked changes upon cation, anion, and neutral molecule complexation, aiming to develop ion-selective and sensitive material. The "hard" ether-oxygen containing macrocycles show a binding preference toward "hard" alkali and alkaline earth metal cation, but the incorporation of "soft" sulfide or amine linkages shifts its preference toward "soft" heavy metal cations.<sup>5</sup> In addition, it has been demonstrated that macrocyclic ligands containing nitrogen-sulfur donor atoms can behave as highly selective complexing agents for transition metal cations.<sup>6,7</sup> In this regards, sulfur containing macrocycles such as thiacycrown ethers have been prepared and their complexation properties have been investigated with various metal cations.<sup>8-10</sup>


BODIPY (boradiazaindacenes) is well known fluorescent

dye in that it gives high quantum yields, large extinction coefficients and narrow emission bands. These properties facilitated their application in many fields, such as fluorescent labeling of biomolecules, ion sensing and signaling, energy transfer cassettes, light harvesting systems and fluorescent stains.<sup>11</sup>


Currently, considerable attention has been focused on fluorescent chemosensors for the selective and rapid determination of the toxic heavy metal ions, such as the Pb<sup>2+</sup>, Cd<sup>2+</sup>, and Hg<sup>2+</sup> ions.<sup>12</sup> Especially, in this regard, the Hg<sup>2+</sup> ion is considered highly dangerous because both elemental and ionic mercury can be converted into methyl mercury by bacteria in the environment, which subsequently bioaccumulates through the food chain.<sup>13</sup> Therefore, there is a high demand for the detection of the Hg<sup>2+</sup> ion both in environmental analysis and in industrial waste treatment.

In this paper, we report the synthesis and the fluorometric properties of BODIPY appended thiacycrown **1** and crown ethers **2** and **3**. Compound **1** displayed a highly selective chelation enhanced fluorescence (CHEF) effect only with Hg<sup>2+</sup>. Similar CHEF effect was observed upon the addition of Ag<sup>+</sup> to **1**, suggesting that silver ions also bind to the thiacycrown, but less strongly than does Hg<sup>2+</sup>.

Synthesis of **1-3** was achieved by adaptation of procedures reported earlier.<sup>14</sup> **4-6** were treated with 2,4-dimethylpyrrole in the presence of TFA, which were subsequently oxidized (*p*-chloranil), neutralized (Et<sub>3</sub>N), and treated with BF<sub>3</sub>·Et<sub>2</sub>O



**Scheme 1.** Synthetic route to fluorogenic ligands **1-3**. Reagents: (a) K<sub>2</sub>CO<sub>3</sub>/CH<sub>3</sub>CN; (b) 2,4-dimethylpyrrole/TFA/*p*-chloranil/Et<sub>3</sub>N/BF<sub>3</sub>·Et<sub>2</sub>O/CH<sub>2</sub>Cl<sub>2</sub>.




**Figure 1.** Absorption spectra of **1** (15.0  $\mu\text{M}$ ) with addition of  $\text{ClO}_4$  salts of  $\text{Li}^+$ ,  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Rb}^+$ ,  $\text{Cs}^+$ ,  $\text{Ag}^+$ ,  $\text{Cd}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Sr}^{2+}$ ,  $\text{Ba}^{2+}$ ,  $\text{Zn}^{2+}$ ,  $\text{Hg}^{2+}$ ,  $\text{Pb}^{2+}$ ,  $\text{Co}^{2+}$ ,  $\text{Cu}^{2+}$ , and  $\text{Al}^{3+}$  (50 equiv, respectively) in  $\text{H}_2\text{O}/\text{CH}_3\text{CN}$  (6:4, v/v).


to produce the desired BODIPY derivatives **1-3**, respectively.

Metal ion binding properties of **1** were investigated by monitoring fluorescence and UV/vis changes upon the addition of  $\text{Li}^+$ ,  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Rb}^+$ ,  $\text{Cs}^+$ ,  $\text{Ag}^+$ ,  $\text{Cd}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Sr}^{2+}$ ,  $\text{Ba}^{2+}$ ,  $\text{Zn}^{2+}$ ,  $\text{Hg}^{2+}$ ,  $\text{Pb}^{2+}$ ,  $\text{Co}^{2+}$ ,  $\text{Cu}^{2+}$ , and  $\text{Al}^{3+}$  ions in  $\text{H}_2\text{O}/\text{CH}_3\text{CN}$  (6:4, v/v). The aqueous media ( $\text{H}_2\text{O}/\text{CH}_3\text{CN}$  (6:4 = v/v)) we have used in this experiment reasoned that water media are all required when fluorescence molecules can be utilized in a biological system as a fluorescent chemosensor. Free **1** showed a sharp and strong absorption band at 498.0 nm (Figure 1). When  $\text{Hg}^{2+}$  is bound to **1**, the absorption band of **1** at 498 nm was slightly red-shifted to 499.5 nm ( $\Delta\lambda = 1.5$  nm). However, other ions did not cause any detectable changes with **1**. To get insight into the binding mode of **1** especially two sulfur atoms for the  $\text{Hg}^{2+}$  binding, corresponding crowns **2** and **3** have been prepared, too. Although the crown-5 (**2**) and crown-6 (**3**) loops are well known to adopt  $\text{Na}^+$  and  $\text{K}^+$  ion, respectively, the UV/Vis band shift could not be observed in this experiment. This is because the BODIPY is positioned in perpendicular to benzocrown unit, resulting in that the ICT change is no longer influenced upon metal ion binding.


The fluorescence change results are represented in Figure 2. **1** shows a pronounced selectivity and sensitivity for  $\text{Hg}^{2+}$  when it is irradiated at 499 nm, whereas **2** and **3** does less selective towards most metal cations. Similar fluorescence change was observed upon the addition of  $\text{Ag}^+$  to **1**, suggesting that the silver ions also bind to the sulfur atoms, but less strongly than does  $\text{Hg}^{2+}$  ion. The fluorescence enhancement phenomena of **1** upon  $\text{Hg}^{2+}$  binding are ascribable to the CHEF (chelation-induced enhanced fluorescence) mechanism.<sup>15</sup> When the  $\text{Hg}^{2+}$  ion interacts with sulfur atoms of the thiacycrown part, the PET (photo-induced electron transfer) from sulfur atoms to fluorescence BODIPY group is inhibited. With  $\text{Cu}^{2+}$  ions, however, a marked quenching effect was observed. Coordination of **1** to Cu (II) having  $d^9$ -electronic configuration is likely to induce the photo-induced electron transfer to give a considerable fluorescence quenching.



**Figure 2.** CHEF: fluorescence emission changes ( $I-I_0$ ) of 0.50  $\mu\text{M}$  solutions of **1-3** in  $\text{H}_2\text{O}/\text{CH}_3\text{CN}$  (6:4, v/v) upon addition of 50 equiv of various metal ions. Excitation at 498 nm;  $I_0$ : fluorescence emission intensity of free **1-3**;  $I$ : fluorescence emission intensity of metal complexes of **1-3**. (+) and (-) denote fluorescence increase and decrease, respectively.



**Figure 3.** Fluorescence emission spectra of **1** (0.50  $\mu\text{M}$ ) for  $\text{Hg}^{2+}$  ion titration in  $\text{H}_2\text{O}/\text{CH}_3\text{CN}$  (6:4, v/v). ( $\lambda_{\text{ex}} = 499$  nm).




**Figure 4.** Suggested complexation mechanism of **1** for  $\text{Hg}^{2+}$  ion.

Free **1** showed an intense greenish fluorescence at 507 nm, in  $\text{H}_2\text{O}/\text{CH}_3\text{CN}$  (6:4, v/v) solution ( $\lambda_{\text{ex}} = 498$  nm). With an excitation at 498 nm, the fluorescence intensity of **1** markedly increased in a function of  $[\text{Hg}^{2+}]$  (Figure 3). According to the extent of the fluorescence emission changes, we could obtain the association constants<sup>16</sup> of **1** ( $K_a = 1.59 \times 10^5 \text{ M}^{-1}$ ) for  $\text{Hg}^{2+}$  ion.

We also carried out Job's plot experiment by varying the concentration of both **1** and  $\text{Hg}^{2+}$ . The maximum point at the mole fraction of 0.5 indicates typical ligand mole fraction for 1:1 (ligand : metal) complexes (Figure 5).

An important feature of chemosensor has to show high selectivity toward specific analyte over other competitive



**Figure 5.** Job plot of a 1:1 complex of **1** and  $\text{Hg}^{2+}$  ion.

species. For the  $\text{Hg}^{2+}$  ion selectivity of **1**, competition experiments using miscellaneous cations including  $\text{Li}^+$ ,  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Rb}^+$ ,  $\text{Cs}^+$ ,  $\text{Ag}^+$ ,  $\text{Cd}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Sr}^{2+}$ ,  $\text{Ba}^{2+}$ ,  $\text{Zn}^{2+}$ ,  $\text{Pb}^{2+}$ ,  $\text{Co}^{2+}$ ,  $\text{Cu}^{2+}$ , and  $\text{Al}^{3+}$  in  $\text{H}_2\text{O}/\text{CH}_3\text{CN}$  (6:4, v/v) were carried out and their results are recorded in Figure 6. The miscellaneous competitive cations did not lead to any significant fluorescence changes. Moreover, in the presence of miscellaneous competitive cations, the  $\text{Hg}^{2+}$  ion still resulted in the similar fluorescence increase. In addition, the increases of fluorescence intensity resulting from the addition of the  $\text{Hg}^{2+}$  ion was not influenced by the subsequent addition of miscellaneous cations.

### Experimental Section


Precursors **5** and **6** were prepared by adaptation of reported procedures.<sup>17</sup>

**Preparation of 1.** This compound was prepared from **4** (300 mg, 1.1 mmol) according to a similar procedure outlined above for **3**. The compound was purified by flash column chromatography on silica gel using ethylacetate as eluent. Evaporation of the solvent yielded pale yellow vis-

cous oil which upon refrigeration gave the desired product as a pale yellow solid. Yield: 250 mg (53%).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  6.94 (d, 2H, ArH), 6.77 (d, 1H, ArH), 5.97 (s, 2H, ArH), 4.20 (m, 4H,  $-\text{OCH}_2\text{CH}_2\text{O}-$ ), 3.92 (m, 4H,  $-\text{OCH}_2\text{CH}_2\text{O}-$ ), 3.80 (m, 4H,  $-\text{OCH}_2\text{CH}_2\text{O}-$ ), 2.88 (s, 4H,  $-\text{SCH}_2\text{CH}_2\text{S}-$ ), 2.80 (q, 4H,  $-\text{SCH}_2\text{CH}_2\text{O}-$ ), 2.54 (s, 6H, -ArH), 1.46 (s, 6H, -ArH).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  146.9, 143.5, 143.2, 138.2, 129.0, 126.1, 125.6, 121.3, 119.4, 119.2, 72.6, 69.7, 46.4, 32.8, 29.9, 14.7, 8.8 ppm. FAB MS  $m/z$  ( $\text{M}^+$ ): calcd, 590.23 Found, 590.0.

**Preparation of 2.** This compound was prepared from **5** (300 mg, 1 mmol) according to a similar procedure outlined above for **3**. The compound was purified by flash column chromatography on silica gel using ethylacetate as eluent. Evaporation of the solvent yielded pale yellow viscous oil which upon refrigeration gave the desired product as a pale yellow solid. Yield: 300 mg (57%).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  6.95 (d, 2H, ArH), 6.79 (d, 1H, ArH), 5.97 (s, 2H, ArH), 4.20 (m, 4H,  $-\text{OCH}_2\text{CH}_2\text{O}-$ ), 3.96 (m, 4H,  $-\text{OCH}_2\text{CH}_2\text{O}-$ ), 3.79 (m, 4H,  $-\text{OCH}_2\text{CH}_2\text{O}-$ , 4H,  $-\text{OCH}_2\text{CH}_2\text{O}-$ ), 2.54 (s, 6H, -ArH), 1.46 (s, 6H, -ArH).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  155.7, 143.4, 130.0, 128.2, 121.3, 113.6, 70.9, 70.7, 70.0, 69.5, 68.9, 14.7 ppm. FAB MS  $m/z$  ( $\text{M}^+$ ): calcd, 514.25 Found, 514.0.

**Preparation of 3.** To a solution of 2,4-dimethylpyrrole (0.18 g, 1.89 mmol) and **6** (0.25 mg, 0.73 mmol) in dried dichloromethane, 2 drops of  $\text{CF}_3\text{CO}_2\text{H}$  was added. The yellow solution was stirred for 3 h at room temperature under  $\text{N}_2$ . A solution of *p*-chloranil (0.47 g, 1.89 mmol) in  $\text{CH}_2\text{Cl}_2$  (100 mL) was then added. After stirring for 30 min,  $\text{Et}_3\text{N}$  and  $\text{BF}_3\cdot\text{OEt}_2$  were subsequently added until a bright-green fluorescence was observed. The solution was washed with water, and the organic layer was dried over anhydrous  $\text{MgSO}_4$ . Removal of the organic solvent *in vacuo* afforded a reddish solid. Column chromatography on silica gel with EtOAc-hexane (2:1) as eluents gave 0.29 g (70%) of **3**.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  6.95 (d, 2H, ArH), 6.76 (d, 1H, ArH), 5.97 (s, 2H, ArH), 4.20 (m, 4H,  $-\text{OCH}_2\text{CH}_2\text{O}-$ ), 3.98 (m, 4H,  $-\text{OCH}_2\text{CH}_2\text{O}-$ ), 3.80 (m, 4H,  $-\text{OCH}_2\text{CH}_2\text{O}-$ ), 3.73



**Figure 6.** Fluorescence spectra of **1** (10  $\mu\text{M}$ ) in  $\text{H}_2\text{O}/\text{CH}_3\text{CN}$  (6:4, v/v) in the presence of the  $\text{Hg}^{2+}$  ion and miscellaneous cations including  $\text{Li}^+$ ,  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Rb}^+$ ,  $\text{Cs}^+$ ,  $\text{Ag}^+$ ,  $\text{Cd}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Sr}^{2+}$ ,  $\text{Ba}^{2+}$ ,  $\text{Zn}^{2+}$ ,  $\text{Pb}^{2+}$ ,  $\text{Co}^{2+}$ ,  $\text{Cu}^{2+}$ , and  $\text{Al}^{3+}$  (10 equiv, respectively; excitation wavelength at 498 nm). All spectral data were recorded at 10 min after  $\text{Hg}^{2+}$  addition.

(m, 4H, -OCH<sub>2</sub>CH<sub>2</sub>O-, 4H, -OCH<sub>2</sub>CH<sub>2</sub>O-), 2.54 (s, 6H, -ArH), 1.46 (s, 6H, -ArH). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 155.6, 143.4, 131.9, 131.8, 121.4, 121.3, 121.0, 70.4, 70.3, 70.2, 14.8, 14.7 ppm. FAB MS *m/z* (M<sup>+</sup>): calcd, 558.27 Found, 558.0.

**Preparation of 4.** A mixture of 3,4-dihydroxybenzaldehyde (1 g, 7.24 mmol) and K<sub>2</sub>CO<sub>3</sub> (1 g, 7.24 mmol) was taken in CH<sub>3</sub>CN (50 mL). Compound 7 (1.43 g, 7.23 mmol) was added to it with stirring and refluxed under nitrogen atmosphere for 18 hours. The cooled mixture was evaporated to dryness and the residue was extracted with CHCl<sub>3</sub> for 3 or 4 times. The extract was evaporated to yield a viscous oil which was purified by column chromatography on silica gel column using CHCl<sub>3</sub> (ethyl acetate/hexane, 1:2) as eluent. Yield: 2.1 g (77%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 9.83 (s, 1H, CHO), 7.46 (d, 2H, ArH), 6.9 (d, 1H, ArH), 4.36 (m, 4H, -OCH<sub>2</sub>CH<sub>2</sub>O-), 3.93 (m, 8H, -OCH<sub>2</sub>CH<sub>2</sub>O-), 3.86 (m, 4H, -OCH<sub>2</sub>CH<sub>2</sub>S-), 2.95 (s, 4H, -SCH<sub>2</sub>CH<sub>2</sub>S-), 1.22 (s, 4H, -OCH<sub>2</sub>CH<sub>2</sub>O-). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 190.9, 155.1, 150.2, 131.1, 127.4, 447.9, 72.8, 72.2, 71.9, 68.9, 32.9, 32.0, 29.9 ppm. FAB MS *m/z* (M<sup>+</sup>): calcd, 372.11 Found, 372.0.

**Preparation of 7.** Under nitrogen, a solution of 1,2-ethanedithiol (0.5 g, 5.3 mmol), and NaOH (0.44 g, 10.6 mmol) in ethanol (100 mL) was stirred at room temperature. Then 2-(2-chloroethoxy)ethanol (1.38 g, 10.6 mmol) was added to it with stirring and refluxed under N<sub>2</sub> atmosphere for 2 hours. After stirring for 24 h, the solvent was evaporated *in vacuo*. The resulting brownish oil was used directly for the next reaction. The resulting residue treated with tosyl chloride (1.47 g, 7.4 mmol) and NaOH (0.3 g, 7.7 mmol) in THF (100 mL). After stirring at room temperature for 24 h, the solvent was evaporated *in vacuo*, and the mixture was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (100 mL). The organic layer was washed with water (300 mL) and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and filtered. Purification by column chromatography on silica gel (ethyl acetate/hexane, 1:4) provided 1.5 g of yellowish oil 7 in 50% yield. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.74 (d, 4H, ArH), 7.30 (d, 4H, ArH), 4.06 (m, 4H, -OCH<sub>2</sub>CH<sub>2</sub>O-), 3.59 (m, 4H, -OCH<sub>2</sub>CH<sub>2</sub>O-), 3.51 (m, 4H, -OCH<sub>2</sub>CH<sub>2</sub>O-), 2.66 (s, 4H, -SCH<sub>2</sub>CH<sub>2</sub>S-), 2.58 (m, 4H, -SCH<sub>2</sub>CH<sub>2</sub>O-), 2.37 (s, 6H, -OCH<sub>2</sub>CH<sub>2</sub>O-). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 145.1, 133.0, 130.1, 128.1, 71.3, 69.4, 68.5, 38.7, 37.7, 32.8, 32.1, 31.6, 21.8 ppm. FAB MS *m/z* (M<sup>+</sup>): calcd, 578.11 Found, 578.0.

**Acknowledgments.** The authors wish to acknowledge the financial support from the basic science research grant of KRF 2008-C00426 and SRC program (R11-2005-008-02001-0(2008)).

## References

- (a) *Chemosensors for Ion and Molecule Recognition. NATO ASI Series*; Desvergnès, J. P.; Czarnik, A. W., Eds; Kluwer Academic: Dordrecht, 1997. (b) de Silva, A. P.; Gunaratne, H. Q. N.; Gunnlaugsson, T.; Huxley, A. J. M.; McCoy, C. P.; Rademacher, J. T.; Rice, T. E. *Chem. Rev.* **1997**, *97*, 1515. (c) Schmidtchen, F. P.; Berger, M. *Chem. Rev.* **1997**, *97*, 1609. (d) Kim, J. S.; Quang, D. T. *Chem. Rev.* **2007**, *107*, 3780.
- (a) Gale, P. A. *Coord. Chem. Rev.* **2000**, *199*, 181. (b) Fabrizzi, L. *Coord. Chem. Rev.* **2000**, *205*, 1. (c) Kuswandi, B. *Jurnal ILMU DASRA* **2000**, *1*, 18. (d) Bren, V. A. *Russ. Chem. Rev.* **2001**, *70*, 1017. (e) Valeur, B. *Molecular Fluorescence*; Wiley-VCH: Weinheim, 2002. (f) Diamond, D.; Nolan, K. *Anal. Chem.* **2001**, *73*, 22A. (g) Lee, Y. O.; Choi, Y. H.; Kim, J. S. *Bull. Kor. Chem. Soc.* **2007**, *28*(1), 151. (h) Kim, H. J.; Kim, S. H.; Quang, D. T.; Kim, J. H.; Suh, I. W.; Kim, J. S. *Bull. Kor. Chem. Soc.* **2007**, *28*(5), 811.
- (a) Beer, P. D.; Gale, P. A. *Angew. Chem. Int. Ed. Engl.* **2001**, *40*, 486. (b) Ludwig, R.; Dzung, N. T. K. *Sensors* **2002**, *2*, 397. (c) Callan, J. F.; de Silva, A. P.; Magri, D. C. *Tetrahedron* **2005**, *61*, 8551. (d) Dallali, N.; Darabi, A.; Agrawal, Y. K. *Rev. Anal. Chem.* **2005**, *24*, 263. (e) Amendola, V.; Fabrizzi, I.; Forti, F.; Pallavicini, P.; Poggi, A.; Sacchi, D.; Tagleitti, A. *Coord. Chem. Rev.* **2006**, *250*, 273. (f) Mancin, F.; Rampazzo, E.; Tecilla, P.; Tonellato, U. *Chem. Eur. J.* **2006**, *12*, 1844. (g) Kungwandi, B.; Nuriman; Verboom, W.; Reinhoudt, D. N. *Sensors* **2006**, *6*, 978. (h) Zhu, L.; Anslyn, E. V. *Angew. Chem., Int. Ed.* **2006**, *45*, 1190. (i) Valeur, B.; Leray, I. *Inorg. Chem. Acta* **2007**, *360*, 765. (j) Anslyn, E. V. *J. Org. Chem.* **2007**, *72*, 687.
- Löhr, H.-G.; Vögtle, F. *Acc. Chem. Res.* **1985**, *18*, 65.
- Chartres, J. D.; Davies, M. S.; Lindoy, L. F.; Meehan, G. V.; Wei, G. *Inorg. Chem. Commun.* **2006**, *9*, 751.
- Ikeda, K.; Abe, S. *Anal. Chim. Acta* **1998**, *363*, 165.
- Matsushita, M. T.; Shono, T. *Polyhedron* **1984**, *3*, 1357.
- Groth, A. M.; Lindoy, L. F.; Meehan, G. V. *J. Chem. Soc., Perkin Trans. 1* **1996**, 1553.
- Lindoy, L. F.; Baldwin, D. S. *Pure Appl. Chem.* **1989**, *61*, 909.
- Tanaka, M.; Nakamura, M.; Ikeda, T.; Ikeda, K.; Ando, H.; Shibutani, Y.; Yajima, S.; Kimura, K. *J. Org. Chem.* **2001**, *66*, 7008.
- (a) Beer, G.; Rurack, K.; Daub, J. *Chem. Commun.* **2001**, 1138. (b) Moon, Y. S.; Cha, N. R.; Kim, Y. H.; Chang, S. K. *J. Org. Chem.* **2004**, *69*, 181. (c) Gareis, T.; Huber, C.; Wolfbeis, O. S.; Daub, J. *Chem. Commun.* **1997**, 1717. (d) Rurack, K.; Kollmannsberger, M.; Daub, J. *New J. Chem.* **2001**, *25*, 289. (e) DiCesare, N.; Lakowicz, J. R. *Tetrahedron Lett.* **2001**, *42*, 9105. (f) Baki, C. N.; Akkaya, E. U. *J. Org. Chem.* **2001**, *66*, 1512. (g) Turfan, B.; Akkaya, E. U. *Org. Lett.* **2002**, *4*, 2857.
- (a) Kim, S. K.; Lee, S. H.; Lee, J. Y.; Lee, J. Y.; Bartsch, R. A.; Kim, J. S. *J. Am. Chem. Soc.* **2004**, *126*, 16499. (b) Kim, S. K.; Bok, J. H.; Bartsch, R. A.; Lee, J. Y.; Kim, J. S. *Org. Lett.* **2005**, *7*, 4839. (c) Kim, S. K.; Kim, S. H.; Kim, H. J.; Lee, S. H.; Lee, S. W.; Ko, J.; Bartsch, R. A.; Kim, J. S. *Inorg. Chem.* **2005**, *44*, 7866. (d) Kim, H. J.; Park, S. Y.; Yoon, S.; Kim, J. S. *Tetrahedron* **2008**, *64*, 1294. (e) Lee, M. H.; Kim, H. J.; Yoon, S.; Park, N.; Kim, J. S. *Org. Lett.* **2008**, *10*, 213. (f) Lee, M. H.; Wu, J.-S.; Lee, J. W.; Jung, J. H.; Kim, J. S. *Org. Lett.* **2007**, *9*, 2501. (g) Wu, J.-S.; Hwang, I.-C.; Kim, K. S.; Kim, J. S. *Org. Lett.* **2007**, *9*, 907.
- (a) Wolfe, M. F.; Schwarzbach, S.; Sulainman, R. A. *Environ. Toxicol. Chem.* **1998**, *17*, 146. (b) Renzoni, A.; Zino, F.; Franchi, E. *Environ. Res.* **1998**, *77*, 68. (c) Feng, X. B.; Tang, S. L.; Li, Z. G.; Wang, S. F.; Liang, L. *Chin. Sci. Bull.* **2004**, *49*, 2068. (d) Chen, C. Y.; Zhang, P. Q.; Chai, Z. F. *Chin. Sci. Bull.* **2005**, *50*, 113. (e) Wang, N.; Zhu, Y. M.; Sheng, L. X.; Meng, D. *Chin. Sci. Bull.* **2005**, *50*, 2166. (f) Harris, H. H.; Pickering, I. J.; George, G. N. *Science* **2003**, *301*, 1203.
- Kim, H. J.; Kim, J. S. *Tetrahedron Lett.* **2006**, *47*, 7051.
- Lee, H. N.; Kim, H. N.; Swamy, K. M. K.; Park, M. S.; Kim, J.; Lee, H.; Lee, K.-H.; Park, S.; Yoon, J. *Tetrahedron Lett.* **2008**, *49*, 1261.
- Association constants were obtained using the computer program ENZFITTER, available from Elsevier-BIOSOFT, 68 Hills Road, Cambridge CB2 1LA, United Kingdom; Connors, K. A. *Binding Constants*; Wiley: New York, 1987.
- (a) Sousa, C.; Freire, C.; de Castro, B. *Molecules* **2003**, *8*, 894. (b) Reinhoudt, D. N.; de Jong, F.; Tomassen, H. P. M. *Tetrahedron Lett.* **1979**, *20*, 2067.