

Quantitative and Pattern Recognition Analyses for the Quality Evaluation of *Cimicifugae Rhizoma* by HPLC

Zhe Fang, Dong Cheul Moon,[†] Kun Ho Son,[‡] Jong Keun Son,[§] Byung Sun Min, and Mi Hee Woo*

College of Pharmacy, Catholic University of Daegu, Gyeongsan 712-702, Korea. *E-mail: woomh@cu.ac.kr

[†]College of Pharmacy, Chungbuk National University, Cheongju 361-763, Korea

[‡]College of Life Science, Andong National University, Andong 760-740, Korea

[§]College of Pharmacy, Yeungnam University, Gyeongsan 712-749, Korea

Received July 6, 2010, Accepted November 15, 2010

In this study, quantitative and pattern recognition analysis for the quality evaluation of *Cimicifugae Rhizoma* using HPLC/UV was developed. For quantitative analysis, three major bioactive phenolic compounds were determined. The separation conditions employed for HPLC/UV were optimized using ODS C₁₈ column (250 × 4.6 mm, 5 μM) with isocratic elution of acetonitrile and water with 0.1% phosphoric acid as the mobile phase at a flow rate of 1.0 mL/min and a detection wavelength of 323 nm. These methods were fully validated with respect to the linearity, accuracy, precision, recovery, and robustness. The HPLC/UV method was applied successfully to the quantification of three major compounds in the extract of *Cimicifugae Rhizoma*. The HPLC analytical method for pattern recognition analysis was validated by repeated analysis of twelve reference samples corresponding to five different species of *Cimicifugae Rhizoma* and seventeen samples purchased from markets. The results indicate that the established HPLC/UV method is suitable for the quantitative analysis and quality control of multi-components in *Cimicifugae Rhizoma*.

Key Words: *Cimicifugae Rhizoma*, HPLC, Pattern recognition, Quality control

Introduction

Herbal medicines have a long history in therapeutic field and they are attracting considerable attention because of low toxicity and excellent therapeutic benefit. Quality control for herbal drugs is difficult than for synthetic drugs because of the chemical complexity of the ingredients. In herbal drugs, no single active constituent is responsible for the overall pharmacological efficacy. For quality control of complex systems, the determination of only a few compounds cannot give a comprehensive and accurate assessment of all active compounds in herbal drugs. Pattern recognition analysis can provide the information of overall chemical composition of herbal medicines traditionally used for quality control.¹

Cimicifugae Rhizoma is the root of *Cimicifuga heracleifolia* Komarov or the other species in the Korean Pharmacopeia (K.P.), of *C. simplex* Wormskjold, *C. dahurica* Maximowicz, *C. foetida* Linne, or *C. heracleifolia* Komarov in the Japanese Pharmacopoeia (J.P.), and of *C. heracleifolia* Komarov, *C. dahurica* Maxim., or *C. foetida* L. in the Chinese Pharmacopoeia (C.P.). *Cimicifugae Rhizoma* is controlled to contain not less than 0.1% of isoferulic acid in C.P. There are ten different *Cimicifuga* species: *C. heracleifolia*, *C. dahurica*, *C. foetida*, *C. simplex*, *C. japonica*, *C. acerina*, *C. biternat*, *C. racemosa*, *C. americana* and *C. rubifolia*. We collected five different species of *Cimicifugae Rhizoma* for this study from Korea and China. Five different species included *C. heracleifolia*, *C. dahurica*, *C. foetida* and *C. simplex*, which have been used in K.P., J.P. and C.P., and *C. japonica*. Pharmacological studies have revealed in to have bone resorption inhibition,² vasoactive effect,³ estrogenic activity,⁴ antiinflammatory,⁵ cytotoxicity,⁶ anti-microbial and cytokine modulation effect⁷ and anticomplement activity.⁸

Some HPLC/UV analytical methods have been developed for the analysis of *Cimicifugae Rhizoma* and its related products.⁹⁻¹¹ Kan He *et al.* reported the triterpene glycoside cimigenol 3-O-arabinoside, cimifugin and cimifugin-3-O-glucoside as specific markers for the distinction of *C. racemosa* from the other *Cimicifuga* species.¹¹ Black cohosh, one of the most important herbal products in the US dietary supplements market, is manufactured from roots and rhizomes of *C. racemosa*. *Cimicifugae Rhizoma* samples used in this study didn't include *C. racemosa* which is native to the Eastern United States, growing as far south as Florida. Furthermore, these studies were focused only quantitative analysis of selected marker compounds which are not promising approaches for the quality control of finger printing analysis of herbal drugs. In the present study, a simple, sensitive and precise reverse-phase HPLC/UV method has been developed for the quantitative determination of three marker phenolic components, caffeic acid (1), ferulic acid (2) and isoferulic acid (3) along with pattern-recognition method for the quality control of *Cimicifugae Rhizoma* extract. The twelve *Cimicifugae Rhizoma* authentic samples collected from China and Korea and seventeen *Cimicifugae Rhizoma* samples purchased from the markets were analyzed by HPLC after extraction with 50% ethanol. *Cimicifugae Rhizoma* samples exhibited very different triterpenoid chromatogram patterns in HPLC according to their different *Cimicifuga* species. Quantitative analysis of isoferulic acid single compound in C.P. would not be an adequate approach for quality control of *Cimicifugae Rhizoma*. Therefore in pattern recognition analysis we used three phenolic marker compounds as caffeic acid (1), ferulic acid (2) and isoferulic acid (3) instead of triterpenoids. In pattern analysis with multivariate statistical analysis we used R-2.11.0 program (downloaded from web <http://www.r-project.org>) to analyze eleven authentic samples of *Cimicifugae Rhizoma*.

and seventeen commercial ones. Subsequent pattern analysis was applied to assess the comprehensive quality of *Cimicifugae Rhizoma*.

Experimental Section

Plant Material. *Cimicifugae Rhizoma* samples collected in 2006 for this study include the following accessions: a *C. foetida* (c8) collected from China, and four *C. heracleifolia* (c1~c4), two *C. dahurica* (c5 and c6), two *C. foetida* (c7 and c9), two *C. simplex* (c10 and c11) and a *C. japonica* (c12) collected from Korea, and seventeen commercial *Cimicifugae Rhizoma* samples (c13 and c14 from Yangyeongsi, Daegu, Korea; c15~c19 from Gyongdong market, Korea; c20~c24, c26 and c27 from Yeongcheon market, Korea; c25 from Xining, China; c28, c29 from Seoul market, Korea) purchased from markets of Korea and China.

Reagents. All of the standard compounds were provided by Prof. Kun Ho Son, Andong National University, Andong, Korea. Their structures were unambiguously identified by NMR and MS data, with the published data, such as caffeic acid,¹² ferulic acid¹³ and isoferulic acid.¹⁴ The standard compound structures were shown in Fig. 1. Purity of standard compounds was estimated to be higher than 95% based on HPLC and LC-MS/MS analysis. Internal standard, 4-hydroxycinnamic acid (**4**), was purchased from Sigma Chemicals (St. Louis, MO, USA). Methanol and acetonitrile of HPLC grade were purchased from Merck K GaA (Darmstadt, Germany). All other chemicals used were of analytical grade unless otherwise noted. Distilled water was prepared using Milli-Q purification system (Millipore, Bedford, MA, USA).

Sample Preparation. To determine the content of three marker compounds and pattern recognition analysis of *Cimicifugae Rhizoma* samples, the dried rhizome powder were used for each extraction. *Cimicifugae Rhizoma* samples were powdered and sieved through 50 mesh, and about 0.3 g of the powder were accurately weighed and added 50 mL of 50% ethanol, accurately measured weight and refluxed for 4 hours at 80 °C. The solution was cooled, weighed again, and made up the loss in weight with 50% ethanol. The solution was filtered through a 0.45 µm membrane filter and the filtrate was used as the test solution. Sample solution of 10 µL was subjected to injection into the HPLC system.

HPLC/UV Condition. The HPLC equipment was a Waters HPLC system (Waters, Milford, MA, USA) with Waters 600 pumps, a Waters 486 UV detector and a Waters 717 autosampler. YMC ODS-H80 (250 × 4.6 mm, 4 µm), Shiseido capcell pak (250 × 4.6 mm, 5 µm) and Shodex ODS pak (250 × 4.6 mm, 5 µm) columns were tested with the guard columns filled with the same stationary phase. A (100% acetonitrile) and B (0.1% phosphoric acid in water) were used as the mobile phase using a isocratic condition (A : B = 15 : 85, v/v %) to analyze samples. The mobile phase was filtered under vacuum through a 0.45 µm membrane filter and degassed prior to use. The analysis was carried out at a flow rate of 1.0 mL/min with the detection wavelength set to 323 nm, and the total run time was 40 min. All compounds could be resolved with baseline separation at 323 nm with the maximum absorption. Hence, characteristic chromatographic patterns were obtained at 323 nm. The chromatograms were processed using software Empower pro software, build 1154 (Waters, Milford, MA).




Figure 1. Chemical structure of marker compounds caffeic acid (A), ferulic acid (B) and isoferulic acid (C) along with UV spectra.

graphic patterns were obtained at 323 nm. The chromatograms were processed using software Empower pro software, build 1154 (Waters, Milford, MA).

Analytical Method Validation. The standards (4 mg) of caffeic acid, ferulic acid and isoferulic acid were each accurately weighed and then dissolved with 10 mL of 100% methanol to produce stock standard solutions of 400 ppm, respectively. The internal standard (4-hydroxycinnamic acid) of 500 mg was accurately weighed and then dissolved with 100 mL of 100% methanol to produce stock solution of 5,000 ppm. The calibration curves were made by diluting the stock solutions with 100% methanol. The reference solution of the three phenolic compounds at concentrations of 0.2 ~ 40 µg/mL was analyzed by HPLC/UV. The regression equations were calculated in the form of $y = ax + b$, where y and x correspond to peak area ratio for internal standard and compound concentration, respectively.

The recovery tests were executed by mixing a powdered sample (0.3 g) with three control levels (20%, 50%, and 100% concentrations of the each compound contained in the samples) of the reference compounds. The mixture was then extracted by reflux with 50 mL of 50% ethanol at 80 °C for 4 hours. The extract solution was filtered through a 0.45 µm membrane. The HPLC/UV analysis experiments were performed in triplicate for each control level. The data was compared with those from the standard solution and extracted sample. Precision and accuracy were determined by multiple analysis ($n = 5$) of quality control samples prepared at lower, medium and higher concentrations spanning the calibration range (0.2, 20, and 40 µg/mL).

Pattern Recognition Analysis. The twelve authentic samples

of *Cimicifugae Rhizoma* were chosen as references for the quality control of *Cimicifugae Rhizoma*. To evaluate the phytochemical equivalency among the twenty-nine samples corresponding to twelve authentic and seventeen commercial ones, pattern recognition analysis was conducted. In this study we used three marker compound peaks (caffeic acid (1), ferulic acid (2) and isoferulic acid (3)) for pattern recognition analysis. Pattern recognition analysis was conducted using software package R-2.11.0.

Results and Discussion

Optimization of Chromatographic Condition. The HPLC conditions were selected by the requirement for obtaining the chromatograms with a better resolution of the adjacent peaks within a short retention time. For the optimization of chromatographic condition, the effect of the composition of mobile phase on the separation was examined. Mobile phase of water-methanol did not result in the satisfactory separation of structurally similar compounds. Acetonitrile as an organic modifier demonstrated a significant improvement on separation. We had tested the addition of 0.1%, 1% and 10% acid (acetic acid, formic acid and phosphoric acid) to the mobile phase to do experiment. The addition of 0.1% phosphoric acid to the mobile phase to all of the compounds resulted in a good resolution, as well as satisfactory peak symmetry and shape. The typical chromatograms of samples and standard mixture are shown in Fig. 2, from which one can observe that all target compounds and an internal standard are completely separated within 40 minutes. 4-Hydroxycinnamic acid (4) was selected as an internal standard. The chromatographic peaks of the analytes in sample solution were identified by comparing their retention times with those of the reference standards and further confirmed by spiking samples with the reference compounds (Fig. 2). For the choice of detection wavelength, extract sample was scanned between 200 - 400 nm using PDA detector. All compounds

could be resolved with baseline separation at 323 nm with the maximum absorption shown for three major constituents (Fig. 1). Hence, characteristic chromatographic patterns were obtained at 323 nm.

Optimization of Sample Preparation Condition. Nine extracting solvents, 100% ethanol, 75% ethanol, 50% ethanol, 25% ethanol, 100% H₂O, 100% methanol, 75% methanol, 50% methanol and 25% methanol were compared with regard to sample assays using reflux extraction for 4 hours at 80 °C. When sample was extracted with 50% ethanol, the sample assay was higher than the other solvent samples. Therefore, we employed 50% ethanol as an extracting solvent throughout this work. Two extraction methods, ultra-sonication and reflux using 50% ethanol extraction solvent, were compared with regard to sample assays. When used for reflux extraction method, the sample assay was higher than sonication one. To determine the time needed to obtain complete extractions, extractions of a sample were performed for six different lengths of time (1, 2, 3, 4, 5 and 6 hours). The rest of the variables employed were: 50% ethanol solvent and reflux extraction method at 80 °C. When extraction time was 4 hours, the sample assay was same as 5 and 6 hours, and higher than 1, 2 and 3 hours. Therefore, when extraction time was 4 hours, all of the compounds were sufficiently extracted.

Validation. Each coefficient of correlation (r^2) was > 0.999, as determined by least square analysis, suggesting good linearity between the peak area ratio and the compound concentrations (Table 1). The limits of detection (LOD) and limits of quantitation (LOQ) were evaluated based on the lowest detectable peak in the chromatogram having a signal-to-noise (S/N) ratio of 3 and 10, respectively. Under our experimental conditions, we listed LOD and LOQ in Table 1. The obtained values for both LOD and LOQ for these three standards were shown to be low enough to detect traces of these compounds in either crude extract or its preparation.

The extraction recovery test was performed by extracting a




Figure 2. HPLC chromatograms of standard mixture and five different species of *Cimicifugae Rhizoma*. 1 caffeic acid, 2 ferulic acid, 3 isoferulic acid, 4 4-hydroxycinnamic acid (I.S.).

Table 1. Calibration graphs, linear ranges, LOD and LOQ

Analytes	linear range (μg/mL)	Slope (a)	Intercept (b)	Correlation coefficient (r)	LOD (ng/mL)	LOQ (ng/mL)
Caffeic acid (1)	0.2 ~ 40	0.2710	0.0105	0.9999	12	36
Ferulic acid (2)	0.2 ~ 40	0.2680	0.0190	0.9997	13	39
Isoferulic acid (3)	0.2 ~ 40	0.2494	0.0277	0.9999	15	45

Table 2. Recovery of marker compounds through standard addition (n = 5)

Analyte	Fortified conc. (μg/mL)	Observed conc. (μg/mL)	Recovery mean (%)	Recovery cv (%)
Caffeic acid (1)	0	3.03	-	-
	1.5	4.57	102.88	0.17
	3.0	5.98	98.39	0.11
	4.5	7.30	94.97	0.07
Ferulic acid (2)	0	4.11	-	-
	2.0	6.14	101.63	0.07
	4.0	8.00	97.18	0.12
	6.0	9.88	96.23	0.10
Isoferulic acid (3)	0	18.70	-	-
	4.0	22.58	97.05	0.35
	8.0	26.95	103.14	0.41
	16.0	34.66	99.77	0.44

Table 3. Precision and accuracy of analytical results

Analyte	Nominal conc. (μg/mL)	Intraday (n = 5)				Inter-day (n = 5)			
		Observed (μg/mL)	SD	Accuracy (%)	Precision (%)	Observed (μg/mL)	SD	Accuracy (%)	Precision (%)
Caffeic acid (1)	0.20	0.19	1.78	96.29	1.85	0.19	0.89	96.04	0.93
	20.00	20.69	0.97	103.43	4.34	20.92	2.00	104.60	2.09
	40.00	41.07	2.48	102.68	2.45	40.69	1.53	101.72	1.51
Ferulic acid (2)	0.20	0.21	2.44	103.03	2.37	0.19	1.96	97.14	2.03
	20.00	19.53	1.37	97.63	1.41	19.59	1.64	97.94	1.68
	40.00	40.95	1.00	102.38	0.98	41.22	1.52	103.05	1.48
Isoferulic acid (3)	0.20	0.19	2.32	98.57	2.35	0.19	1.50	98.33	1.53
	20.00	19.25	3.60	96.23	3.74	19.10	2.94	95.48	3.09
	40.00	42.75	2.71	106.88	2.54	42.60	2.70	106.49	2.53

known amount of the three compounds from the *Cimicifugae Rhizoma* powder samples. A known amount of each standard compound at three levels was mixed with the sample powder and extracted with 50% ethanol, as described in the experimental section. The % recovery of each standard ranged from 94.97 to 103.14%, and the RSD was less than 0.44% (Table 2). The average recovery was represented by the formula: $R(\%) = [(\text{amount from the sample spiked standard} - \text{amount from the sample}) / \text{amount from the spiked standard}] \times 100$. Precision and accuracy were determined by multiple analysis (n = 5) of quality control samples prepared at lower, medium and higher concentrations spanning the calibration range. Intra-assay precision and accuracy were determined from the variability of multiple

analyses (n = 5) of quality control samples analyzed within the same analytical run. The remaining quality control samples had the intra-assay precision below 4.34% and accuracy between 96.23% and 106.88%. Inter-assay precision and accuracy were evaluated from the variability of multiple analyses (n = 5) of quality control samples analyzed on single analytical run and extended for consecutive five days. The remaining quality control samples had the inter-assay precision lower than 3.09% and accuracy between 95.48% and 106.49%. The above data reflects that the developed method is highly reproducible and precision and accuracy data are presented in Table 3.

The robustness was determined in order to evaluate the reliability of the established HPLC methods. All of the parameters

Table 4. Robustness of marker compounds through standard addition

		Analytes					
		caffeic acid		ferulic acid		isoferulic acid	
		mean	SD	mean	SD	mean	SD
Column	Theoretical plate (N)						
	YMC	1121	128	4146	143	7214	209
	Shiseido	1001	48	2927	16	2677	125
	Sodex	543	27	1834	98	2243	62
	Capacity factor (k')						
	YMC	3.62	0.04	8.88	0.10	10.39	0.12
	Shiseido	3.43	0.16	7.56	0.29	8.92	0.35
	Sodex	4.39	0.02	10.17	0.03	11.84	0.03
	Separation factor (α)						
	YMC	0.48	0.01	1.17	0.02	1.16	0.00
	Shiseido	0.45	0.02	1.00	0.03	1.18	0.00
	Sodex	0.58	0.00	1.34	0.01	1.16	0.00
	Resolution (Rs)						
	YMC	16.47	0.85	4.32	0.45	2.89	0.03
	Shiseido	16.24	1.16	3.23	0.79	2.17	0.03
	Sodex	7.66	0.35	3.35	0.43	2.89	0.03
Temperature	Theoretical plate (N)						
	25	1122	129	4146	143	7214	210
	30	1183	109	4226	138	9217	597
	35	1015	42	4183	129	4450	239
	40	1003	49	3844	193	4540	64
	Capacity factor (k')						
	25	3.75	0.04	9.20	0.12	10.80	0.14
	30	3.61	0.04	8.88	0.10	10.39	0.12
	35	3.51	0.01	8.56	0.02	9.99	0.02
	40	3.40	0.07	8.27	0.18	9.60	0.22
	Separation factor (α)						
	25	0.49	0.00	1.21	0.01	1.17	0.00
	30	0.48	0.01	1.17	0.02	1.16	0.00
	35	0.46	0.00	1.13	0.00	1.17	0.00
	40	0.45	0.01	1.09	0.02	1.16	0.00
	Resolution (Rs)						
	25	16.52	0.95	4.32	0.44	2.90	0.03
	30	15.81	0.86	5.26	0.20	3.14	0.03
	35	16.43	0.27	3.35	0.09	2.53	0.03
	40	15.28	0.72	2.32	0.52	2.42	0.04
Flow rate	Theoretical plate (N)						
	0.9	1125	119	4166	143	7315	410
	1.0	1120	125	4124	145	7244	410
	1.1	1116	105	4093	139	7187	439
	Capacity factor (k')						
	0.9	3.72	0.04	9.11	0.12	10.80	0.15
	1.0	3.59	0.05	8.90	0.13	10.59	0.14
	1.1	3.50	0.03	8.76	0.12	10.39	0.12
	Separation factor (α)						
	0.9	0.49	0.03	1.21	0.01	1.25	0.02
	1.0	0.47	0.02	1.16	0.02	1.19	0.01
	1.1	0.45	0.02	1.13	0.02	1.16	0.01
	Resolution (Rs)						
	0.9	16.88	0.72	4.42	0.52	3.11	0.04
	1.0	16.43	0.75	4.22	0.40	2.99	0.03
	1.1	15.91	0.86	4.16	0.20	2.74	0.03

were maintained so there would not be any interference with the other peaks for the *Cimicifugae Rhizoma*. The experimental conditions, such as the column temperature, column species and mobile phases, were purposely altered. The theoretical plate (N), capacity factor (k'), separation factor (α) and resolution (Rs) were evaluated. To evaluate the suitability three different columns, YMC, Shiseido and Shodex, were compared with regard to four analytical factors (N, k' , α and Rs) on the column temperature of 30 °C. The result showed that four analytical factors did not differ greatly, depending on the column species (YMC, Shiseido and Shodex). Four different column temperatures, 25, 30, 35 and 40 °C, were compared with regard to four analytical factors using YMC column. The result showed that four analytical factors did not differ greatly, depending on the column temperature (25, 30, 35 and 40 °C). Three different flow rates, 0.9, 1.0 and 1.1 mL/min, were compared with regard to four analytical factors using YMC column on 25 °C. The result showed that four analytical factors did not differ greatly, depending on the flow rates (0.9, 1.0 and 1.1). We optimized by changing the chromatographic parameters, but the four analytical factors did not differ greatly, therefore this experiment condition were sufficiently robust.

The sample stability test was determined with a standard mixture solution at 0, 0.5, 1, 2, 5, 10, 15 and 30 days. During this period, the solution was stored at no light conditions, and at room temperature and 4 °C, respectively. The resulting data indicated that all marker analytes remained stable more than 98% during the experimental period.

Sample Analysis. The developed HPLC/UV method was then applied to the simultaneous determination of the three compounds, caffeic acid (1), ferulic acid (2) and isoferulic acid (3) in the *Cimicifugae Rhizoma*. Twelve authentic *Cimicifugae Rhizoma* samples corresponding to five different *Cimicifugae Rhizoma* species and seventeen commercially available *Cimicifugae Rhizoma* samples were obtained from Korea and China. The developed analytical method was subsequently applied to the simultaneous determination of the three components in *Cimicifugae Rhizoma* extract. The quantity of each compound present in samples was determined and the results are summarized in Table 5. Each sample was analyzed in triplicate to ensure the reproducibility of the quantitative result. The results indicated that, the contents (mg/g) of caffeic acid showed 1.5 ~ 3.5 (mg/g) in *C. foetida* and *C. simplex* and 0.1 ~ 1.0 (mg/g) in *C. heracleifolia*, *C. dahurica*, *C. foetida* (mutation) and *C. japonica*. In the commercial samples, the assays of caffeic acid showed 0.1 ~ 0.4 (mg/g) except c25 (0.9 mg/g), corresponding to lower quantities than *C. foetida* and *C. simplex*. The contents of isoferulic acid showed 0.001 ~ 0.3 (mg/g) in *C. foetida*, *C. simplex* and *C. japonica*, and 1.6 ~ 4.6 (mg/g) in *C. heracleifolia*, *C. dahurica* and *C. foetida* (mutation). In the commercial samples, the contents of isoferulic acid showed 1.6 ~ 4.3 (mg/g), nearly the same as *C. heracleifolia*, *C. dahurica* and *C. foetida* (mutation). Additionally, the contents of ferulic acid showed 0.2 ~ 0.7 (mg/g) in all of *Cimicifugae Rhizoma* species and commercial samples except c12 (0.04 mg/g) and c25 (0.08 mg/g). Consequently, *C. foetida* and *C. simplex* species samples were higher than *C. heracleifolia*, *C. dahurica*, *C. foetida* (mutation) and *C. japonica* species samples in the caffeic acid assays, whereas


Table 5. Contents (mg/g) of phenolic components in Cimicifugae Rhizoma (n = 3)

Sample	Contents (mg/g)								
	Caffeic acid			Ferulic acid			Isoferulic acid		
	Mean	SD	RSD	mean	SD	RSD	mean	SD	RSD
c1	0.118	0.002	1.99	0.516	0.013	2.61	1.575	0.023	1.48
c2	0.149	0.001	0.13	0.658	0.002	0.28	2.237	0.007	0.30
c3	0.611	0.004	0.64	0.233	0.001	0.10	5.243	0.003	0.63
c4	1.018	0.015	1.50	0.373	0.004	1.08	2.520	0.039	1.56
c5	0.333	0.003	0.94	0.490	0.002	0.49	4.550	0.044	0.96
c6	0.592	0.007	1.13	0.305	0.001	1.13	3.457	0.006	0.18
c7	1.537	0.015	0.97	0.376	0.006	0.15	0.210	0.003	1.53
c8	3.454	0.028	0.80	0.320	0.002	0.56	0.001	0.002	1.95
c9	0.292	0.003	0.99	0.324	0.002	0.75	2.147	0.022	1.04
c10	2.054	0.001	0.06	0.329	0.005	1.57	0.063	0.002	3.17
c11	1.928	0.019	1.01	0.252	0.006	1.12	0.019	0.007	3.56
c12	0.555	0.014	2.45	0.039	0.001	2.74	0.298	0.008	2.66
c13	0.258	0.004	1.44	0.406	0.001	0.18	2.246	0.017	0.75
c14	0.389	0.002	0.48	0.453	0.001	0.07	2.205	0.001	0.07
c15	0.148	0.001	0.96	0.270	0.002	0.86	1.885	0.031	1.66
c16	0.222	0.004	1.79	0.344	0.010	1.01	2.263	0.022	0.96
c17	0.178	0.001	0.48	0.350	0.004	1.05	2.126	0.022	1.01
c18	0.151	0.001	0.99	0.220	0.001	0.34	1.566	0.007	0.48
c19	0.247	0.002	0.66	0.320	0.001	0.16	2.067	0.017	0.83
c20	0.189	0.002	0.99	0.291	0.002	0.64	2.329	0.017	0.72
c21	0.147	0.002	1.27	0.278	0.002	0.67	1.694	0.016	0.99
c22	0.215	0.001	0.91	0.325	0.002	0.57	1.859	0.020	1.06
c23	0.183	0.002	1.02	0.403	0.003	0.46	2.230	0.018	0.80
c24	0.225	0.002	0.89	0.393	0.002	0.48	2.309	0.021	0.91
c25	0.926	0.031	3.37	0.076	0.022	2.87	4.340	0.127	2.93
c26	0.346	0.008	2.24	0.573	0.012	2.01	3.132	0.069	2.20
c27	0.299	0.005	1.62	0.521	0.007	1.39	2.515	0.041	1.65
c28	0.378	0.013	3.50	0.429	0.012	2.76	2.483	0.101	4.09
c29	0.404	0.04	0.90	0.459	0.006	1.30	2.733	0.021	0.78

C. heracleifolia, *C. dahurica*, and *C. foetida* (mutation) species samples were higher than *C. foetida*, *C. simplex* and *C. japonica* ones in the isoferulic acid assays. In the ferulic acid assay, we couldn't find any differences among species. In the quantitative analysis of Cimicifugae Rhizoma we indicated that, the Cimicifugae Rhizoma samples clustered two groups same as mentioned below. It was considered that *C. japonica* was clustered together with *C. foetida* and *C. simplex* due to a low quantity (0.3 mg/g) of isoferulic acid.


Pattern Recognition Analysis. To evaluate the phytochemical equivalency between the twenty nine samples corresponding to twelve authentic and seventeen commercial ones, pattern recognition analysis was conducted. In this study we used three marker compound peaks (caffeic acid (1), ferulic acid (2) and isoferulic acid (3)) for pattern recognition analysis. For pattern recognition analysis, three common peaks were selected based on the relative retention time for 4-hydroxycinnamic acid (I.S.) peak and used for a dataset. From the pattern analysis of Pam analysis (Fig. 3 and Fig. 4) and Hclust analysis (Fig. 5 and

Fig. 6) we indicated that, all of the samples were clustered to two groups (A, B). In authentic specimen analysis, we could build two clusters (A and B), and all of the species were successfully clustered two groups (A and B). In Pam analysis (Fig. 3) of twelve authentic specimen samples, we also found two clustering groups, exhibiting the same result as Pam analysis (Fig. 4). C1~c4 (*C. heracleifolia*) and c5 and c6 (*C. dahurica*) samples belong to the group A, and c7 and c8 (*C. foetida*), c10 and c11 (*C. simplex*) and c12 (*C. japonica*) belong to the group B. A mutation sample of *C. foetida*, c9, was clustered with group A because of the contents of caffeic acid (0.29 mg/g) and isoferulic acid (2.15 mg/g). The contents of caffeic acid and isoferulic acid in *C. foetida* corresponding to Group B were 1.5 ~ 3.5 mg/g and 0.001 ~ 0.3 mg/g, respectively. There are low caffeic acid and high isoferulic acid contents in Group A, whereas high caffeic acid and low isoferulic acid contents in Group B. Thus, the contents of caffeic acid and isoferulic acid in sample were important factors to build two clusters, A and B. In Pam analysis (Fig. 4) the commercial samples (c13~c29) were clustered



These two components explain 86.92% of the point variability.

Figure 3. Pam of 12 authentic specimens of Cimicifugae Rhizoma. A: (*C. heracleifolia* and *C. dahurica*), B: (*C. foetida*, *C. simplex* and *C. japonica*)



These two components explain 83.43% of the point variability.

Figure 4. Pam of 29 specimens of Cimicifugae Rhizoma including 17 commercial samples. A: (*C. heracleifolia* and *C. dahurica*), B: (*C. foetida*, *C. simplex* and *C. japonica*)

together with the group A. The commercial samples were purchased at different market and have different producer, but c13~c29 were clustered with group A. Therefore, all of the commercial samples were *C. heracleifolia*, or *C. dahurica*. Therefore, this result demonstrated that pattern recognition analysis can provide more comprehensive information for the chemical equivalency which can be omitted in the general simultaneous quantitative analysis. Thus, the pattern analysis result will be used to check the quality control of Cimicifugae Rhizoma.

Conclusions

A rapid and optimized chromatographic method with UV detection was designed for the quality control of Cimicifugae Rhizoma, well-known Korean traditional medicine. Validation




Figure 5. Hclust of 12 authentic specimens of Cimicifugae Rhizoma. A: (*C. heracleifolia* and *C. dahurica*), B: (*C. foetida*, *C. simplex* and *C. japonica*)




Figure 6. Hclust of specimens of Cimicifugae Rhizoma including 17 commercial samples. A: (*C. heracleifolia* and *C. dahurica*), B: (*C. foetida*, *C. simplex* and *C. japonica*)

data indicates that the developed analytical methods are suitable to measure the concentration of three compounds to apply to pattern recognition analysis of Cimicifugae Rhizoma. The developed HPLC/UV method for quantitative analysis of major bioactive compounds, along with a pattern-recognition method, can provide the promising prospect to comprehensive quality control of Cimicifugae Rhizoma and its related herbal medicine.

C. foetida and *C. simplex* species samples were higher than *C. heracleifolia*, *C. dahurica*, *C. foetida* (mutation) and *C. japonica* species samples in the caffeic acid assays, whereas *C. heracleifolia*, *C. dahurica*, and *C. foetida* (mutation) species samples were higher than *C. foetida*, *C. simplex* and *C. japonica* ones in the isoferulic acid assays. In the ferulic acid assay, we couldn't find any differences among species. Our results confirm that caffeic acid can serve as the species-specific marker

compounds to distinguish authentic *C. foetida* and *C. simplex*, and isoferulic acid to distinguish authentic *C. heracleifolia* and *C. dahurica* species in a pattern-recognition analysis. Therefore it is considered that caffeic acid, ferulic acid and isoferulic acid are adequate as marker compounds of quality control to distinguish the different species of Cimicifugae Rhizoma. In the pattern recognition analysis we indicated that, all of the samples were clustered to two groups A (*C. heracleifolia* and *C. dahurica*) and B (*C. foetida*, *C. simplex* and *C. japonica*), and the commercial samples (c13~c29) purchased from the markets were clustered together with group A. Therefore, the commercial samples were *C. heracleifolia* or *C. dahurica*.

Acknowledgments. This work was supported by a grant from the Korea Food and Drug Administration for Studies on Standardization of Herbal Medicine (2007).

References

- Islam, M. N.; Lee, S. K.; Jeong, S. Y.; Kim, D. H.; Jin, C.; Yoo, H. H. *Bull. Korean Chem. Soc.* **2009**, *30*, 137.
- Ajisawa, Y.; Kitazawa, M.; Uchida, M.; Kobayashi, M. *Yakugaku Zasshi* **1996**, *116*, 50.
- Noguchi, M.; Nagai, M.; Koeda, M.; Nakayama, S.; Sakurai, N.; Takahira, M.; Kusano, G. *Biol. Pharm. Bull.* **1998**, *21*, 1163.
- Kruse, S. O.; Lohning, A.; Paull, G. F.; Winterhoff, H.; Nahrstedt, A. *Planta Med.* **1999**, *65*, 763.
- Sakai, S.; Kawamata, H.; Kogure, T.; Mantani, N.; Terasawa, K.; Umatake, M.; Ochiai, H. *Mediators Inflamm.* **1999**, *8*, 173.
- Watanabe, K.; Mimaki, Y.; Sakagami, H.; Sashida, Y. *Chem. Pharm. Bull.* **2002**, *50*, 121.
- Lin, S. J.; Chen, C. S.; Lin, S. S.; Chou, M. Y.; Shih, H. C.; Lee, I. P.; Kao, C. T.; Ho, C. C.; Chen, F. L.; Ho, Y. C.; Hsieh, K. H.; Huang, C. R.; Yang, C. C. *Food Chem. Toxicol.* **2006**, *44*, 2078.
- Qiu, M.; Kim, J. H.; Lee, H. K.; Min, B. S. *Phytother. Res.* **2006**, *20*, 945.
- Jiang, B.; Yang, H.; Nuntanakorn, P.; Balick, M. J.; Kronenberg, F.; Kennelly, E. J. *J. Ethnopharmacol.* **2005**, *96*, 521.
- Jiang, B.; Kronenberg, F.; Balick, M. J.; Kennelly, E. J. *Phyto-medicine* **2006**, *13*, 477.
- He, K.; Pauli, G. F.; Zheng, B.; Wang, H.; Bai, N.; Peng, T.; Roller, M.; Zheng, Q. Y. *J. Chromatogr. A* **2006**, *1112*, 241.
- Lin, L. C.; Kuo, Y. C.; Chou, C. J. *J. Nat. Prod.* **1999**, *62*, 405.
- Wandji, J.; Nkengfack, A. E.; Fomum, Z. T.; Ubillas, R.; Killday, K. B.; Tempesta, M. S. *J. Nat. Prod.* **1990**, *53*, 1425.
- Usia, T.; Banskota, A. H.; Tezuka, Y.; Midorikawa, K.; Matsushige, K.; Kadota, S. *J. Nat. Prod.* **2002**, *65*, 673.