

Identification of a New Biflavonoid from *Selaginella doederleinii* HieronAh-Reum Han, Na-Youn Lee, Joo-Won Nam, Adam Wiryawan,[†] and Eun-Kyoung Seo^{*}

The Global Top5 Research Program, College of Pharmacy, Ewha Womans University, Seoul 120-750, Korea

^{*}E-mail: yuny@ewha.ac.kr[†]Faculty of Sciences, Brawijaya University, Malang 65145, Indonesia

Received July 1, 2013, Accepted July 12, 2013

Key Words : *Selaginella doederleinii* Hieron, Selaginellaceae, Biflavonoid

A higher plant, *Selaginella doederleinii* Hieron (Selaginellaceae) has been used as a traditional herbal medicine to treat cancer and cardiovascular disease in Southeast Asia.^{1,2} Previous phytochemical reports on this plant are including several biflavonoids, lignans, and alkaloids,^{1-3,6,7} together with biological activities such as cytotoxic,^{2,3} anti-viral,⁶ and hypertensive effects.⁷ In the present phytochemical study on *S. doederleinii*, a new biflavonoid, 2,3-dihydrorobustaflavone 7,7"-dimethyl ether (**1**) was isolated along with two known compounds, 2,3-dihydrorobustaflavone 7,4',7"-trimethyl ether (**2**) and 2",3"-dihydrorobustaflavone 7,4',7"-trimethyl ether (**3**), which were found in this plant for the first time (Figure 1). The structure of **1** was elucidated by physical and spectroscopic data analysis.

Compound **1** gave a molecular ion peak $[M+Na]^+$ at m/z 591.1286 in the HRESIMS, corresponding to an elemental formula of $C_{32}H_{24}O_{10}Na$. The absorption bands at 3376 and 1644 cm^{-1} in the IR spectrum of **1** suggested the presence of hydroxyl and conjugated carbonyl functionalities, respectively.⁸ The UV spectrum exhibited absorption maxima at 333 and 285 nm, indicating the presence of aromatic ring system(s).⁸ The ^1H -NMR spectrum of **1** showed a hydrogen-bonded hydroxyl(s) proton at δ_{H} 12.14 (2H, weak br s, OH).³ The ^1H - and ^{13}C NMR spectra of **1** indicated the presence of a flavanone skeleton as in **1**, which showed characteristic signals at δ_{H} 5.51 (1H, dd, $J = 12.8, 2.8\text{ Hz}$, H-2)/ δ_{C} 78.6 (C-2) and 3.33 (1H, H-3a) and 2.75 (1H, dd, $J = 17.2, 2.8\text{ Hz}$, H-3b)/41.9 (C-3),⁹ together with signals for aromatic systems. In the ^1H -NMR spectrum of **1**, signals for A_2X_2 coupling system at δ_{H} 7.96 (2H, d, $J = 8.6\text{ Hz}$, H-2" and 6") and 6.90 (2H, d, $J = 8.6\text{ Hz}$, H-3" and 5"), signals for 1,3,4-trisubstituted aromatic ring system at δ_{H} 7.32 (1H, dd, $J = 8.4, 2.2\text{ Hz}$, H-2'), 7.21 (1H, d, $J = 2.2\text{ Hz}$, H-6'), and 6.90 (1H, d, $J = 8.4\text{ Hz}$, H-3'), an aromatic singlet at δ_{H} 6.88 (1H, s, H-8"), and two *meta*-coupled aromatic protons at δ_{H} 6.12 (1H, d, J

$= 2.2\text{ Hz}$, H-6) and 6.07 (1H, d, $J = 2.2\text{ Hz}$, H-8), were observed. The above data suggested that there were two sets of flavonoid, 5,7,3,4'-tetrasubstituted flavanone and 5",6", 7",4"-tetrasubstituted flavone, supported by the following HMBC correlations: H-2/C-4, C-6', H-3/C-2, C-2', H-6/C-10, H-8/C-6, C-9, H-3"/C-2", C-4", -10", C-1"', H-2"' and H-6"/C-2", and H-3"' and H-5"/C-1"' (Figure 2). They are connected by a C-3'-C-6" interflavonoid linkage, as evidenced by the HMBC cross-peaks of H-2'/C-6" and H-8"/C-6". Two methoxy protons at δ_{H} 3.78 (3H, s) and 3.81 (3H, s) were assigned at C-7 and C-7", respectively, according to the HMBC correlations with carbon signals at δ_{C} 167.4 (C-7) and 162.8 (C-7"), respectively. This was also supported by the NOESY correlations of OCH_3 -7/H-6 and H-8 and OCH_3 -7"/H-8". Further detailed analysis of ^1H - ^1H COSY, ^1H - ^1H NOESY, HSQC and HMBC NMR data allowed unambiguous assignments for all of the ^1H and ^{13}C NMR signals of **1**. Therefore, the structure of **1** corresponded to the robustaflavone series¹⁰ and determined as a new compound, 2,3-dihydrorobustaflavone 7,7"-dimethyl ether. The circular dichroism (CD) experiment was performed to determine an absolute configuration at C-2 of **1**. However, the CD spectrum of **1** exhibited a flat CD band without a positive or negative cotton effect, indicating the presence of a racemic mixture, but not in a 1:1 mixture due to the slightly positive specific rotation observed.




Figure 2. Selected ^1H - ^1H COSY (—), ^1H - ^1H NOESY (---), and HMBC (→) correlations of compound **1**.




Figure 1. Structures of the isolates **1-3** from the whole plants of *S. doederleinii*.

Compounds **2** and **3** were identified as 2,3-dihydrorobustaflavone 7,4',7''-trimethyl ether (**2**)¹¹ and 2'',3''-dihydrorobustaflavone 7,4',7''-trimethyl ether (**3**),¹² respectively, by analyses of their physical and spectroscopic data as well as by comparison of their data with the published values. Compound **2** was optically active ($[\alpha]_D^{25}$: -10.60), however, the absolute configuration at C-2 could not be determined due to the flat CD band shown in the CD spectrum of **2**. This observation indicated that **2** present as a racemic mixture. The configuration at C-2'' of **3** was determined as *S*, due to its CD spectrum (a positive cotton effect at 333 nm and a negative cotton effect at 287 nm).¹³ To the best of our knowledge, **2** and **3** were isolated from *S. doederleinii* for the first time.

Experimental Section

General Experimental Procedures. Optical rotations were measured with a P-1010 polarimeter (JASCO, Japan) at 25 °C. UV spectra (λ_{\max}) were recorded on a U-3000 spectrophotometer (Hitachi, Japan). IR spectra (ν_{\max}) were determined on a FTS 135 FT-IR spectrometer (Bio-Rad, CA). Circular dichroism measurements were performed using JASCO J-715 CD/ORD spectropolarimeter. The 1D and 2D NMR experiments were conducted on a UNITY INOVA 400 MHz FT-NMR (varian, CA, USA), and TMS was used as an internal standard. ESIMS and HRESIMS were obtained on a JMS 700 Mastation HRMS spectrometer (JEOL, Japan). Thin-layer chromatographic (TLC) analysis was performed on Kieselgel 60 F₂₅₄ and RP-18 F_{254s} silica gel plates (Merck, Germany) with visualization under UV light (254 and 365 nm) and 10% (v/v) sulfuric acid spray followed by heating (120 °C, 5 min). Silica gel (230-400 mesh, Merck, Germany), YMC*GEL ODS-A (S-150 μ m, YMC Co., Ltd., Japan), and Sephadex LH-20 (Amersham Pharmacia Biotech, U.K.) were used for column chromatography.

Plant Material. The whole plants of *S. doederleinii* were collected in Batu Medica Centre, East Java, Indonesia, in May 2005 and were identified by prof. Adam Wiryawan (Brawijaya University, Malang 65145, Indonesia). A voucher specimen has been deposited at the Batu Herba Medica Centre.

Extraction and Isolation. The dried whole plants of *S. doederleinii* (335 g) was extracted with 95% MeOH (3 \times 8 L) overnight at room temperature. The solvent was evaporated *in vacuo* to afford a MeOH extract (25 g), which was subjected to silica gel column chromatography (CC) using gradient mixtures of hexanes-EtOAc (49:1 \rightarrow 0:1) to give 13 fractions (FI-FXIII). Fraction FXII (7.4 g) was chromatographed on Sephadex LH-20 (100% MeOH), providing five sub-fractions (FXII-1 – FXII-5). The sub-fraction FXII-1 (15 mg) was subjected to reversed-phase CC eluted with gradient mixtures of MeOH-H₂O (3:2 \rightarrow 4:1), yielding **3** (1.8 mg). The sub-fraction FXII-2 (90 mg) was separated by reversed-phase CC eluted with gradient mixtures of MeOH-H₂O (4:1 \rightarrow 2:1), affording five sub-fractions (FXII-2-1 – FXII-2-5). The sub-fraction FXII-2-5 (30 mg) was carried

out preparative TLC (CHCl₃-MeOH = 49:1) to afford **2** (2.0 mg). The combined fractions of FXII-3 and FXII-4 (300 mg) were chromatographed over silica gel eluted with gradient mixtures of CHCl₃-MeOH (99:1 \rightarrow 1:1), yielding eight sub-fractions (FXII-3-1 – FXII-3-8). FXII-3-3 (24 mg) was purified by preparative TLC using a solvent system of CHCl₃-MeOH (49:1) to provide **1** (1.5 mg).

2,3-Dihydrorobustaflavone 7,7''-dimethyl ether (1): Yellow powder. $[\alpha]_D^{25}$: +13.00 (*c* 0.1, MeOH); UV (MeOH) λ_{\max} (log ϵ) 333 (4.6), 285 (4.7) nm; IR (KBr) ν_{\max} cm⁻¹: 3376, 1644, 1604, 1450, 1349, 1202, 1156; ¹H-NMR (DMSO-*d*₆, 400 MHz) δ 12.14 (2H, s, OH), 7.96 (2H, d, *J* = 8.6 Hz, H-2''' and 6'''), 7.32 (1H, dd, *J* = 8.4, 2.2 Hz H-6'), 7.21 (1H, d, *J* = 2.2 Hz, H-2'), 6.90 (1H, d, *J* = 8.4 Hz, H-5'), 6.90 (2H, d, *J* = 8.6 Hz, H-3''' and 5'''), 6.88 (1H, s, H-8'''), 6.80 (1H, s, H-3''), 6.12 (1H, d, *J* = 2.2 Hz, H-6), 6.07 (1H, d, *J* = 2.2 Hz, H-8), 5.51 (1H, dd, *J* = 12.8, 2.8 Hz, H-2), 3.81 (3H, s, OCH₃-7''), 3.78 (3H, s, OCH₃-7), 3.33 (1H, H-3), 2.75 (1H, dd, *J* = 17.2, 2.8 Hz, H-3); ¹³C-NMR (DMSO-*d*₆, 100 MHz) δ 197.0 (C-4), 181.6 (C-4''), 167.4 (C-7), 163.8 (C-2''), 163.2 (C-9), 162.9 (C-5), 162.8 (C-7''), 162.5 (C-4'''), 156.8 (C-9''), 156.2 (C-4'), 131.1 (C-6'), 128.4 (C-2''' and 6'''), 128.0 (C-2'), 127.2 (C-1'), 120.1 (C-5' and 1'''), 116.2 (C-3', 5''' and 3'''), 110.3 (C-6''), 104.9 (C-10''), 102.8 (C-10), 102.6 (C-3''), 94.6 (C-8), 93.7 (C-6), 90.1 (C-8''), 78.6 (C-2), 56.1 (OCH₃-7''), 55.8 (OCH₃-7), 41.9 (C-3); HRESIMS *m/z* = 591.1286 [M+Na]⁺ (calcd for C₃₂H₂₄O₁₀Na: 591.1261).

2,3-Dihydrorobustaflavone 7,4',7''-Trimethyl Ether (2): $[\alpha]_D^{25}$: -10.60 (*c* 0.15, MeOH; literature values: -13.3, *c* 0.75, dioxane¹¹).

2'',3''-Dihydrorobustaflavone 7,4',7''-Trimethyl Ether (3): $[\alpha]_D^{25}$: -5.55 (*c* 0.18, MeOH; literature values: -2.3, *c* 0.44, dioxane¹²); CD (*c* 2.0 \times 10⁻³ mM, MeOH) $\Delta\epsilon_{287}$ -0.48, $\Delta\epsilon_{333}$ +0.22 nm.

Acknowledgments. This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (MEST) (2011-0003682). The work was also supported in by the National Research Foundation of Korea (NRF) grant funded by the Korea Government (MEST) (No. 2012-0000952).

Supporting Information. The spectral data of compound **1** are available on request from the correspondence author.

References

- Lin, R. C.; Skaltsounis, A.-L.; Seguin, E.; Tillequin, F.; Koch, M. *Planta Med.* **1994**, *60*, 168.
- Chao, L. R.; Seguin, E.; Tillequin, F.; Koch, M. *J. Nat. Prod.* **1987**, *50*, 422.
- Lee, N.-Y.; Min, H.-Y.; Lee, J.; Nam, J.-W.; Lee, Y.-J.; Han, A.-R.; Wiryawan, A.; Suprpto, W.; Lee, S. K.; Seo, E.-K. *Chem. Pharm. Bull.* **2008**, *56*, 1360.
- Cheng, X.-L.; Ma, S.-C.; Yu, J.-D.; Yang, S.-Y.; Xiao, X.-Y.; Hu, J.-Y.; Lu, Y.; Shaw, P.-C.; But, P. P.-H.; Lin, R.-C. *Chem. Pharm. Bull.* **2008**, *56*, 982.
- Silva, G. L.; Chai, H.; Gupta, M. P.; Farnsworth, N. R.; Cordell, G.

- A.; Pezzuto, J. M.; Beecher, C. W. W.; Kinghorn, A. D. *Phytochemistry* **1995**, *40*, 129.
6. Ma, S.-C.; But, P. P.-H.; Ooi, V. E.-C.; He, Y.-H.; Lee, S. H.-S.; Lee, S.-F.; Lin, R.-C. *Biol. Pharm. Bull.* **2001**, *24*, 311.
7. Liu, R. C.; Peyroux, J.; Seguin, E.; Koch, M. *Phytother. Res.* **1991**, *5*, 188.
8. Pavia, D. L.; Lampman, G. M.; Kriz, G. S. *Introduction to Spectroscopy: A Guide for Students of Organic Chemistry*; Thomson Learning, Ltd.: London, 2001; p 13, 353.
9. Mabry, T. J.; Markham, K. R.; Thomas, M. B. *The Systematic Identification of Flavonoids*; Springer: New York, 1970.
10. Markham, K. R.; Sheppard, C.; Geiger, H. *Phytochemistry* **1987**, *26*, 3335.
11. Lin, L.-C.; Chou, C.-J. *Chin. Pharm. J. (Taipei)* **2000**, *52*, 211.
12. Lin, L.-C.; Kuo, Y.-C.; Chou, C.-J. *J. Nat. Prod.* **2000**, *63*, 627.
13. Gaffield, W. *Tetrahedron* **1970**, *26*, 4093.
-