

The relationship between some endometrial secretion cytokines and in vitro fertilization

Mohammad Ehsan Rahiminejad¹ M.D., Amirhossein Moaddab² M.D., Mehrnoosh Ebrahimi¹ M.D., Soghra Rabiee¹ M.D., Alireza Zamani³ Ph.D., Mohammad Ezzati⁴ M.D., Alireza Abdollah Shamshirsaz² M.D.

1. Department of Obstetrics & Gynecology, Hamadan University of Medical Sciences and Health Services, Hamadan, Iran.

2. Department of Obstetrics and Gynecology, Baylor College of Medicine, Houston, Texas, USA.

3. Department of Immunology and Hematology, Hamadan University of Medical Sciences and Health Services, Hamadan, Iran.

4. Department of Obstetrics and Gynecology, UT Southwestern Medical Center, Dallas, TX, USA.

Corresponding Author:

Amirhossein Moaddab, F1040-01, 6621 Fannin St, Pavilion for women, Baylor College of Medicine, Houston, TX 77030.
Email: moaddab@bcm.edu;
amirh_moaddab@yahoo.com
Tel: (+98) 832766-6029

Received: 29 October 2014

Revised: 5 February 2015

Accepted: 21 June 2015

Abstract

Background: Endometrial secretion analysis is a non-invasive and promising method in evaluation of endometrial receptivity.

Objective: The aim of the present study was to assess the relationship between the success rate of IVF procedures and some endometrial secretion cytokines, including interleukin-1 β (IL-1 β), tumor necrosis factor (TNF- α), interferon gamma-induced protein 10 (IP-10), and monocyte chemoattractant protein (MCP).

Materials and Methods: In a prospective cohort study, 50 women selected for IVF met the study inclusion criteria. All the patients underwent endometrial secretion aspiration prior to embryo transfer. The level of IL-1 β , TNF- α , IP-10 and MCP were analyzed by enzyme-linked immunosorbent assay method using special standard kits. To detect successful implantation and pregnancy patients underwent serum human chorionic gonadotropin measurements and ultrasound evaluation.

Results: Five samples were excluded. Nine women (20%) had successful clinical pregnancies, which resulted in live birth. Other 36 women (80%) were classified as failed pregnancy. Comparison of cytokine levels showed lower concentrations of TNF- α , IP-10, and MCP in the group with successful clinical pregnancy compared to the group with failed pregnancy ($p=0.007$, 0.005 and 0.001 , respectively). However, no significant difference was revealed in IL-1 β levels between two groups ($p=0.614$).

Conclusion: The current study suggested that lower concentrations of TNF- α , IP-10, and MCP in endometrial secretions might be associated with improved endometrial receptivity and IVF outcome. Regarding IL-1 β , no statistically significant differences were seen between the groups with and without successful pregnancy.

Key words: Cytokine, In vitro fertilization, Outcome, Endometrium, Implantation.

Introduction

Infertility is a common condition, affecting marital relationship, mental health and the couples' quality of life. Recent advances in assisted reproductive technologies (ARTs) such as In Vitro Fertilization (IVF) provide more efficient methods for infertility treatment (1-3). Despite the increasing prevalence, most infertility treatment modalities are very expensive and impose a heavy financial burden on families. Moreover, a significant number of IVF procedures does not result in a live birth (3-5).

Successful implantation depends on the quality of the embryo and the endometrial receptivity. (6) Cytokines produced by fetal and uterine mucus are responsible for regulating interactions between the mother and fetus, which improve uterus receptivity by

controlling the expression of proteins involved in the adhesion process (7, 8). The known cytokines and growth factors that may have contributed to the increased receptivity of human endometrial implantation include leukemia inhibitory factor, interleukin-1, keratinocyte growth factor and colony stimulating factor-1. An ideal biomarker for increased endometrial receptivity must be present in the endometrium, should be close to the site of implantation and should appear during the implantation period and disappear after it (9).

Identification of such biomarkers may provide more information about implantation in both normal and ART cycles. Furthermore, the identification of these markers might be useful in determining the best time for embryo transfer. The aim of the present study was to assess the relationship between the success

rate of IVF procedures and some endometrial secretion cytokines, including interleukin-1 β (IL-1 β), tumor necrosis factor (TNF- α), interferon gamma-induced protein 10 (IP-10), and monocyte chemoattractant protein (MCP). These cytokines were chosen based on controversial results of previous studies focused on describing the immunological dialogue between a vital embryo and a receptive endometrium in implantation (1, 6-8).

Materials and methods

Study design

This prospective non-randomized study were performed at Fatemeh Women's Hospital from September 2011 to March 2013. Fifty women less than 35 years of age who were candidates for IVF due to tubal factor infertility were enrolled after providing sufficient explanation and obtaining informed consent. The study protocol was approved by the Research Ethics Committee of Hamadan University of Medical Sciences. To confine the effects of factors which affect IVF outcome, all women met the study inclusion criteria, including a) normal menstrual cycles between 25 to 35 days, b) female aged less than 35 years old, c) BMI less than 30, d) TSH <10 on the third day of the cycle and the number of antral follicles more than 6 on the third day ultrasound, and e) normal semen analysis. Also, exclusion criteria were as follows: a) metabolic and/or endocrine disorders (diabetes, metabolic syndrome, PCOS and Thyroid Disorders), b) previous gynecological/pelvic surgery except for salpingectomy, c) more than once previous failed IVF cycle, and d) smoking.

IVF technique

To stimulate ovulation, recombinant follicular stimulating hormone (rFSH) (Puregon, Organon, Germany) was injected daily from the third day of the menstrual cycle with an appropriate dosage. In order to prevent premature luteinization, daily injection of gonadotropin-releasing hormone (GnRH) antagonist (Ganirelix acetate, Organon, Netherland) (450 IU once daily) was considered from the fifth day of the IVF cycle. In order to stimulate final oocyte maturation, an intramuscular dose (10000 IU) of human chorionic gonadotropin (hCG) (Pregnyle,

Organon, Germany) was injected. A maximum of three embryos were transferred, 4-5 days after the ovarian puncture. Embryo transfer was performed in the luteal phase. An embryo with stage-specific cell size, <10% of fragmentation and no multinucleation was considered as grade A. Embryos with stage-specific cell size for the majority of cells, 10–25% of fragmentation and no evidence of multinucleation considered as grade B and embryos with cell size not stage-specific, severe fragmentation (more than 25%) and evidence of multinucleation counted as grade C (10).

During the first 5 days after embryo transfer, three hCG injections (each containing 5000 IU) and intramuscular progesterone (Progestan, Organon, Germany) injection (100 mg/day) were administered for luteal phase support.

Serum β -hCG was measured 10 days after embryo transfer in order to detect implantation. In case of a positive test result, the test was repeated 48 hours later. A double increase in β -hCG level was considered as successful implantation. Women with successful implantation underwent ultrasound evaluation of the pregnancy from four weeks after oocyte retrieval. Pregnancy was defined by visualization of an embryonic sac at vaginal ultrasound examination at 5 weeks after embryo transfer. Losing pregnancy products diagnosed by ultrasound any time after pregnancy detection was considered as a failed IVF cycle.

Aspiration and analysis of endometrial secretions

Aspiration of endometrial secretion was performed by the method described by Boomsma (1). Endometrial secretions were aspirated by insertion a trans-cervical catheter and gentle suction with a 2-mL syringe prior to embryo transfer. In previous studies, the safety of the uterine secretions aspirated prior to embryo transfer have shown [11]. All the samples were stored in liquid nitrogen at a temperature of -80°C. The level of cytokines including interleukin-1 β (IL-1 β), tumor necrosis factor (TNF- α), interferon gamma-induced protein 10 (IP-10), and monocyte chemoattractant protein (MCP) in the aspirated fluid were analyzed by enzyme-linked immunosorbent assay (ELISA) method

using special standard kits. (R&D Systems, Minneapolis, Minnesota).

Statistical analysis

Values were mean \pm SD/median (25; 75th percentiles) and data were analyzed statistically by χ^2 test or Mann–Whitney U-test. Statistical Package for the Social Sciences, version 19.0, SPSS Inc, Chicago, Illinois, USA (SPSS software) was used for statistical analysis. $P < 0.05$ was considered statistically significant.

Results

Fifty patients were recruited for this study. Five samples were excluded because of contact bleeding in the course of sampling. Thus, aspirated endometrial secretions of 45

enrolled women were analyzed to assess levels of IL-1 β , TNF- α , IP-10, and MCP. According to serum β -hCG levels and ultrasound studies, 9 women (20%) had successful clinical pregnancies which resulted in live birth. Other 36 women (80%) were classified as the group with failed pregnancy. There was no significant difference between women with successful pregnancy and failed pregnancy in terms of baseline and clinical characteristics (Table I). The concentrations of TNF- α , IP-10, and MCP in the group with successful clinical pregnancy were significantly different compared to the group with failed pregnancy (p -values 0.007, 0.005 and 0.001, respectively) (Table II). However, no significant difference was revealed in IL-1 β levels between two groups.

Table I. Baseline and clinical characteristics in groups with successful pregnancy and failed pregnancy

Characteristics	Successful pregnancy (n = 9)	Failed pregnancy (n = 36)	p-value
Age, yr (Mean \pm SD)	28.0 \pm 3.3	29.7 \pm 3.5	0.13
Duration of infertility, yr (Mean \pm SD)	5.7 \pm 2.2	5.9 \pm 2.6	0.69
Number of transferred embryos (Mean \pm SD)	2.2 \pm 0.8	2.2 \pm 0.9	1.00
Grade of transferred embryo			
A*	9 (100%)	29 (80.6%)	0.06
B $^{\Delta}$	0	6 (16.7%)	-
C †	0	1 (2.7%)	-

* Embryo with stage-specific cell size, <10% fragmentation and no multinucleation

Δ Embryo with stage-specific cell size for the majority of cells, 10–25% fragmentation and no evidence of multinucleation

\dagger Embryo with cell size not stage-specific, severe fragmentation (25%) and evidence of multinucleation

Table II. Comparison of cytokine levels in aspirated endometrial secretions in with successful pregnancy and failed pregnancy

Cytokine	Successful pregnancy (n = 9)	Failed pregnancy (n = 36)	p-value
TNF- α *	13.5(10.3;230.45) ■	25.1(17.9;32.7)	<0.01
IL-1 β $^{\Delta}$	363.7(102.3;636.4)	49.2(35.3;887.2)	0.61
MCP †	250.5(134.2;372.5)	1050.3(106.3;1995.6)	<0.01
IP-10 $^{\circ}$	298.2(267.6;407.4)	634.7(290.7;996.3)	<0.01

* Tumor necrosis factor- α

$^{\circ}$ Interferon gamma-induced protein 10

Δ Interleukin-1 β

■ All levels were measured in pg/mL and represents median (25; 75th percentiles)

† Monocyte chemoattractant protein

Discussion

The current study suggested that lower concentrations of TNF- α , IP-10, and MCP in endometrial secretions were associated with improved uterine receptivity and IVF outcome. Regarding IL-1 β , no statistically significant

difference was seen between the groups with and without successful pregnancy.

The interactions between maternal immune system and conceptus tissues at implantation are necessary for successful implantation and progression to pregnancy (12). It seemed that these interactions at the level of endometrium

regulated cytokines concentrations and Th1/Th2 ratio. Recent studies have shown that some of the secreted inflammatory cytokines not only make no disturbance in implantation, but also are necessary for endometrial reconstruction and modulate the events subsequent to implantation by recruiting the macrophages and dendritic cells (13-16). Some studies hypothesized that slight damage to the endometrium in women with repeated IVF failure led to a greater clinical pregnancy rate by modifying endometrial secreted cytokines (17-20).

Previous studies reported some controversial results on the role of different cytokines in providing a proper physiological function for implantation (1, 6-8). TNF- α is not only a key regulator in interactions prior to the implantation, but also higher levels of it plays a crucial role in some reproductive diseases such as endometrial infections and recurrent spontaneous abortions (21-24). Boomsma *et al* study revealed a positive relationship between successful pregnancy and higher levels of TNF- α in endometrial secretions. Similar to earlier animal studies, we found that TNF- α concentrations in pregnant women with successful pregnancy were significantly lower than those with IVF failure (25).

While previous studies have shown that inappropriate expression of IL-1 β was associated with repeated abortions and pregnancy failure, the current study failed to find any statistically significant difference between the study groups (25-28).

In a study by Boomsma *et al*, MPC1 and IP10 levels had significant negative and positive association with initial implantation of embryos, respectively (25). Some studies showed that the embryo expressed some receptors for MPC1 and IP10 (29-35). Furthermore, it was suggested that MPC1 was a strong absorber and activator of uterine natural killer cells, which were associated with abortion and infertility (32-33). Given the relationship between higher rates of implantation with lower levels of these cytokines, these findings were probably consistent with the results of the current study.

Comparing to other studies, the chance to achieve a successful IVF in this study was low (10, 36). We determined the concentrations of the endometrial secretions by ELISA because of its feasibility and regarding the results of previous studies, which demonstrated the

similar results of endometrial cytokine measurements by immunohistochemistry, ELISA and semiquantitative reversed transcription-polymerase chain reaction (semiquantitative RT-PCR) (37). In addition to theoretical uncertainties, some factors such as technical differences between this study and previous ones in secretions aspiration or timing of aspiration as well as lower volume of samples could be possible reasons for this difference.

Considerable evidence suggests that pregnancy is a complex event related to complex immunological processes. While infertility and failed ART cycles are tied to emotional distress, further understanding of this phenomenon might lead to more control on factors contributing to the success and failure of pregnancy process and maximize the estimation of success rate (38). The non-invasive nature of analysis of uterine secretion aspiration enabled the researchers of this study to evaluate the uterine receptivity before the embryo transfer in IVF process without inflicting any damage comparing to alternative methods such as endometrial dating.

Conclusion

The current study suggests that lower concentrations of TNF- α , IP-10, and MCP in endometrial secretions might be associated with improved endometrial receptivity and IVF outcome. This non-invasive approach may improve understanding of immunological events, which embryo experiences during its early days.

Acknowledgements

This study was supported by a grant from Vice Chancellor of Research of Hamadan University of Medical Sciences.

Conflict of interest

The authors declare that they have no conflict of interest.

References

1. Boomsma CM, Kavelaars A, Eijkemans MJ, Fauser BC, Heijnen CJ, Macklon NS. Ovarian stimulation for in vitro fertilization alters the intrauterine cytokine, chemokine, and growth factor milieu encountered by the embryo. *Fertil Steril* 2009; 94: 1764-1768.

2. Campbell KL, Rockett JC. Biomarkers of ovulation, endometrial receptivity, fertilization, implantation and early pregnancy progression. *Paediatr Perinat Epidemiol* 2006; 20: 13-25.
3. Cavagna M, Mantese JC. Biomarkers of endometrial receptivity-a review. *Placenta* 2003; 24: 39-47.
4. Dikareva LV, Shvarev EG, Shvarev GE, Teplyi DL. Age, structural and biochemical characteristics of endometrial secretion in patients with hysteromyoma. *Adv Gerontol* 2008; 21: 596-601.
5. Florio P, Bruni L, Galleri L, Reis FM, Borges LE, Bocchi C, et al. Evaluation of endometrial activin A secretion for prediction of pregnancy after intrauterine insemination. *Fertil Steril* 2009; 93: 2316-2320.
6. Haouzi D, Mahmoud K, Fourar M, Bendhaou K, Dechaud H, De Vos J, et al. Identification of new biomarkers of human endometrial receptivity in the natural cycle. *Hum Reprod* 2009; 24: 198-205.
7. Lieberman JA, Moscicki AB, Sumerel JL, Ma Y, Scott ME. Determination of cytokine protein levels in cervical mucus samples from young women by a multiplex immunoassay method and assessment of correlates. *Clin Vaccine Immunol* 2008; 15: 49-54.
8. Nieuwenhuizen L, Khalil MK, Venkatesh N, Othman NH. Endometrial and endocervical secretion: the search for histochemical differentiation. *Anal Quant Cytol Histol* 2006; 28: 87-96.
9. van der Gaast MH, Macklon NS, Beier-Hellwig K, Krusche CA, Fauser BC, Beier HM, et al. The feasibility of a less invasive method to assess endometrial maturation--comparison of simultaneously obtained uterine secretion and tissue biopsy. *BJOG* 2009; 116: 304-312.
10. Alpha Scientists in Reproductive Medicine and ESHRE Special Interest Group of Embryology. The Istanbul consensus workshop on embryo assessment: proceedings of an expert meeting. *Hum Reprod* 2011; 26: 1270-1283.
11. van der Gaast MH, Beier-Hellwig K, Fauser BC, Beier HM, Macklon NS. Endometrial secretion aspiration prior to embryo transfer does not reduce implantation rates. *Reprod Biomed Online* 2003; 7: 105-109.
12. Robertson SA, Moldenhauer LM. Immunological determinants of implantation success. *Int J Dev Biol* 2014; 58: 205-217.
13. Rajaei S, Zarnani AH, Jeddi-Tehrani M, Tavakoli M, Mohammadzadeh A, Dabbagh A, et al. Cytokine profile in the endometrium of normal fertile and women with repeated implantation failure. *Iran J Immunol* 2011; 8: 201-208.
14. Wegmann TG, Lin H, Guilbert L, Mosmann TR. Bidirectional cytokine interactions in the maternal-fetal relationship: is successful pregnancy a TH2 phenomenon. *Immunol Today* 1993; 14: 353-356.
15. Chaouat G, Ledee-Bataille N, Dubanchet S, Zourbas S, Sandra O, Martal J. TH1/TH2 paradigm in pregnancy: paradigm lost? Cytokines in pregnancy/early abortion: reexamining the TH1/TH2 paradigm. *Int Arch Allergy Immunol* 2004; 134: 93-119.
16. Chaouat G. The Th1/Th2 paradigm: still important in pregnancy? *Semin Immunopathol* 2007; 29: 95-113.
17. Granot I, Gnainsky Y, Dekel N. Endometrial inflammation and effect on implantation improvement and pregnancy outcome. *Reproduction* 2012; 144: 661-668.
18. Karimzadeh MA, Ayazi Rozbahani M, Tabibnejad N. Endometrial local injury improves the pregnancy rate among recurrent implantation failure patients undergoing in vitro fertilisation/intra cytoplasmic sperm injection: a randomised clinical trial. *Aust N Z J Obstet Gynaecol* 2009; 49: 677-680.
19. Zhou L, Li R, Wang R, Huang HX, Zhong K. Local injury to the endometrium in controlled ovarian hyperstimulation cycles improves implantation rates. *Fertil Steril* 2008; 89: 1166-1176.
20. El-Toukhy T, Sunkara S, Khalaf Y. Local endometrial injury and IVF outcome: a systematic review and meta-analysis. *Reprod Biomed Online* 2012; 25: 345-354.
21. Toder V, Fein A, Carp H, Torchinsky A. TNF-alpha in pregnancy loss and embryo maldevelopment: a mediator of detrimental stimuli or a protector of the fetoplacental unit. *J Assist Reprod Genet* 2003; 20: 73-81.
22. Meisser A, Chardonnens D, Campana A, Bischof P. Effects of tumour necrosis factor-alpha, interleukin-1 alpha, macrophage colony stimulating factor and transforming growth factor on trophoblastic matrix metalloproteinases. *Mol Hum Reprod* 1999; 5: 252-260.
23. Cohen M, Meisser A, Haenggeli L, Bischof P. Involvement of MAPK pathway in TNF-alpha induced MMP-9 expression in human trophoblastic cells. *Mol Hum Reprod* 2006; 12: 225-232.
24. Lockwood CJ, Oner C, Uz YH, Kayisli UA, Huang SJ, Buchwalder LF, et al. Matrix metalloproteinase 9 expression in preeclamptic decidua and MMP9 induction by tumor necrosis factor alpha and interleukin 1 beta in human first trimester decidual cells. *Biol Reprod* 2008; 78: 1064-1072.
25. Boomsma CM, Kavelaars A, Eijkemans MJ, Lentjes EG, Fauser BC, Heijnen CJ, et al. Endometrial secretion analysis identifies a cytokine profile predictive of pregnancy in IVF. *Hum Reprod* 2009; 24: 1427-1435.
26. Inagaki N, Stern C, McBain J, Lopata A, Kornman L, Wilkinson D. Analysis of intra-uterine cytokine concentration and matrix-metalloproteinase activity in women with recurrent failed embryo transfer. *Hum Reprod* 2003; 18: 608-615.
27. von Wolff M, Thaler CJ, Strowitzki T, Broome J, Stolz W, Tabibzadeh S. Regulated expression of cytokines in human endometrium throughout the menstrual cycle: dysregulation in habitual abortion. *Mol Hum Reprod* 2000; 6: 627-634.
28. Jones RL, Hannan NJ, Kaitu'u TJ, Zhang J, Salamonsen LA. Identification of chemokines important for leukocyte recruitment to the human endometrium at the times of embryo implantation and menstruation. *J Clin Endocrinol Metab* 2004; 89: 6155-6167.
29. Inngjerdigen M, Damaj B, Mghazachi AA. Expression and regulation of chemokine receptors in human natural killer cells. *Blood* 2001; 97: 367-375.
30. Dominguez F, Pellicer A, Simon C. The chemokine connection: hormonal and embryonic regulation at the human maternal-embryonic interface-a review. *Placenta* 2003; 24: 48-55.
31. Caballero-Campo P, Domínguez F, Coloma J, Meseguer M, Remohí J, Pellicer A, et al. Hormonal

- and embryonic regulation of chemokines IL-8, MCP-1 and RANTES in the human endometrium during the window of implantation. *Mol Hum Reprod* 2002; 8: 375-384.
32. Loetscher P, Seitz M, Clark-Lewis I, Baggiolini M, Moser B. Monocyte chemotactic proteins MCP-1, MCP-2, and MCP-3 are major attractants for human CD4+ and CD8+ T lymphocytes. *FASEB J* 1994; 8: 1055-1060.
 33. Moffett A, Regan L, Braude P. Natural killer cells, miscarriage, and infertility. *BMJ* 2004; 329: 1283-1285.
 34. Sela HY, Goldman-Wohl DS, Haimov-Kochman R, Greenfield C, Natanson-Yaron S, Hamani Y, et al. Human trophoblast apposition is regulated by interferon gamma-induced protein 10 (IP-10) during early implantation. *Placenta* 2013; 34: 222-230.
 35. Nagaoka K, Nojima H, Watanabe F, Chang KT, Christenson R, Sakai S, et al. Regulation of blastocyst migration, apposition and initial adhesion by a chemokine, IFN-gamma -inducible protein 10 kDa (IP-10), during early gestation. *J Biol Chem* 2003; 278: 29048-29056.
 36. Gianaroli L, Magli MC, Ferraretti AP, Iammarrone E. Preimplantation genetic diagnosis increases the implantation rate in human in vitro fertilization by avoiding the transfer of chromosomally abnormal. *Fertil Steril* 1997; 68: 1128-1131.
 37. Von Wolff M, Classen-Linke I, Heid D, Krusche CA, Beier-Hellwig K, Karl C, et al. Tumour necrosis factor-alpha (TNF-alpha) in human endometrium and uterine secretion: an evaluation by immunohistochemistry, ELISA and semiquantitative RT-PCR. *Mol Hum Reprod* 1999; 5: 146-152.
 38. Rahiminejad ME, Moaddab A, Rabiee S, Esna-Ashari F, Borzouei S, Hosseini SM. The relationship between clinicobiochemical markers and depression in women with polycystic ovary syndrome. *Iran J Reprod Med* 2014; 12: 811-816.

Archive of SID