

SOME PROPERTIES OF PROSTACYCLIN SYNTHASE FROM PIG AORTA

Paulina WLODAWER and Sven HAMMARSTRÖM

Department of Chemistry, Karolinska Institutet, S-104 01 Stockholm, Sweden

Received 1 November 1978

1. Introduction

Two potent bioregulators can be formed from the prostaglandin endoperoxide PGH_2 : thromboxane A_2 (TXA_2), a platelet-aggregating and smooth muscle-contracting agent [1] and prostacyclin (PGI_2) which inhibits platelet aggregation and relaxes smooth muscles [2]. The thromboxane synthase has been partially purified from platelets [3] and lung [4], and shown to also catalyze the conversion of PGH_2 to 12-hydroxy-5,8,10-heptadecatrienoic acid (HHT) [5,6]. A possible mechanism of these transformations is shown in fig.1. It is postulated that the reactions are initiated by protonation of the oxygen at C-9 of PGH_2 and that the charged intermediate (I) rearranges to TXA_2 or cleaves to HHT plus malondialdehyde.

On the other hand, protonation of the oxygen at C-11 could initiate a rearrangement of PGH_2 to PGI_2 . The intermediate (II) might also undergo a similar cleavage as (I) to give HHT and malondialdehyde.

This paper describes the solubilization and fractionation by DEAE cellulose chromatography of prostacyclin synthase. On the basis of these experiments, the use of enzyme inhibitors, and protein denaturing agents, it is concluded that PGI_2 synthase does not enzymatically form HHT.

Abbreviations EMPA, 9 α ,11 α -epoxymethano-15(S)hydroxyprosta-5(cis)-13(trans)-dienoic acid, HHT, 12-hydroxy-5,8,10-heptadecatrienoic acid; L-8027, 2-isopropyl-3-nicotinylindole, N-0164, *p*-benzyl-4-[1-oxo-2-(4-chlorobenzyl)-3-phenylpropyl]-phenyl phosphonate; PG, prostaglandin; TLC, thin-layer chromatography, TXB_2 , thromboxane B_2 , 15-HPETE, 15-hydroperoxy 5,8,11,13-eicosatetraenoic acid




Fig.1. Possible mechanisms of thromboxane A_2 (TXA_2) and prostacyclin (PGI_2) formation from PGH_2 .

2. Materials and methods

2.1. Chemicals

Prostaglandins $\text{F}_{2\alpha}$, E_2 , 6-keto- $\text{PGF}_{1\alpha}$ and EMPA (9 α ,11 α -epoxymethano-15(S)hydroxyprosta-5(cis), 13(trans)-dienoic acid) were obtained from the Upjohn Company, Kalamazoo, MI. 2-Iso-propyl-3-nicotinylindole (L-8027) and sodium *p*-benzyl-4-[1-oxo-2-(4-chlorobenzyl)-3-phenylpropyl]phenyl phosphonate (N-0164) were given by Labaz and Nelson Research, respectively. [$1\text{-}^{14}\text{C}$] PGH_2 (spec. act. 1.0 Ci/mol) and 12L-hydroxy-5,8,10-heptadecatrienoic acid (HHT) were prepared by the methods in [7,8]. DEAE cellulose was from Whatman, Silica gel G, imidazole and SnCl_2 from Merck.

2.2. Enzyme preparation

Pig aorta was obtained from slaughter house and immediately frozen on dry ice. Portions of tissue (~50 g) were frozen in liquid nitrogen, crushed into a fine powder and homogenized in 2 vol. 0.1 M potassium phosphate buffer (pH 7.4). Subcellular fractions were obtained by differential centrifugation, as for bovine lung [4]. The mitochondrial and microsomal fractions (sedimented at $9000 \times g$, 15 min and $100\,000 \times g$, 60 min, respectively) were resuspended in the same buffer to a concentration corresponding to 2 g tissue/ml.

Solubilization of microsomes and DEAE-cellulose chromatography were performed as for platelet [3] and bovine lung [4] thromboxane synthase.

2.3. Enzyme assay

Incubations were performed in 0.1 M potassium phosphate buffer (pH 7.4) and details are shown in appropriate legends. The reaction was stopped with 5 vol. ethanol containing 5 mg/ml SnCl_2 and products were extracted with ethyl acetate [9]. They were analyzed by TLC in ethyl acetate/2,2,4-trimethylpentane/acetic acid/water, 55.25:10:50 (organic

phase, solvent I). When the plate was developed twice in the same solvent, excellent separation of products was obtained, with the following R_F values: 6-keto-PGF $_{1\alpha}$ (the stable end-product of PGI $_2$), 0.24; PGF $_{2\alpha}$, 0.37; PGE $_2$, 0.55; HHT, 0.90. Unlabelled compounds were used as internal (6-keto-PGF $_{1\alpha}$) or external (PGF $_{2\alpha}$, PGE $_2$ and HHT) references. In solvent II (ethyl acetate/2,2,4-trimethylpentane/water 5:10:10) all prostaglandins (methyl esters) remained close to the origin and HHT had an R_F of about 0.5.

The reference compounds were visualized by iodine vapor and the chromatograms were scanned for radioactivity with a Berthold Dünnschicht Scanner II. The radioactive zones were scraped off and counted in a Packard Tri-Carb liquid scintillation spectrometer model 3375.

3. Results

3.1. Subcellular distribution of PGI $_2$ synthase

Table 1 shows the conversion of PGH $_2$ by subcellular fractions of pig aorta.

Table 1
Conversion of PGH $_2$ by subcellular fractions of pig aorta – effect of boiling

	Product formation (% conversion)				
	6-Keto-PGF $_{1\alpha}$	PGF $_{2\alpha}$	PGE $_2$	PGD $_2$	HHT
Homogenate	72	3	12	3	10 (11)
Boiled homogenate	3	16	7	11	63 (62)
Mitochondria	7	29	28	13	21 (20)
Boiled mitochondria	4	13	45	21	17 (15)
Microsomes	35	17	19	6	22 (20)
Boiled microsomes	5	33	16	11	35 (36)
100 000 $\times g$ supernatant	10	25	26	11	28 (25)
Boiled supernatant	6	19	9	12	54 (56)
Buffer alone	3	9	55	24	9

Last column (in parentheses) shows HHT determinations by TLC in solvent II

All samples contained material corresponding to 0.5 g tissue in 1 ml 0.1 M potassium phosphate buffer, pH 7.4. Incubation was for 10 min, at 37°C with 5 μg PGH $_2$ (~30 000 cpm) and terminated with 5 ml ethanol containing 25 mg SnCl_2 . Products were extracted and analyzed as in section 2

PGI₂ forming activity was localized mainly in the microsomes as in [10]. However, the microsomes produced less PGI₂ (determined as 6-keto-PGF_{1α}), compared to an equivalent amount of homogenate, suggesting the presence in the latter of an activating or stabilizing factor. Table II shows that addition of supernatant to the microsomes stimulated PGI₂ formation. The factor responsible for this effect was not destroyed by boiling nor removed by dialysis (table 2).

3.2. Effects of denaturing agents on PGI₂ and HHT formation

Boiling (5–7 min at 100°C) destroyed the PGI₂

Table 2
Effect of supernatant on 6-keto-PGF_{1α} formation by aorta microsomes

Addition	Products	
	6-Keto-PGF _{1α} (%)	HHT (%)
None	32	23
Native supernatant (0.25 ml) ^a	62	17
Boiled supernatant (0.25 ml) ^a	54	34
Dialyzed supernatant (0.25 ml) ^a	47	19

^a Corresponding to 0.125 g tissue

Experimental details same as in table 1

forming activity in all preparations. Formation of HHT was, however, either not affected (mitochondria) or 2–6-fold enhanced (high-speed supernatant and homogenate, respectively). High production of HHT in those preparations contrasted with the small amount of HHT formed in buffer alone, under the same experimental conditions. Treatment with 6 M guanidine-HCl had similar effects as boiling: using microsomes, the formation of PGI₂ was abolished whereas HHT formation increased from 31–45%. In the supernatant guanidine-HCl raised the HHT formation from 31–66%.

3.3. Effects of inhibitors

Four inhibitors of thromboxane synthase were used in concentrations which inhibit TXB₂ formation by platelet microsomes by 80–90% [5]. Table 3 shows that none of these compounds appreciably influenced PGI₂ formation. On the other hand, 15-HPETE, an inhibitor of PGI₂ synthesis [10], prevented the formation of 6-keto-PGF_{1α}, but had no effect on HHT formation.

3.4. Partial purification of PGI₂ synthase

Aorta microsomes were treated with Triton X-100 as for platelet [3] and lung [4] microsomes. The 100 000 × g supernatant, which contained most of the PGI₂ forming activity, was chromatographed on a DEAE-cellulose column (table 4). Prostacyclin syn-

Table 3
Conversion of [1-¹⁴C]PGH₂ by aorta microsomes in the presence and absence of additions

Addition	Conversion of [1- ¹⁴ C]PGH ₂			
	6-Keto-PGF _{1α}		HHT	
	%	relative ^a	%	relative ^a
None	42	100	26	100
L-8027 (10 ⁻⁴ M)	35	84	23	88
N-0164 (2 × 10 ⁻⁴ M)	33	80	25	96
EMPA (10 ⁻⁴ M)	36	86	24	92
Imidazole (10 ⁻³ M)	37	89	23	88
15-HPETE (6 μg/ml)	6	14	24	92

^a Relative to the conversion by microsomes alone, taken as 100

Each sample contained 0.25 ml microsomal suspension (corresponding to 0.5 g tissue) in potassium phosphate buffer and the total volume was 0.5 ml. Other details as in table 1

Table 4
Product formation from [$1\text{-}^{14}\text{C}$]PGH₂ by aorta microsomal preparations

	Products	
	6-Keto-PGF _{1α} (%)	HHT (%)
Microsomal	49	24
Solubilized microsomes (supernatant)	43	28
Solubilized microsomes (sediment)	8	30
Fraction 1	5	18
Fraction 2	6	10
Fraction 3	37	29
Boiled fraction 3	4	32

Microsomal suspension 2 ml (corresponding to 4 g tissue) were solubilized with 0.5% Triton X-100 and the 100 000 \times g supernatant was applied to a DEAE-cellulose column (5 ml). Elution 30 ml 10 mM potassium phosphate buffer (pH 7.4), 0.1% Triton X-100 (fraction 1), 10 ml 20 mM buffer; 0.1% Triton X-100 (fraction 2) and 10 ml 0.2 M buffer, 0.1% Triton X-100 (fraction 3). Eluates were concentrated by ultrafiltration and portions corresponding to 0.5 g initial tissue were incubated with 3 μ g [$1\text{-}^{14}\text{C}$]PGH₂ (\sim 20 000 cpm) for 15 min at 37°C in a total volume of 0.5 ml. Products were extracted and analyzed as in section 2

thase was eluted from the column with 0.2 M buffer and almost no activity appeared in the preceding two fractions. Boiling of the partially purified enzyme (fraction 3) abolished 6-keto-PGF_{1 α} formation, but had no effect on HHT production.

4. Discussion

The two bioregulators, thromboxane and PGI₂, are formed from PGH₂ by the action of enzymes differently distributed in animal tissues [11]. While the thromboxane synthesizing enzyme systems in human platelets [3,5,6] and bovine lung [4] have been studied in some detail, less is known, so far, about the properties of PGI₂ synthase. The present results show that aorta microsomal PGI₂ synthase is stimulated by the high-speed supernatant. In contrast, lung thromboxane synthase was more active in isolated microsomes than in the unfractionated homogenate [4]. The effect of the supernatant on PGI₂ synthesis by pig aorta microsomes resembles the activation by supernatant of PGE₂ formation from arachidonic acid by sheep vesicular gland microsomes [in the absence of reduced glutathione, (unpublished observations)]. In either case, the soluble factor was resistant to boiling and non-dialyzable.

Four inhibitors of thromboxane synthase had almost no effect on PGI₂ formation by aorta microsomes. This indicates that these compounds can inhibit thromboxane formation selectively in systems which produce both thromboxanes and PGI₂. 15-HPETE inhibited PGI₂ formation, as has been reported [10] but had no effect on HHT formation. PGI₂ formation and HHT formation are therefore not coupled in the way that thromboxane and HHT formation are. In the latter case several structurally unrelated compounds inhibited TXB₂ and HHT formation identically [6]. HHT production from PGH₂ by aorta preparations was furthermore not prevented by boiling (see also [12]). Instead, its formation in certain fractions was enhanced by heating (table 1). In contrast to [12], we did not find much HHT formation from PGH₂ in buffer (table 1 and [4]), even when the incubation period was prolonged to 30 min (not shown). The major products were PGE₂ and PGD₂. It seems probable that a factor is present in the soluble fraction of the cell, which promotes HHT formation from PGH₂. When enzymes which utilize PGH₂ are denatured by heating, more PGH₂ becomes available for conversion to HHT. It has also been reported [13] that boiled homogenate of rat kidney has several-fold

increased HHT formation relative to the fresh homogenates which enzymatically form mostly PGE₂. Thus, thromboxane synthase is, so far, the only enzyme described which forms HHT.

Acknowledgements

This work was supported by a grant from the Swedish Medical Research Council (03X-217).

References

- [1] Hamberg, M., Svensson, J. and Samuelsson, B. (1975) *Proc. Natl. Acad. Sci. USA* 72, 2994–2998.
- [2] Moncada, S., Gryglewski, R., Bunting, S. and Vane, J. R. (1976) *Nature* 263, 663–665.
- [3] Hammarström, S. and Falardeau, P. (1977) *Proc. Natl. Acad. Sci. USA* 74, 3691–3695.
- [4] Wlodawer, P. and Hammarström, S. (1978) *Biochem. Res. Commun.* 80, 525–532.
- [5] Diczfalussy, U. and Hammarström, S. (1977) *FEBS Lett.* 82, 107–110.
- [6] Diczfalussy, U., Falardeau, P. and Hammarström, S. (1977) *FEBS Lett.* 84, 271–274.
- [7] Hamberg, M., Svensson, J., Wakabayashi, T. and Samuelsson, B. (1974) *Proc. Natl. Acad. Sci. USA* 71, 345–349.
- [8] Hamberg, M. and Samuelsson, B. (1974) *Proc. Natl. Acad. Sci. USA* 71, 3400–3404.
- [9] Hamberg, M. and Samuelsson, B. (1973) *Proc. Natl. Acad. Sci. USA* 70, 899–903.
- [10] Moncada, S., Gryglewski, R. J., Bunting, S. and Vane, J. R. (1976) *Prostaglandins* 12, 715–737.
- [11] Pace-Asciak, C. R. and Rangaraj, G. (1977) *Biochim. Biophys. Acta* 486, 579–582.
- [12] Skidgel, R. A. and Printz, M. P. (1978) *Prostaglandins* 16, 1–16.
- [13] Pace-Asciak, C. R. and Nashat, M. (1975) *Biochim. Biophys. Acta* 388, 243–253.