

Proteolysis of liver acetyl coenzyme A carboxylase by cathepsin B

Kenji Wada and Tadashi Tanabe*

Laboratory of Cell Biology, National Cardiovascular Center Research Institute, Fujishiro-dai, Suita, Osaka 565, Japan

Received 13 November 1984

Proteolysis of acetyl-CoA carboxylase was examined with cathepsin B. When chicken liver acetyl-CoA carboxylase was incubated with cathepsin B at pH 6.3, the native 220-kDa polypeptide was primarily cleaved into two polypeptides of 125 and 115 kDa, and further degraded to polypeptides of 100–50 kDa.

Protein degradation Acetyl-CoA carboxylase Cathepsin B

1. INTRODUCTION

Acetyl coenzyme A carboxylase (acetyl-CoA: carbon dioxide ligase (ADP-forming), EC 6.4.2.1) plays a central role in the regulation of the fatty acid biosynthesis [1,2]. The hepatic content of acetyl-CoA carboxylase depends on both the rates of synthesis and degradation of the enzyme [1,3]. Although the synthetic rate of acetyl-CoA carboxylase varies under different metabolic conditions, the degradation rate of this enzyme is hardly changed by metabolic perturbations except nutritional depletion [1,3]. Recent studies suggested that lysosomal thiol proteinases, including cathepsin B, H and L, play important roles in the degradation of endogenous proteins in the basal and nutritionally depleted states of cells [4–6]. The degradation mechanisms of individual proteins, however, have been poorly understood.

Our previous work has demonstrated that animal acetyl-CoA carboxylase is constituted of a unique multifunctional 220–230-kDa polypeptide [7,8]. It has also been shown that incubation of the native acetyl-CoA carboxylase isolated from rat liver with lysosomal extracts yields smaller polypeptides (100–130 kDa) [7]. Here, degradation of acetyl-CoA carboxylase was investigated

with the most abundant thiol proteinase, cathepsin B (EC 3.4.22.1).

2. MATERIALS AND METHODS

Acetyl-CoA carboxylase from chicken liver was purified as described previously [8]. The specific activity of the enzyme was 3.5 units/mg protein. Porcine kidney cathepsin B (type 1) was isolated according to [9]; the authentic sample was kindly donated by S. Takahashi of this institute. Specific activity of the enzyme was 4.3 units/mg protein as measured with a synthetic substrate, benzoyloxycarbonyl-phenylalanyl arginine-4-methyl-7-coumarylamide. Acetyl-CoA carboxylase (50 μ g) was incubated for 1 or 20 h with 5 μ g of cathepsin B in a reaction mixture (0.1 ml) containing 0.1 M potassium phosphate buffer, 5 mM cysteine and 2 mM EDTA, pH 6.3. The reaction was terminated by adding 25 μ l of 10 μ g/ml leupeptin. Aliquots (10 μ l) were removed from the reaction mixtures and treated with an equal volume of 2% SDS containing 8 M urea and 10% 2-mercaptoethanol (v/v). The mixture was heated at 100°C for 2 min and subjected to 7.5% polyacrylamide gel electrophoresis (PAGE) in the presence of 0.1% SDS as described previously [8]. The gel was stained by Coomassie brilliant blue R-250. Molecular masses of the polypeptides generated by

* To whom correspondence should be addressed

cathepsin B were determined with the standards from Bio-Rad, which are myosin heavy chain, β -galactosidase, phosphorylase *b*, bovine serum albumin and ovalbumin. Protein was determined by [10] with bovine serum albumin as the standard.

3. RESULTS AND DISCUSSION

The properties of cathepsin B are not greatly different among animal species [4]. Thus, the degradation of acetyl-CoA carboxylase from chicken liver was studied with porcine kidney cathepsin B. After the incubation of native acetyl-CoA carboxylase with cathepsin B for 1 h, the native 220-kDa polypeptide almost disappeared and two polypeptides (125 and 115 kDa) were newly formed (fig.1). The prolonged incubation for 20 h resulted in the formation of additional smaller polypeptides (100–50 kDa). Therefore, the initial steps of the catheptic degradation of acetyl-CoA carboxylase seem to proceed in a sequential manner. The primary degradation products of the 125- and 115-kDa polypeptides are quite similar to the polypeptides which have been found in the enzyme preparations purified at pH 7.0–7.5 in the absence of protease inhibitors [7,8,11,12]. These polypeptides found in the animal acetyl-CoA carboxylase preparations are likely due to the pro-

teolysis by cathepsin B, because cathepsin B exhibits a partial activity at neutral pH despite the pH optimum being 6.0–6.5, and additions of cathepsin B inhibitors, such as leupeptin and anti-pain [4,13] in the purification buffers markedly reduced the production of the smaller polypeptides [8].

Accumulated indirect evidence has suggested that cathepsin B may play an important role in the general catabolism of proteins in the lysosomal system [4]. Recently it has been clearly shown that 30–45% of the degradation of endogenous proteins in cultured macrophages and fibroblasts is inhibited by a cathepsin B inhibitor [5,6]. Furthermore, it has also been reported that the breakdown of pyruvate carboxylase in 3T3-L1 cells is inhibited by leupeptin [14]. Therefore, available evidence suggests the possibility that cathepsin B has a role in the initial steps of the degradation of acetyl-CoA carboxylase *in vivo*.

ACKNOWLEDGEMENTS

This investigation was supported in part by research grants from the Ministry of Education, Science, and Culture of Japan and Mitsui Life Social Welfare Foundation.

REFERENCES

- [1] Numa, S. and Tanabe, T. (1984) in: *Fatty Acid Metabolism and Its Regulation* (Numa, S. ed.) *New Comprehensive Biochemistry*, vol.7, pp.1–27, Elsevier, Amsterdam, New York.
- [2] Wakil, S.J., Stoops, J.K. and Joshi, V.C. (1983) *Annu. Rev. Biochem.* 52, 537–579.
- [3] Numa, S. and Yamashita, S. (1974) *Curr. Top. Cell. Regul.* 8, 197–246
- [4] Barrett, A.J. (1977) in: *Proteinases in Mammalian Cells and Tissues* (Barrett, A.J. ed) pp.181–208, North-Holland, Amsterdam.
- [5] Shaw, E. and Dean, R.T. (1980) *Biochem. J.* 186, 385–390.
- [6] Cockle, S.M. and Dean, R.T. (1982) *Biochem. J.* 208, 795–800.
- [7] Tanabe, T., Wada, K., Okazaki, T. and Numa, S. (1975) *Eur. J. Biochem.* 57, 15–24.
- [8] Wada, K. and Tanabe, T. (1983) *Eur. J. Biochem.* 135, 17–23.
- [9] Takahashi, S., Murakami, K. and Miyake, Y. (1981) *J. Biochem.* 90, 1677–1684.




Fig. 1. SDS-PAGE of acetyl-CoA carboxylase digests by cathepsin B. For experimental details see section 2. 4 μ g of acetyl-CoA carboxylase was run in each lane. Lane 1, marker proteins; lane 2, acetyl-CoA carboxylase without incubation, lane 3, acetyl-CoA carboxylase alone incubated for 20 h; lanes 4–6, the enzyme treated with cathepsin B for 0, 1 and 20 h, respectively.

- [10] Lowry, O.H., Rosebrough, N.J., Farr, A.L. and Randall, R.J. (1951) *J. Biol. Chem.* 193, 265–275
- [11] Inoue, H. and Lowenstein, J.M. (1972) *J. Biol. Chem.* 247, 4825–4832.
- [12] Guchhait, R.B., Zwergel, E.E. and Lane, M.D. (1974) *J. Biol. Chem.* 249, 4776–4780.
- [13] Umezawa, H. and Aoyagi, T. (1977) in: *Proteinases in Mammalian Cells and Tissues* (Barrett, A.J. ed.) pp 637–662, North-Holland, Amsterdam.
- [14] Chandler, C.S. and Ballard, F.J. (1983) *Biochem. J.* 210, 845–853