

# Diacylglycerol stimulates phospholipase A<sub>2</sub> from Swiss 3T3 fibroblasts

Ronald M. Burch

*Nova Pharmaceutical Corporation, 6200 Freeport Centre, Baltimore, MD 21224, USA*

Received 10 May 1988

We recently demonstrated that diacylglycerol induced arachidonate release and prostaglandin E<sub>2</sub> synthesis in 3T3 fibroblasts, and greatly augmented prostaglandin E<sub>2</sub> synthesis in response to submaximal and maximal concentrations of bradykinin. We have now partially purified a phospholipase A<sub>2</sub> from the cells. When phosphatidyl[<sup>3</sup>H]choline was used as substrate, several diacylglycerols augmented phospholipase A<sub>2</sub> activity. Diacylglycerol was effective at concentrations as low as 30 nM. Protein kinase C inhibition did not affect diacylglycerol's stimulation of phospholipase A<sub>2</sub>. Diacylglycerol did not alter the calcium requirement for phospholipase A<sub>2</sub> or its pH optimum. The present study demonstrates that the effect of diacylglycerol to augment arachidonate metabolism is at the level of phospholipase A<sub>2</sub>, itself.

Phospholipase A<sub>2</sub>; Arachidonic acid; Diacylglycerol; Protein kinase C; (Fibroblast)

## 1. INTRODUCTION

We have shown that bradykinin stimulates phospholipase A<sub>2</sub> (PLA<sub>2</sub>) activity in Swiss 3T3 fibroblasts to release arachidonate for prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) synthesis [1]. The primary substrate for the PLA<sub>2</sub> appeared to be phosphatidylcholine. We have also shown that in these cells diacylglycerols stimulate basal arachidonate release and PGE<sub>2</sub> synthesis, and that diacylglycerols markedly augment PGE<sub>2</sub> synthesis in response to bradykinin [2]. The site of diacylglycerol action was not defined, although protein kinase C (PKC) did not appear to be involved.

In the present study, we have used a partially purified membrane-bound PLA<sub>2</sub> from Swiss 3T3 fibroblasts to demonstrate that the effect of diacylglycerol is at the level of PLA<sub>2</sub>.

*Correspondence address:* R.M. Burch, Nova Pharmaceutical Corporation, 6200 Freeport Centre, Baltimore, MD 21224, USA

## 2. MATERIALS AND METHODS

### 2.1. Purification of PLA<sub>2</sub>

Swiss 3T3 fibroblasts were maintained as previously described [1]. Cells were harvested by scraping from the plates in phosphate-buffered saline and pelleted at 50000 × g for 10 min. The pellet was resuspended in 25 mM Tris-Cl (pH 7.4), 1 mM EGTA, 1 mM deoxycholate and 1 M KCl, and sonicated with a probe sonicator 4 times for 5 s each, with 30 s between each burst, in an ice bath. The lysate was incubated at room temperature for 30 min, then centrifuged at 200000 × g for 60 min. The supernatant was diluted in 100 vols of the same buffer without KCl, concentrated to 10 ml by ultrafiltration through an Amicon PM10 membrane, then applied to a DEAE-cellulose (Whatman DE-52) column (1 × 10 cm) that had been equilibrated with 25 mM Tris-Cl (pH 7.4), 1 mM EGTA, and 1 mM deoxycholate. The column was eluted with 25 ml of the same buffer followed by a linear gradient of 0–500 mM NaCl in the same buffer. Fractions of 2 ml were collected and PLA<sub>2</sub> activity was assayed at pH 7 and 9. A single peak of PLA<sub>2</sub> activity (pH 9) was eluted at 100–200 mM NaCl. No PLA<sub>2</sub> was eluted that was more active at pH 7 than at pH 9. This partially purified enzyme was dialyzed against 20 mM Tris, pH 7, and used within 2 days.

### 2.2. PLA<sub>2</sub> activity

Phosphatidyl[<sup>3</sup>H]choline was prepared by incubation of Swiss 3T3 fibroblasts with [<sup>3</sup>H]choline for 30–48 h. Phospholipids were extracted [3] and the lipids were subjected

to thin-layer chromatography [4]. The phosphatidylcholine fraction was scraped from the plate and was eluted with methanol, evaporated and stored in toluene/ethanol (50:50). Phosphatidylcholine concentration was estimated by measuring lipid phosphorus content [5].

For assay, a reaction mixture was prepared by adding labeled phosphatidylcholine to a 20 ml glass vial, with diacylglycerol where appropriate. The substrate was evaporated under  $N_2$ , then resuspended in 100 mM Tris, pH 9 (or other pH in selected experiments), containing 1 mM  $CaCl_2$  and 1 mM deoxycholate to a final phosphatidylcholine concentration of 10  $\mu M$ . The mixture was sonicated by probe 3 times for 5 s each, separated by 15 s intervals in an ice bath. For assay, 75  $\mu l$  of this mixture was added to 12  $\times$  75 mm glass tubes. The assay was begun by the addition of 25  $\mu l$  of enzyme preparation and subsequent incubation at 37°C. The reaction was linear for at least 90 min. The reactions were terminated by the addition of 200  $\mu l$  of methanol/chloroform/1 N HCl (2:1:0.01), followed by 100  $\mu l$  of 2 M KCl. The phospholipids were separated by thin-layer chromatography [4], and lysophosphatidylcholine was scraped from the plate, eluted with methanol and radioactivity was quantitated by liquid scintillation spectroscopy.

### 2.3. Materials

All lipids were obtained from Sigma. Phorbol myristate acetate was obtained from Calbiochem. H-7 was obtained from Seikagaku America.

## 3. RESULTS

### 3.1. Stimulation of partially purified phospholipase $A_2$ by diacylglycerols

When oleoylacetylgllycerol or dioleoylglycerol was present in the assay mixture,  $PLA_2$  activity was markedly augmented (fig.1). A variety of diacylglycerols were able to stimulate  $PLA_2$  activity (table 1). Among those tested, stearoylarachidonylglycerol has been shown to be a predominant diacylglycerol produced in cells [6]. Most of the diacylglycerols displayed similar effectiveness in stimulation of  $PLA_2$ . However, oleoylacetylgllycerol was consistently less effective than longer chain diacylglycerols (fig.1 and table 1).

The concentration-response for stimulation of  $PLA_2$  by dioleoylglycerol is shown in fig.2. At a concentration as low as 0.03  $\mu M$ , dioleoylglycerol significantly stimulated  $PLA_2$  activity. Other diacylglycerols displayed similar concentration-response patterns (not shown).

### 3.2. Role of protein kinase C in stimulation of $PLA_2$ by diacylglycerols

In our previous study [2], we demonstrated that diacylglycerols markedly augment arachidonate metabolism in Swiss 3T3 cells. While diacyl-




Fig.1. Oleoylacetylgllycerol (OAG) and dioleoylglycerol (diOG) stimulate  $PLA_2$  activity. Data are triplicate tubes assayed at pH 9 from 1 of 3 similar experiments.

glycerols activated PKC in the cells, we excluded this enzyme from playing a major role in stimulation of arachidonate metabolism, since neither inhibition nor down-regulation of PKC blocked diacylglycerols' effects. In our partially purified  $PLA_2$  preparation, under the conditions of the  $PLA_2$  assay, we detected no apparent PKC activity (table 2). However, if assay conditions were altered to be optimal for PKC, we did find evidence for protein kinase activity in the preparation (table 2). Thus, we tested the ability of the PKC inhibitor H-7 to affect diacylglycerol-stimulated  $PLA_2$  activity. Over a wide concentration range, H-7 did not affect diacylglycerol-stimulated  $PLA_2$  activity. In the presence of 100  $\mu M$  H-7, 4835  $\pm$  370 cpm lysoPC were formed, compared to 4886  $\pm$  322 cpm in the presence of dioleoylglycerol alone (1420  $\pm$  235 cpm in the absence of dioleoylglycerol). As a

Table 1

Comparison of the ability of several diacylglycerols to stimulate  $PLA_2$

	LysoPC (cpm)
Control	862 $\pm$ 94
Oleoylacetylgllycerol, 5 $\mu M$	2654 $\pm$ 285
Dioleoylglycerol, 5 $\mu M$	3562 $\pm$ 430
Dipalmitoylglycerol, 5 $\mu M$	3238 $\pm$ 264
Stearoylarachidonylglycerol, 5 $\mu M$	3764 $\pm$ 386

Incubations were at pH 9 for 60 min at 37°C. Data are from 1 of 2 similar experiments

Table 2

Protein kinase C activity in the partially purified PLA<sub>2</sub> preparation

	cpm <sup>32</sup> P into histone
Control	24 ± 20
OAG, 5 μM	30 ± 14
PKC assay buffer + OAG	96 ± 24
Initial homogenate + OAG	370 ± 26

Mean ± SE of triplicate determinations in 1 of 2 similar experiments. Incorporation of <sup>32</sup>P from [ $\gamma$ -<sup>32</sup>P]ATP into histone (type III-S) was assayed in the PLA<sub>2</sub> assay buffer plus radioactive ATP or in PKC assay buffer (400 μg/ml phosphatidylserine, 1 mM CaCl<sub>2</sub>, and 5 μM oleoylacetyl-glycerol (OAG) for 2 min) [14]. Homogenates of 3T3 cells were also assayed in PKC assay buffer and adjusted to the same amount of protein added (10 μg) as the partially purified PLA<sub>2</sub> preparation

final test of whether PKC might mediate diacylglycerols' ability to stimulate PLA<sub>2</sub>, we prepared membranes from Swiss 3T3 cells that had been pretreated with 1 μM phorbol myristate acetate for 48 h, a treatment that completely down-regulates PKC [2]. Down-regulation of PKC had no effect on the ability of diacylglycerols to stimulate PLA<sub>2</sub> activity, since after down-regulation 3260 ± 386 cpm lysoPC was formed, compared to 3654 ±




Fig.2. Concentration-response for stimulation of PLA<sub>2</sub> by dioleoylglycerol (diOG). Incubations were for 60 min at 37°C. Data are from 1 of 3 similar experiments. At 0.03 μM, dioleoylglycerol significantly stimulated PLA<sub>2</sub> ( $p < 0.05$ ) when all 3 experiments were compared by Student's *t*-test for paired observations.




Fig.3. Effect of calcium on the ability of dioleoylglycerol (diOG) to stimulate PLA<sub>2</sub>. The ordinate is the amount of calcium added to the assay buffer. Incubations were for 60 min at 37°C. Data are from 1 of 2 similar experiments.

270 in the presence of dioleoylglycerol alone (control 1128 ± 145 cpm).

### 3.3. Role of calcium in the ability of diacylglycerol to stimulate PLA<sub>2</sub>

Since most PLA<sub>2</sub>s require calcium for activity, we tested the effects of altered calcium concentration on diacylglycerol-stimulated PLA<sub>2</sub> activity. In the absence of added calcium, diacylglycerol did not stimulate PLA<sub>2</sub> activity (fig.3).

## 4. DISCUSSION

The present study extends our previous observation that diacylglycerol stimulates arachidonate metabolism in Swiss 3T3 cells. We have now shown that diacylglycerol acts at the level of PLA<sub>2</sub> itself, markedly stimulating activity of a PLA<sub>2</sub> partially purified from Swiss 3T3 cells.

A variety of diacylglycerols were able to stimulate PLA<sub>2</sub> activity. However, oleoylacetyl-glycerol was consistently less effective than longer-chain diacylglycerols. It has been previously shown that diacylglycerols cause restructuring of phospholipid vesicles [7]. This may allow PLA<sub>2</sub> more ready access to its substrate.

It is particularly noteworthy that stearyl-arachidonoylglycerol, a diacylglycerol that is formed in response to activation of a number of receptors [6], stimulated PLA<sub>2</sub> activity. We have no information about the mass of diacylglycerol formed in

Swiss 3T3 cells in response to agonists. In platelets, however, diacylglycerol (4–11 nmol/10<sup>9</sup> platelets) is generated in response to receptor activation [8], amounting to perhaps 1–2 mol% of total phospholipid. In our experiments, 0.3 mol% diacylglycerol significantly stimulated PLA<sub>2</sub>. Thus, formation of diacylglycerol in response to receptor activation may be expected to stimulate PLA<sub>2</sub>.

In our previous study utilizing intact Swiss 3T3 cells, diacylglycerol stimulation of arachidonate metabolism appeared to be independent of PKC. Under the PLA<sub>2</sub> assay conditions of the present study, PKC activity could not be detected. Further evidence against participation of PKC were the findings that neither PKC inhibition nor down-regulation affected diacylglycerol stimulation of PLA<sub>2</sub> activity. Also, experiments showed that in the absence of added calcium, PLA<sub>2</sub> was not activated. In the absence of added calcium, free calcium in the buffer was likely several micromolar [9]. This concentration would be adequate to fully activate PKC in the presence of diacylglycerol.

The mechanism by which diacylglycerol stimulated PLA<sub>2</sub> activity is unclear. Diacylglycerol did not affect the calcium requirement of the enzyme nor its pH optimum. The mechanism may relate to the ability of diacylglycerol to alter the structural organization of the substrate. It has been suggested that PLA<sub>2</sub> and PKC may contain similar regulatory sequences [10]. Thus, similar to its effect on PKC [11], diacylglycerol may directly bind to PLA<sub>2</sub> and activate it. Testing this possibility awaits further purification of PLA<sub>2</sub>.

Similar ability of diacylglycerol to stimulate phospholipases has been observed for purified phosphatidylinositol-specific phospholipase C [12], and other PLA<sub>2</sub>s [13]. However, this study correlated diacylglycerol activation of PLA<sub>2</sub> to a

biological effect, potentiation of PGE<sub>2</sub> synthesis in response to receptor agonists [2]. This action of diacylglycerol may be physiologically relevant in that diacylglycerol, newly synthesized in response to a variety of agonists may activate PLA<sub>2</sub>, leading to arachidonate release and metabolism, or other biochemical events mediated by PLA<sub>2</sub> metabolites.

*Acknowledgements:* I would like to thank Julius Axelrod and Stephen G. Farmer for critically reading the manuscript and Cheryl Sowards for preparing the final manuscript.

## REFERENCES

- [1] Burch, R.M. and Axelrod, J. (1987) Proc. Natl. Acad. Sci. USA 84, 6374–6378.
- [2] Burch, R.M., Ma, A.L. and Axelrod, J. (1988) J. Biol. Chem., in press.
- [3] Bligh, E.G. and Dyer, W.J. (1959) Can. J. Biochem. Physiol. 37, 911–917.
- [4] Billah, M.M., Lapetina, E.G. and Cuatrecasas, P. (1980) J. Biol. Chem. 255, 10227–10231.
- [5] Bartlett, G.R. (1959) J. Biol. Chem. 234, 466–468.
- [6] Prescott, S.M. and Majerus, P.W. (1981) J. Biol. Chem. 256, 579–582.
- [7] Das, R. and Rand, R.P. (1986) Biochemistry 25, 2882–2889.
- [8] Rittenhouse-Simmons, S. and Deykin, D. (1981) in: Platelets in Biology and Pathology 2 (Gordon, J.L. ed.) pp.349–372, Elsevier, Amsterdam.
- [9] Burch, R.M., Wise, W.C. and Halusha, P.V. (1983) J. Pharmacol. Exp. Ther. 227, 84–91.
- [10] Maraganore, J.M. (1987) Trends Biochem. Sci. 12, 176–177.
- [11] Nishizuka, Y. (1984) Science 225, 1365–1370.
- [12] Hofmann, S.L. and Majerus, P.W. (1982) J. Biol. Chem. 257, 14359–14364.
- [13] Dawson, R.M.C., Hemington, N.L. and Irvine, R.F. (1983) Biochem. Biophys. Res. Commun. 117, 196–201.
- [14] Rodriguez-Pena, A. and Rozengurt, E. (1984) Biochem. Biophys. Res. Commun. 120, 1053–1059.