

Partial sequence of the rat heavy neurofilament polypeptide (NF-H)

Identification of putative phosphorylation sites

Kieran C. Breen, Philip A. Robinson*, Didier Wion[†] and Brian H. Anderton

Dept of Immunology, St. Georges Hospital Medical School, Cranmer Terrace, London SW17 0RE, England

Received 29 September 1988

A 3 kb cDNA clone has previously been isolated in this laboratory corresponding to the rat heavy neurofilament polypeptide (NF-H). This clone, equivalent to approximately 70% of the total mRNA of the protein has been sequenced and shown to contain the carboxy-terminal region of the message. This contains 51 of the Lys-Ser-Pro repeat triplets which are reported to be the site of neurofilament phosphorylation. The sequence obtained was subsequently compared to those of mouse and human NF-H, showing a homology of approximately 85%. There is, however, one region which is variable between the species, this being the highly phosphorylated region of the protein containing the Lys-Ser-Pro triplet repeat.

Neurofilament; cDNA sequence; Phosphorylation site; Interspecies comparison; (Rat brain)

1. INTRODUCTION

Neurofilaments (NF), the neuronal intermediate filaments, are composed of three polypeptides of apparent molecular mass, determined by SDS-PAGE, approx. 70 kDa (NF-L), 160 kDa (NF-M) and 200 kDa (NF-H) and are involved in the maintenance of neuronal caliber [1]. Phosphorylation probably plays a major role in the functioning of the two larger neurofilament polypeptides (NF-M and NF-H), the levels of phosphorylation being altered developmentally and coincident with a change in neurofilament function [2,3]. Both NF-M and NF-H contain a heavily phosphorylated carboxy region which may influence the formation of filament-filament axonal cross bridges [4,5].

The NF-L and NF-M genes and cDNAs have

been cloned and sequenced [6–11] and also those of the mouse and human NF-H [12,13]. More recently, a partial cDNA clone for rat NF-H has been sequenced [14]. We have previously reported the isolation of a partial cDNA clone for rat NF-H [15] which extends over 200 bases further towards the N-terminal and here we describe the sequence of the clone which corresponds to approx. 70% of the total message. The clone contains the triplet repeat Lys-Ser-Pro sequence which is present in the carboxy-terminal region and is repeated 51 times, this being the putative phosphorylation region of the protein. The sequence of the rat triplet repeat region is then compared to those of human and mouse NF-H.

2. MATERIALS AND METHODS

2.1. Preparation and screening of a rat brain cDNA library

The cDNA library was prepared and screened as previously described [15] yielding a clone of 3 kb in length.

2.2. DNA sequencing

The clone obtained was inserted into the Bluescript expression vector. Deletion mutants were generated using the Exo III/mung bean nuclease system (Stratagene) and these were sequenced by the dideoxy chain termination method [16].

Correspondence address: K.C. Breen, Dept of Immunology, St. Georges Hospital Medical School, Cranmer Terrace, London SW17 0RE, England

* Present address: Dept of Dental Surgery, School of Dentistry, University of Leeds, Leeds LS2 9LU, England

[†] Present address: INSERM U298, Faculté de Médecine, Rue Haute de Reculée, 49045 Angers, Cedex, France

1 CAC CAG GAG GAG GTG GGC GAG CTG CTC GGT CAG ATT CAG GGC TGC GGT GCC GCG CAG GCG CAG GCT CAG GCC GAG GCT CGG GAC GCC CTC AAC TCC 96
 His Glu Glu Val Glu Glu Leu Leu Gly Glu Glu Ile Glu Glu Ala Ala Glu Ala Glu Ala Arg Asp Ala Leu Lys Cys
 97 GAC GTG ACG TCG GCG CTG CGG GAG ATC CGC CGG CAG CTC GAA GGA CAC ACC GTG CAG ACT ACC CTG CAG TCA GAG GAG TGG TTC CGA GTG AGA TTG 192
 Asp Val Thr Ser Ala Leu Arg Glu Ile Arg Ala Glu Leu Glu Gly His Thr Val Glu Ser Thr Leu Glu Ser Glu Glu Trp Phe Arg Val Arg Leu
 193 GAC CGA CTC TCA GAG GCA GCC AAA GTG AAC CAG CAT GCT ATT CCC TCT GGC CAA GAG GAG ATA ACT GAG TAC CGG CCG CGC CTG CAG GCG ACC 288
 Asp Arg Leu Ser Glu Ala Ala Lys Val Asn Thr Asp Ala Met Arg Ser Ala Glu Glu Ile Thr Glu Tyr Arg Arg Glu Leu Glu Ala Arg Thr
 289 ACA GAG TTG GAG GCA CTG AAA AGC ACC AAG GAG TCA CTG GAG AGG CAG CGC TCT GAG GAG GAC CGT CAT CAG GTA GAC ATG GCC TCC TAC CAG 384
 Thr Glu Leu Glu Ala Leu Lys Ser Thr Lys Glu Ser Leu Glu Arg Glu Asn Arg Ser Glu Leu Glu Arg His Glu Val Asp Arg Met Ala Ser Tyr Glu
 385 GAT GCA ATT CAG CAG CTG GAC AAT GAG CTG AGA AAC ACC AAA TGG GAG ATG GCC CGG CAG CTC CGA GAG TAC CAG GAC CTC CTC AAC GTC AAG ATG 480
 Asp Ala Ile Glu Leu Asp Arg Asn Glu Leu Arg Asn Thr Lys Trp Glu Met Ala Glu Leu Arg Glu Tyr Glu Asp Leu Leu Asn Val Lys Met
 481 GCC CTG GAT CTT GAG ATC GCT TAC AGA AAA CTC CTG GAA GGC GAA GAG TGT CGG ATT GGC TTT GGA CCC ATT CCC TTC TCT CTT ACT GAG GGA 576
 Ala Leu Asp Leu Glu Ile Ala Asp Tyr Arg Lys Leu Leu Glu Gly Glu Cys Arg Ile Glu Phe Gly Pro Ile Pro Phe Ser Leu Thr Glu Gly
 577 CTC CCA AAA ATT CCC TCC ATG TCC ACT AAC ATA AAA GTG AAA GCA GAA GAG AAC TGA ATA AAA GTA GCA GAA AAA TCG GAG AAG GAA ACC GTC ATT GTC 672
 Leu Pro Lys Ile Pro Ser Met Ser Thr His Ile Lys Val Lys Ser Glu Glu Lys Ile Val Val Glu Lys Ser Glu Lys Glu Thr Val Ile Val
 673 GAG GAA CAG ACA GAA GAG ATC CAG GTG ACA GAA GAG GAG GAC AAA GAG GCC CAA GGG GAG GAA GAA GAG GCA GAA GAG GGA 768
 Glu Glu Glu Glu Thr Glu Glu Ile Glu Val Thr Glu Glu Glu Asp Lys Glu Ala Glu Glu Glu Ala Glu Glu Glu Ala Glu Glu
 769 GGA GAA GAA GCA GCA ACT ACG TCT CCC CCT GCA GAA GAG GCT GCA TCT GCA GAA AAC GAA GCA ACC AAC AAG TCT CCT GTG AAA GAA GAG GCC AAG TCC CCA 864
 Gly Glu Glu Ala Glu Ala Thr Thr Ser Pro Ala Glu Ala Asp Ser Pro Glu Lys Glu Thr Lys Ser Pro Val Lys Glu Ala Lys Ser Pro
 865 GCT GAG GCC AAG TCC CCA GCT GAG GCC AAC TCA CCA GCT CAG GCC AAC TCC CCA GCT GAG GTC AAA TCT CCA GCT GTG GCC AAC TCC CCA GCT GAA 960
 Ala Glu Ala Lys Ser Pro Ala Glu Ala Lys Ser Pro Ala Glu Ala Lys Ser Pro Ala Glu Val Lys Ser Pro Ala Val Ala Lys Ser Pro Ala Glu
 961 GTC AAA TCT CCA GCT GTC AAA TCT CCA GCT GAG GCC AAC TCA CCA GCT GAG GCC AAC TCA CCA GCT GAG GTC AAC TCT CCA GCT ACA GTC AAG 1056
 Val Lys Ser Pro Ala Glu Val Lys Ser Pro Ala Glu Ala Lys Ser Pro Ala Glu Ala Lys Ser Pro Ala Glu Val Asn Ser Pro Ala Lys Ser Pro
 1057 TCT CCA GGT GAG GGC AAG TCC CCA GCT GAG TCA CCA GCT GAG CTC AAA TCT CCA GCT GAG GCC AAC TCA CCA GCT GAG GGC AAG TCT CCA 1152
 Ser Pro Glu Ala Lys Ser Pro Ala Glu Ala Lys Ser Pro Ala Glu Val Lys Ser Pro Val Glu Ala Lys Ser Pro Ala Glu Ala Lys Ser Pro
 1153 GCT TCA GTG AAG TCC CCA GGT GAG GCC AAG TCA CCA GCT GAG GCC AAC TCA CCA GCT GAG GTC AAC TCT CCA GCT ACA GTC AAG TCC CCA GTT GAG 1248
 Ala Ser Val Lys Ser Pro Glu Ala Lys Ser Pro Ala Glu Ala Lys Ser Pro Ala Glu Val Lys Ser Pro Ala Glu Val Lys Ser Pro Ala Val Glu
 1249 GCC AAG TCA CCA GCT GAG GTC AAA TCT CCA GTT ACA GTG AAG TCC CCA GCT GAG GCC AAC TCA CCA GCT GAG GTC AAA TCT CCA GCT TCG GAG 1344
 Ala Lys Ser Pro Ala Glu Val Lys Ser Pro Ala Glu Ala Lys Ser Pro Ala Glu Ala Lys Ser Pro Ala Glu Ala Lys Ser Pro Ala Val Lys
 1345 TCC CCA AGT GAA GCC AAG TCA CCT GCT GGA GCC AAC TCA CCA GCT GAG GCC AAC TCA CCA GTT GTG GCC AAA TCA CCA GCT GAG GCC AAC TCA CCA 1440
 Ser Pro Ser Glu Ala Lys Ser Pro Ala Glu Ala Lys Ser Pro Ala Glu Ala Lys Ser Pro Val Val Ala Lys Ser Pro Ala Glu Ala Lys Ser Pro
 1441 GCT GGA GCC AAG CCT CCA GCT GAG CCC AAC TCA CCA GCT GAG GCC AAC TCA CCA GCT GAG GTC AAC TCT CCA GCT GAG GCC AAC TCA CCA CCT GAG 1536
 Ala Glu Ala Lys Pro Pro Ala Glu Ala Lys Ser Pro Ala Glu
 1537 GCC AAG TCA CCT GTT GAG GTC AAA TCT CCA GAG AAC GCC AAC CCC GTG AAG GAA GAG GGT GCA AAA TCC CTA GCT GAG GCC AAC TCC CCT GAG 1632
 Ala Lys Ser Pro Val Glu Val Lys Ser Pro Glu Lys Ala Lys Ser Pro Val Lys Glu Gly Ala Lys Ser Leu Ala Glu Ala Lys Ser Pro Glu Lys
 1633 GCC AAG TCC CCT GTG AAG GAA GAG ATC GCA GCT GAG AAC TCC CCC GAG AAC GGC AAC CCC ATG AGG AAC GAG GGC AAC TCT CCT 1726
 Ala Lys Ser Pro Val Lys Glu Glu Ile Lys Pro Pro Ala Glu Val Lys Ser Pro Glu Lys Ala Lys Ser Pro Met Arg Lys Glu Ala Lys Ser Pro
 1729 GAG AAG GCC AAG ACT CTG GAT GTG AAG TCT CCA GAA GCC AAC CCT CCA GCG AAC GAG GAA GCA AAC GGC CCC GCA GAC ATC AGA TCC CCT GAG CAG 1824
 Glu Lys Ala Lys Thr Leu Asp Val Lys Ser Pro Glu Ala Lys Pro Pro Ala Lys Glu Glu Ala Lys Arg Pro Ala Asp Ile Arg Ser Pro Glu Glu
 1825 GTC AAA AGT CCT GCC AAG GAG GAG CCC AAC TCC CCC GAG AAC GAA GAG ACC AGG ACT GCA AAC GAG GTC GGT CCC AAC AGG AAC GAA GAG GTG AAG TCC CCT 1920
 Val Lys Ser Pro Ala Lys Glu Glu Ala Lys Ser Pro Glu Lys Glu Glu Thr Arg Thr Glu Lys Val Lys Ala Pro Lys Glu Glu Val Lys Ser Pro
 1921 GTG GAG GAG GTC AAA GCC AAA GCA GGC AAC GAG GTG GAG GAG AAC ACA CCA GGC ACA AAC ACA GAG GTG AAC GAG AGC AAC GAT 2016
 Val Glu Glu Val Lys Ala Lys Pro Pro Lys Glu Val Glu Glu Lys Thr Pro Ala Thr Pro Lys Thr Pro Ala Thr Pro Lys Glu Ser Lys Lys Asp
 2017 GAA GCT CCC AAG GAG GCC CAG AAC CCC AAC GCG GAG AAC GAG GAG CCT CTC ACA GAA AAC CCC AAC GAG GAC TCT CCT CGG GGG GAA GCC AAC ARG GAA GAG 2112
 Glu Ala Pro Lys Glu Ala Glu Lys Pro Lys Glu Ala Glu Lys Pro Lys Glu Leu Thr Glu Lys Pro Lys Asp Ser Pro Glu Glu Ala Lys Lys Glu
 2113 GCT AAA GAG AAC GGC GCG CCA GAG GAG AAC CCC GGC AAC TGT GGC GTG AAC GAA GAG GCT AAA CCC AAA GAG AAC GCA GAA GAC GCC 2208
 Ala Lys Glu Lys Ala Ala Pro Glu Glu Glu Thr Pro Ala Lys Leu Glu Val Lys Glu Glu Ala Lys Pro Lys Glu Lys Ala Glu Asp Ala
 2209 AAC GCC AAA GAA CCT AGC AAA CCC TCA GAG AAC GAG AAA CCG AAC AAC GAG GAG GAG GTG CCG GCA GCA GCA GAG AAC AAA GAC ACC AAC GAG GAG AAG 2304
 Lys Ala Lys Glu Pro Ser Lys Pro Glu Lys Glu Lys Pro Lys Glu Lys Pro Lys Glu Lys Asp Ser Thr Pro Lys Glu Lys Asp Thr Lys Glu Lys
 2305 ACT ACG GAG TCC AAG CGT GAG AAA CCC AAA ATG GAG GCA AGG CCA AGG AGG GCA AGG ACA AGG GCC TTC CCC CCC AGG AGC CTA GCA AAC CCA AGA 2400
 Thr Thr Glu Ser Lys Lys Arg Lys Pro Lys Glu Met Glu Pro Arg Pro Arg Arg Arg Arg Ala Pro Lys Ser Leu Ala Asn Asn Pro Arg
 2401 CAG AAA AGG CGT AAA AGT CCT CTA GCA CAG ACC AAA AAC GCA GGC AGC CCT CAG AGA AGG CCC CAG AGG ACA AGG CTG CCA AGG GAG ACA AGT AAG 2496
 Glu Lys Arg Leu Lys Ser Pro Leu Ala Glu Ile Lys Lys Glu Asp Gly Pro Glu Gly The Glu Gly Val Val *
 2497 AGG ACG AGA GGG ACA CCC AGA ATA GCA AAA GAA ACT CAG GAC GGC CCC GGT ACT CAA GGG TTG CTG TAA TAA AGT TTA TTT CCT CCT CCT CCG 2592
 Arg Thr Arg Gly Thr Pro Arg Ile Lys Lys Glu Asp Gly Pro Glu Gly The Glu Gly Val Val *
 2593 TAA GAA GAA ACA CCA CCT AGA TTG CGG GCC CGC CCT CAC CAA ACA GGA TTT CTA TTA GGA TTA AGT TAG CAA GAG AAC ATC AAC CCT GAG CCC TCC 2688
 2689 CTC CCC AAC ACC AAA GCC CTC CCC AAC TTG ATG GAC ATT TAT GAT AGC TTA TCG TAG CGG AAC GGA GAT GTC TTG CTG AAC GCT CCA CGT AAA AAC 2784
 2785 CGT GAC TAA AAA CTG CCC CTC CCT TCC RAG TAA GTG CAT TCA CCT CCC GTC TGT CCT ACC GAC AGG TCA CGG CAG TAA TGA ATG ACC AGT TAG 2880
 2881 AAA TGC ATT ATG CCT GAA ATG TTG TAA CCT ATT CCC GAA TGC CCT TTG TTG CAA AGG AGC GGT CAG GCC CTT GCC CGG TAC ACC CTC CTC GAA 2976
 2977 GAG CTG CAG CAG CTG AGG CAG GGC GCT GGC TGA ACC AGG CCA GGG TGT GCT GTC CAC TGA AGG CCA CTT TCG ATT GCT TCC GTG CAA TAA AAC 3072
 3073 CCA ACT GCT TCT GA{41} 3086

Fig.1. Nucleotide sequence of the 3 kb clone of rat NF-H, and the corresponding deduced amino acid sequence. The untranslated region is 529 bases long and includes the poly(A)⁺ tail.

3. RESULTS AND DISCUSSION

The 3 kb clone obtained begins at the end of α -helix coil 1B and continues through the carboxy-terminal (stop codon) to the poly(A) tail (fig.1). The amino acid sequence of coil 2 was found to be very highly conserved with respect to both rat NF-L and NF-M and also mouse and human NF-H, demonstrating a homology of approx. 80% and 90%, respectively [6,11–13]. This is in good agreement with previous results which have shown a high degree of conservation between the coiled regions of intermediate filaments from a number of different species [4–6]. There is also a good homology between the three species in the last 200 amino acids or so at the carboxy-terminal of the NF-H protein.

In between these two regions, there is a section containing a series of 51 repeated amino acid triplets: Lys-Ser-Pro. These occur largely in a sequence of 6 amino acids, in which the Lys-Ser-Pro is highly conserved in positions 3–5 of the sextet (fig.2). The amino acid in position 2, preceding the triplet, is also highly conserved being either Ala or Val. There is, however, a greater divergence in the first and sixth amino acids of the sextet, but this difference is unlikely to influence greatly the secondary structure of the region, this probably being determined by the triplet repeat sequence. The Lys-Ser-Pro sequence has previously been identified as the phosphorylation site within NF-H [18] and has also been recognized in other neurofilament proteins as well as in neurofilament and microtubule-associated proteins including

	<u>RAT</u>	<u>MOUSE</u>	<u>HUMAN</u>
Glu-Ala-Lys-Ser-Pro-Ala	21	15	6
Glu-Val-Lys-Ser-Pro-Ala	4	1	—
Glu-Ala-Lys-Ser-Pro-Val	3	2	2
Glu-Val-Lys-Ser-Pro-Val	3	1	1
Val-Ala-Lys-Ser-Pro-Ala	2	—	—
Glu-Ala-Lys-Ser-Pro-Ala	2	—	—
Glu-Val-Lys-Ser-Pro-Ser	2	—	—
Glu-Val-Lys-Ser-Pro-Glu	2	—	4
Glu-Ala-Lys-Ser-Pro-Glu	2	—	10
Glu-Thr-Lys-Ser-Pro-Val	1	1	—
Thr-Val-Lys-Ser-Pro-Gly	1	1	—
Ser-Val-Lys-Ser-Pro-Val	1	—	—
Thr-Val-Lys-Ser-Pro-Val	1	—	—
Thr-Val-Lys-Ser-Pro-Ala	1	—	—
Gly-Ala-Lys-Ser-Pro-Ala	1	1	—
Gly-Ala-Lys-Ser-Pro-Glu	1	1	—
Lys-Ala-Lys-Ser-Pro-Met	1	—	—
Asp-Val-Lys-Ser-Pro-Glu	1	1	—
Gln-Val-Lys-Ser-Pro-Ala	1	—	—
Ala-Val-Lys-Ser-Pro-Gly	—	1	—
Glu-Ala-Lys-Ser-Pro-Ile	—	1	—
Gln-Val-Lys-Ser-Pro-Glu	—	1	—
Glu-Ala-Lys-Ser-Pro-Gly	—	7	—
Glu-Ala-Lys-Ser-Pro-Ser	—	1	—
Glu-Pro-Lys-Ser-Pro-Ala	—	3	—
Glu-Thr-Lys-Ser-Pro-Pro	—	—	1
Lys-Ala-Lys-Ser-Pro-Ala	—	—	2
Glu-Ala-Lys-Ser-Pro-Pro	—	—	1
Lys-Ala-Lys-Ser-Pro-Val	—	—	8
Lys-Ala-Lys-Ser-Pro-Thr	—	—	1
Lys-Ala-Lys-Ser-Pro-Glu	—	1	3
Lys-Ala-Lys-Ser-Pro-Leu	—	—	1
Glu-Val-Lys-Ser-Pro-Gly	—	3	—
Ala-Val-Lys-Ser-Pro-Ala	—	1	—
TOTAL	51	43	41

Fig.2. Sequence of the amino acid sextets containing the Lys-Ser-Pro triplet repeat of NF-H of rat, mouse [12] and human [13].

MAP-2 and Tau, indicating that it too, is highly conserved [17].

A small number of triplet repeats are also present in NF-M [9–11,18], but are absent from NF-L [6–8]. Previous studies have shown that the number of alkaline phosphatase-sensitive phosphate groups on NF-H and NF-M are approximately proportional to the number of triplet repeats, these phosphate groups being absent from NF-L [19]. This conforms with the proposal that

the triplet is indeed the site of neurofilament phosphorylation.

There is however, a difference between both the total number of triplet repeats present and the intervening amino acids in the heavy neurofilament protein from rat, mouse and human. The mouse and human NF-H messages are fairly similar with 43 and 40 repeat sequences, respectively, whereas the rat NF-H has 51 repeats (fig.3). A similar interspecies difference has also previously been

R	- - - - - T S P P A E E A A S P E K E T K S P V K E E A K S P A E A K S P
H	G G E E E T K S P P A E E A A S P E K E A K S P V K E E A K S P A E A K S P
M	E L A A A - T S P P A E E A A S P E K E T K S P V K E E A K S P G E A K S P
R	A - - E A K S P A E A K S P A E V K S P A - - V A K S P A E V K S P A - - E
H	E K E E A K S P A E V K S P E K A K S P A K E E A K S P P P E A K S P E K E E
M	G - - E A K S P A E A K S P G E A K S P G - - E A K S P G E A K S P A - - E
R	V K S P A E A K S P A E A K S P A - - E A K S P A T V K S P G E A K S P A -
H	A K S P A E V K S P E K A K S P A K E E A K S P A E A K S P E K A K S P V K
M	P K S P A E P K S P A E A K S P A - - E P K S P A T V K S P G E A K S P S -
R	- E A K S P A E V K S P V - - E A K S P A E A K S P A S V K S P V - - E A K
H	E E A K S P A E A K S P V K E E A K S P A E V K S P E K A K S P T K E & A K
M	- E A K S P A E A K S P A - - E A K S P A E A K S P A E V K S P G - - E A K
R	S P A E A K S P A - - E V K S P S T V K S P V - - E A K S P A E V K S P V -
H	S P E K A K S P E K E E A K S P E K A K S P V K A E A K S P E K A K S P V K
M	S P A E P K S P A - - E A K S P A E V K S P A - - E A K S P A E V K S P G -
R	- T V K S P A E A K S P V - - E V K S P A S V K S P S - - E A K S P A G A K
H	A E A K S P E K A K S P V K E E A K S P E K A K S P V K E E A K S P E K A K
M	- E A K S P A A V K S P A - - E A K S P A A V K S P G - - E A K S P G E A K
R	S P A - - E A K S P V V A K S P A - - E A K S P A E A K S P A E A K S P A E
H	S P V K E E A K T P E K A K S P V K E E A K S P E K A K S P E K A K T L D V
M	S P A - - E A K S P A E A K S P I - - E V K S P G G A K T P V - - - - -
R	A K S P A E A K S P A E A K S P A E A K S P V E V K S P E K A K S P V K E G
H	K - E E
M	- E E
R	- A K S P E K A K S P V K E E I K P P A E V K S P E K A - - - - - K S P
H	- - K S P E - A K T P A K E E A - - - - - R S P A D K F F E K A - - K S P
M	G A K S P A G A K S P - - E E A - - - - - K S P V E E D I P P A E A K S P
R	M R K E A K S P E K A K T L D V - - - - - K S P E A K P P A K E E A K
H	V K E E V K S P E K A - - - - - K S P L K A D A K A P E K E
M	- G E A - K S P V K E G A K P P E K A K P L D V K S P E A Q T P V Q E E A T
R	A P A D I R S P E Q V K S P A K E E A K S P E K E E T R - T E K V A P K K E
H	I P K - K E
M	Y P T D I R P P E Q V K S P A K E K A K S P E K E E A K T S E K V A P K K E
R	E V K S P V
H	E V K S P V
M	E V K S P V

Fig.3. Comparison of the amino acid sequences of the repeat region of rat NF-H, human NF-H [13], and mouse NF-H [12]. Deletions (dashes) allow for better alignment of the sequences.

noted in the corresponding region of NF-M [4,9–11].

In the rat NF-H cDNA sequence, 41 of the triplet Lys-Ser-Pro repeats occur in a regular pattern of six amino acids (fig.3). In the mouse, also, this regular pattern was seen with 33 of the 43 triplet repeats present. In the human, however, it can be noted that the spacing of the triplets is slightly altered with them being separated alternately by three and five amino acids. There is also a considerable difference between the amino acid composition of the repeat region between the three species. While Ala and Val are conserved in position 2 of the sextet in all three species, there is a considerable difference in the amino acid composition of positions 1 and 6 (fig.2). The rat NF-H favours Glu in position 1 and a nonpolar amino acid (Ala or Val) in position 6. In the mouse NF-H, however, there is a much greater variation in the amino acids present at positions 1 and 6, with no clear pattern emerging. In the human NF-H sequence, while there is a broad range of amino acids in position 6, two main amino acids, Glu and Lys, dominate position 1. As previously mentioned, the triplet repeats in the human NF-H are separated alternatively by 5 and 3 amino acids. One can notice that when one of the extra two amino acids present in a 5 amino acid bridge is a lysine residue and the subsequent amino acid at position 1 of the sextet is usually a Glu residue. Alternatively, when there are only three residues between the repeats, position 1 is always occupied by a Lys residue. From this, one can conclude that the presence of a basic amino acid between the triplet repeats may be important in human NF-H. The reason for the differences seen between species, considering the high degree of homology seen in the rest of the proteins, is unknown. However, given that the region in question is that of phosphorylation, it is possible that the interspecies differences between the amino acid sequences separating the triplet repeats may reflect different binding requirements for the associated protein kinase enzymes, which may exhibit a high degree of species specificity.

Neurofilament phosphorylation has been shown to be related to the neuronal calibre [19] and is very specifically developmentally regulated [2,20]. In the perikarya and developing axons, NF-H is in a relatively unphosphorylated form. In mature axons, however, NF-H is in a highly phosphorylated

form, and it is thought that phosphorylation of the polypeptide results in the formation of interfilament cross bridges that are important in the maintenance of axonal calibre [1,4,5]. Therefore the triplet repeats in the carboxy region are likely to play an important role in the functioning of neurofilaments. The acidic carboxy tailpiece of neurofilaments has been implicated in the formation of interneurofilament cross bridges [4,5] and the proximity of the potential phosphorylation site to this region would have a major influence on protein structure and function.

In conclusion, therefore, while there are some minor differences in the repeat region of the protein in the three species examined, these are unlikely to greatly affect the functioning of this region of the protein. Overall, therefore, the amino acid sequence of the NF-H protein is highly conserved between the mouse, rat and human species.

Acknowledgements: This work was supported by the Medical Research Council. We acknowledge the technical assistance of Ms Suzanne Payne and Mr Stephen Field.

REFERENCES

- [1] Hoffman, P.N., Cleveland, D.W., Griffin, J.W., Landen, P.W., Cowan, N.J. and Price, D.L. (1987) Proc. Natl. Acad. Sci. USA 84, 3472–3476.
- [2] Foster, G.A., Dahl, D. and Lee, V.M.Y. (1987) J. Neurosci. 7, 2651–2663.
- [3] Oblinger, N.M. (1987) J. Neurosci. 7, 2510–2521.
- [4] Geisler, N., Fischer, S., Vanderkerchove, J., Plessman, U. and Weber, K. (1984) EMBO J. 3, 2701–2706.
- [5] Geisler, N., Fischer, S., Vanderkerchove, J., Van Damme, J., Plessmann, U. and Weber, K. (1985) EMBO J. 4, 57–63.
- [6] Julien, J.P., Ramachandran, K. and Grosfeld, F. (1985) Biochim. Biophys. Acta 825, 398–404.
- [7] Lewis, S.A. and Cowan, N.J. (1986) Mol. Cell Biol. 6, 1529–1534.
- [8] Julien, J.P., Grosfeld, F., Yazdanbaksh, K., Flavell, D., Meijer, D. and Mushynski, W. (1987) Biochim. Biophys. Acta 909, 10–20.
- [9] Levy, E., Liem, R.K.H., D'Eustachio, P. and Cowan, N.J. (1987) Eur. J. Biochem. 166, 71–77.
- [10] Myers, M.W., Lazzarini, R.A., Lee, V.M.Y., Schlaepfer, W.W. and Nelson, D.L. (1987) EMBO J. 6, 1617–1626.
- [11] Napolitano, E.W., Chin, S.S.M., Colman, D.R. and Liem, R.K.H. (1987) J. Neurosci. 7, 2590–2599.
- [12] Julien, J.P., Cote, F., Beaudet, L., Sidky, M., Flavell, D., Grosfeld, F. and Mushynski, W. (1988) submitted.
- [13] Lee, J.F., Shneidman, P.S., Skuntz, S.F., Carden, M.J. and Lazzarini, R.A. (1988) EMBO J. 7, 1947–1955.

- [14] Dautigny, A., Pham-Edinh, D., Roussel, C., Felix, J.M., Nussbaum, J.L. and Jolles, P. (1988) *Biochem. Biophys. Res. Commun.* 154, 1099–1106.
- [15] Robinson, P.A., Wion, D. and Anderton, B.H. (1986) *FEBS Lett.* 209, 203–205.
- [16] Sanger, F., Nicklen, S. and Coulson, A.R. (1977) *Proc. Natl. Acad. Sci. USA* 74, 5463–5467.
- [17] Lee, V.M.Y., Otvos, L., Carden, M.J., Hollosi, M., Dietzschold, B. and Lazzarini, R.A. (1988) *Proc. Natl. Acad. Sci. USA* 85, 1998–2002.
- [18] Geisler, N., Vanderkerchove, J. and Weber, K. (1987) *FEBS Lett.* 221, 403–407.
- [19] Carden, M.J., Schlaepfer, W.W. and Lee, V.M.Y. (1985) *J. Biol. Chem.* 260, 9805–9817.
- [20] Carden, M.J., Trojanowski, J.Q., Schlaepfer, W.W. and Lee, V.M.Y. (1987) *J. Neurosci.* 7, 3489–3504.