

Formation of an ion transport supercomplex in *Escherichia coli*

An experimental model of direct transduction of energy

K.A. Bagramyan and S.M. Martirosov

Orbeli Institute of Physiology, Armenian Academy of Sciences and Yerevan State University, Yerevan, USSR

Received 31 January 1989

Hydrogen gas production was observed to occur during ATP-driven H^+/K^+ exchange in anaerobically grown *E. coli*. Neither process was found in aerobically grown cells or anaerobic cells grown on nitrate medium or when the osmotic pressure was decreased or K^+ removed, or finally when DCCD, arsenate or CCCP was applied. Dithiothreitol restored the process even in the presence of CCCP but not in other cases of inhibition. A model of a multienzyme transport supercomplex is proposed. The supercomplex consists of three genetically independent mechanisms: F_0F_1 H^+ -ATPase to provide energy, the K^+ -transporting Trk system as energy sink and formate-hydrogen lyase as donor of reducing equivalents.

Within this supercomplex direct transduction of energy is accomplished via oxidation of 2 SH to S-S.

Exchange, H^+ - K^+ ; Hydrogen evolution; Transport supercomplex; Transmembrane electrochemical proton gradient; Thiol-disulfide interconversion; (*E. coli*)

1. INTRODUCTION

Experimental evidence was obtained recently that in anaerobically grown *E. coli*, the F_0F_1 H^+ -ATPase may unite with a K^+ transporter, the so-called Trk system [1], into a single supercomplex [2,3]. Such a supercomplex functions as an ion-exchanging H^+/K^+ pump with a ratio of $1ATP:2H^+:1K^+$. This phenomenon was not observed in aerobically grown *E. coli* [4] and in mutant strains of *E. coli* with defects in F_0F_1 [5] or in the Trk system [6]. In other words, in anaerobic bacteria the F_0F_1 enzyme transfers part of the energy of ATP hydrolysis directly to the Trk system for creating a K^+ gradient of 2000:1 [4], without participation of the long-range messenger $\Delta\mu H^+$.

Correspondence address: S.M. Martirosov, Orbeli Institute of Physiology, Armenian Academy of Sciences, 375028 Yerevan, USSR

Abbreviations: DCCD, *N,N'*-dicyclohexylcarbodiimide; CCCP, carbonyl cyanide *m*-chlorophenylhydrazone; DTT, dithiothreitol; FHL, formate-hydrogen lyase


The present data allow us to propose a possible mechanism of interaction between F_0F_1 and the Trk system.

2. MATERIALS AND METHODS

Experiments were carried out with the wild-type *E. coli* K-12 (λ). Anaerobic cells were grown in a medium containing peptone with glucose. To obtain anaerobic cells oxidizing formate through nitrate/nitrite respiration [7], 100 mM $NaNO_3$ was introduced into the growth medium. Aerobic cells were grown in a minimal salt medium with succinate. Bacteria were grown for 18–22 h at 37°C. Preparation of bacteria for experiments was detailed in [8]. Ion fluxes were measured using cation-selective glass electrodes as described [8]. The $\Delta\psi$ values were calculated from the distribution of tetraphenylphosphonium cations (TPP^+) with TPP^+ electrodes in cells treated with 10 mM EDTA [9,10].

To measure the oxidation reduction potentials both a platinum electrode (E_1) and an electrode manufactured from titanium-silicate glass (E_h) at the Laboratory of Glass Electrochemistry (Leningrad State University) were used. In contrast with platinum electrodes, those made from electron-conducting glass are insensitive to gaseous O_2 and H_2 and cannot serve as a catalyst for redox reactions. These distinctions between two types of electrode allowed us to use platinum electrodes to record the kinetics of H_2 evolution in anaerobic *E.*

coli during ATP-driven H^+/K^+ exchange. To verify this approach we also measured H_2 production chemically according to the reaction



where a solution with $KMnO_4$ is bleached in the presence of H_2 .

A closed experimental chamber with electrodes was connected through a tube with a test-tube containing the solution. It was established that only anaerobically grown *E. coli* bleached the solution. This effect was detected during the period when the platinum electrode indicated a sharp reduction (broken lines in figures) in contrast to the true E_h observed with the titanium-silicate electrode. Thus, the kinetics of H_2 production recorded with the platinum electrode were found to be correct. The kinetic curves in the figures represent one of 3–5 independent experiments.

3. RESULTS

3.1. H_2 production by formate-hydrogen lyase

It was shown in [4] that the difference between the rates of H_2 extrusion before and after the bend in the H^+ kinetic curves (between 5 and 10 min, fig.1) is the H^+ efflux passing through the F_0F_1 [5] while the intense accumulation of K^+ during the same period occurs via the Trk system [6]. Both counter-fluxes are interrelated with a rigid stoichiometry of $ATP:2H^+1K^+$ [2,3,9].

The important feature of the findings in fig.1 is the coincidence in time of H^+/K^+ pump activity and H_2 evolution. Anaerobic growth of *E. coli* leads to synthesis of membrane-bound formate-hydrogen lyase [7] which splits formate into CO_2 and H_2 . Synthesis of FHL need not be observed in anaerobic *E. coli* grown in nitrate medium or in aerobically grown *E. coli*. Fig.2 indicates that cells without FHL did not produce H_2 (E_h and E_h' curves are the same in fig.2, cf. fig.1) and showed no H^+/K^+ pump activity (see also [4]).

3.2. Absence of H_2 production without H^+/K^+ pump operation

Another approach demonstrates much more clearly the link between FHL and the H^+/K^+ pump. We showed previously, using appropriate mutant strains of *E. coli*, that a decrease in external osmotic pressure or application of DCCD blocks the F_0 proton channel of the F_0F_1 enzyme, whereas the removal of K^+ from the medium obviously eliminates Trk activity [5]. Fig.3 indicates that production of H_2 is also lacking under these conditions. It is important to note that neither H_2




Fig.1. Simultaneous recording of H^+ and K^+ fluxes, membrane potential ($\Delta\psi$) and redox potential (E_h , E_h') measured with a titanium-silicate and a platinum electrode, respectively, in anaerobically grown *E. coli*. The broken line shows the period when the platinum electrode (E_h') registered H_2 production (cf. E_h and E_h' curves); after growth, bacteria were washed with distilled water and resuspended in the experimental medium containing 3 mM KCl, 1 mM NaCl, 0.4 mM $MgSO_4$, Tris-phosphate buffer (pH 7.8) and 50 mM glucose. Bacterial count: about 5×10^9 /ml. During experiments the pH fell from 7.8 to 6.8. However, the decrease in pH had no effect on the data (except for the slow decrease in $\Delta\psi$) because the characteristics were the same when pH was maintained at 7.8. The transfer of bacteria from distilled water into high-osmolarity solution was equivalent to an increase in external osmolarity. This procedure is necessary for the activation of ATP-driven H^+/K^+ exchange [8].

production nor H^+/K^+ exchange was restored by DTT in the experiments illustrated in fig.3.

3.3. $\Delta\mu H^+$ as a regulator of SH groups in the membrane

Despite the noticeable drop in $\Delta\psi$ during glycolysis (fig.1) the absolute value of the membrane potential remained sufficiently high. According to the theory developed by Robillard and Konings [12] such a $\Delta\psi$ value could still support normal transport activity by maintaining the necessary asymmetric distribution of SH and S-S groups inside the membrane.

Elimination of $\Delta\mu H^+$ by CCCP resulted in the suppression of H_2 evolution and H^+/K^+ exchange (fig.4). DTT completely restored the pump and




Fig.2. Data analogous to those shown in fig.1 but in aerobically grown *E. coli*. Similar data were obtained for anaerobic cells grown in the presence of 100 mM NaNO_3 . In this case, 10 mM NaNO_3 was added to the experimental medium.

FHL operation. However, DTT could not restore $\Delta\psi$ -driven K^+ uptake [1,4] in aerobic cells treated with CCCP (not shown).




Fig.3. Data analogous to those in fig.1 but after application of 0.5 mM DCCD. Similar data were obtained under three other conditions: (i) when bacteria were washed with a sucrose solution of high osmolarity and transferred to a medium of low osmotic pressure [8]; (ii) when K^+ was removed from the medium, and (iii) in the presence of 10 mM sodium arsenate.




Fig.4. Inhibition of H^+/K^+ exchange and H_2 production in anaerobic *E. coli* by 10 μM CCCP and recovery of all the characteristics by 5 mM DTT. To reactivate H^+/K^+ exchange by an osmotic upshift [8] 300 mM sucrose was added. Conditions as in fig.1 except that the initial concentration of K^+ was 1 mM.

4. DISCUSSION

Based on our findings [2–6,8,9], including the present data, we can propose a model of a multi-enzyme transport supercomplex (fig.5). The reaction $2\text{SH} \rightarrow \text{S-S} + \text{H}_2$ is accompanied by evolution of H_2 and is characterized by a ΔG of -35 kJ/mol, 20 kJ/mol of which is evidently utilized for counter-gradient accumulation of K^+ (2000:1 [4]). At the same time, the F_0F_1 H^+ -ATPase transfers one H^+ against the $\Delta\mu\text{H}^+$ and the second H^+ only against ΔpH (electroneutral exchange of H^+ vs K^+ at a H^+/K^+ stoichiometry of 2:1, see [3]). Translocation of 2 H^+ requires about 15 kJ/mol in anaerobic *E. coli* at pH 7.8 and a $\Delta\psi$ of 150 mV. Since the phosphate potential in *E. coli* is about 50 kJ/mol [13], the energy required for operation of the supercomplex does not exceed that of ATP hydrolysis.

A difference in the electrochemical potential of H^+ is essential for maintaining sulfhydryl groups in the system [12]. As shown in fig.4, $\Delta\mu\text{H}^+$ can be substituted with DTT. This means that the artificial chemical regulator of SH groups, DTT, is equivalent to the natural electrochemical form, $\Delta\mu\text{H}^+$. The compulsory occurrence of $\Delta\mu\text{H}^+$ or application of DTT for the transfer of reducing




Fig.5. Model of the multienzyme transport supercomplex composed of F_0F_1 H^+ -ATPase as a donor of energy, formate-hydrogen lyase (FHL) as a donor of reducing equivalents and the K^+ transport Trk system, as an acceptor of energy.

equivalents is somewhat extraordinary because the formate/ CO_2 couple has a redox potential of -0.42 V while that of the $2SH/S-S$ couple is only -0.34 V.

Extrapolation of our findings to other transport systems leads to the conclusion that direct transduction of energy through a dithiol-disulfide interchange is impossible without the participation of $\Delta\mu H^+$ (see also [12]). Putting it another way, oxidative phosphorylation or photophosphorylation cannot proceed through direct transfer of energy but only with mediation of $\Delta\mu H^+$.

Acknowledgements: We wish to thank Drs E.A. Karagulin and G.V. Sumarukov for valuable advice and Dr A.G. Vardanian for assistance with some measurements.

REFERENCES

- [1] Rhoads, D.B. and Epstein, W. (1977) *J. Biol. Chem.* 252, 1394–1401.
- [2] Martirosov, S.M. and Trchounian, A.A. (1982) *Bioelectrochem. Bioenerg.* 9, 459–467.
- [3] Bourd, G.I. and Martirosov, S.M. (1983) *Bioelectrochem. Bioenerg.* 10, 315–333.
- [4] Martirosov, S.M. and Trchounian, A.A. (1986) *Bioelectrochem. Bioenerg.* 15, 417–426.
- [5] Martirosov, S.M. and Trchounian, A.A. (1983) *Bioelectrochem. Bioenerg.* 11, 29–36.
- [6] Martirosov, S.M. and Trchounian, A.A. (1981) *Bioelectrochem. Bioenerg.* 8, 597–603.
- [7] Gottschalk, G. (1979) *Bacterial Metabolism*, Springer, Berlin.
- [8] Durgaryan, S.S. and Martirosov, S.M. (1978) *Bioelectrochem. Bioenerg.* 5, 554–560.
- [9] Khachatryan, A.Z., Durgaryan, S.S. and Martirosov, S.M. (1988) *Biochim. Biophys. Acta* 934, 191–200.
- [10] Grinius, L.L. (1986) *Transport of Macromolecules in Bacteria*, Nauka, Moscow (in Russian).
- [11] Midgley, M., Iscander, N.S. and Dawes, E.A. (1986) *Biochim. Biophys. Acta* 856, 45–49.
- [12] Robillard, G.T. and Konings, W.N. (1982) *Eur. J. Biochem.* 127, 597–604.
- [13] Kashket, E.R. (1983) *FEBS Lett.* 154, 343–346.