

# Identity of $G_{A1}$ , $G_{M1a}$ and $G_{D1b}$ synthase in Golgi vesicles from rat liver

Heinrich Iber, Reinhard Kaufmann, Gottfried Pohlentz, Günter Schwarzmann and Konrad Sandhoff

*Institut für Organische Chemie und Biochemie der Universität Bonn, Gerhard-Domagk-Straße 1, D-5300 Bonn 1, FRG*

Received 31 January 1989

Synthesis of ganglioside  $G_{D1b}$  from ganglioside  $G_{D2}$  was demonstrated using Golgi membranes isolated from rat liver. Competition experiments using gangliosides  $G_{A2}$ ,  $G_{M2}$  and  $G_{D2}$  as substrates, and as mutual inhibitors for ganglioside galactosyltransferase activity in preparations of Golgi vesicles derived from rat liver, suggested that galactosyl transfer to these three compounds, leading to gangliosides  $G_{A1}$ ,  $G_{M1a}$  and  $G_{D1b}$  respectively, is catalyzed by one enzyme. These results strengthen the hypothesis that the main site for the regulation of ganglioside biosynthesis occurs within the reaction sequence  $\text{LacCer} \rightarrow G_{A3} \rightarrow G_{D3} \rightarrow G_{T3}$ .

Glycosphingolipid; Ganglioside; Ganglioside synthesis; Galactosyltransferase

## 1. INTRODUCTION

Gangliosides are a group of complex glycosphingolipids characterized by the presence of one or more sialic acid units in their oligosaccharide chain. Although gangliosides are ubiquitous plasma membrane components, little is known about

their function and the regulation of their metabolism. Ganglioside biosynthesis takes place in the Golgi apparatus where, starting with glucosylceramide, it progresses through the sequential addition of galactose, *N*-acetylglactosamine and *N*-acetylneuraminic acid to the growing oligosaccharide chain [1]. These reactions are

*Correspondence address:* K. Sandhoff, Institut für Organische Chemie und Biochemie der Universität Bonn, Gerhard-Domagk-Straße 1, D-5300 Bonn 1, FRG

*Abbreviations:* Cer, ceramide (*N*-acylsphingosine); Gal, galactose; GalNAc, *N*-acetylglactosamine; Glc, glucose; NeuAc, *N*-acetylneuraminic acid; UDP-Gal, uridine 5'-diphosphogalactose; UDP-GalNAc, uridine 5'-diphospho-*N*-acetylglactosamine; CDP-choline, cytidine 5'-diphosphocholine; CMP-NeuAc, cytidine 5'-monophospho-*N*-acetylneuraminic acid; GlcCer, glucosylceramide,  $\text{Glc1} \rightarrow 1\text{Cer}$ ;  $\text{LacCer}$ ,  $\text{Gal}\beta 1 \rightarrow 4\text{Glc}\beta 1 \rightarrow 1\text{Cer}$ . In the nomenclature of Svennerholm [19] for gangliosides: G, ganglioside; M, monosialo; D, disialo; T, trisialo; and arabic numerals indicate the sequence of migration in thin-layer chromatograms

*Enzymes:* UDP-*N*-acetylglactosamine: galactosyl-glucosylceramide  $\beta 1 \rightarrow 4$  *N*-acetylglactosaminyltransferase (EC 2.4.1.-) or  $G_{A2}$  synthase; UDP-*N*-acetylglactosamine: (*N*-acetylneuraminyl)-galactosyl-glucosylceramide  $\beta 1 \rightarrow 4$  *N*-acetylglactosaminyltransferase (EC 2.4.1.92) or  $G_{M2}$  synthase; UDP-*N*-acetylglactosamine: (*N*-acetylneuraminyl)-galactosyl-glucosylceramide  $\beta 1 \rightarrow 4$  *N*-acetylglactosaminyltransferase (EC 2.4.1.-) or  $G_{D2}$  synthase; CMP-*N*-acetylneuramate: (*N*-acetylneuraminyl)-galactosyl-glucosylceramide  $\alpha 2 \rightarrow 8$  sialyltransferase (EC 2.4.99.8) or  $G_{D3}$  synthase; UDP-galactose: *N*-acetylglactosaminyl-galactosyl-glucosylceramide  $\beta 1 \rightarrow 3$  galactosyltransferase (EC 2.4.1.-) or  $G_{A1}$  synthase; UDP-galactose: *N*-acetylglactosaminyl- (*N*-acetylneuraminyl)-galactosyl-glucosylceramide  $\beta 1 \rightarrow 3$  galactosyltransferase (EC 2.4.1.62) or  $G_{M1a}$  synthase; UDP-galactose: *N*-acetylglactosaminyl- (*N*-acetylneuraminyl- *N*-acetylneuraminyl)-galactosyl-glucosylceramide  $\beta 1 \rightarrow 3$  galactosyltransferase (EC 2.4.1.-) or  $G_{D1b}$  synthase; CMP-*N*-acetylneuramate: asialoganglioside ( $G_{A1}$ )  $\alpha 2 \rightarrow 3$  sialyltransferase (EC 2.4.99.-) or  $G_{M1b}$  synthase; CMP-*N*-acetylneuramate: monosialoganglioside ( $G_{M1a}$ )  $\alpha 2 \rightarrow 3$  sialyltransferase (EC 2.4.99.2) or  $G_{D1a}$  synthase; CMP-*N*-acetylneuramate: disialoganglioside ( $G_{D1b}$ )  $\alpha 2 \rightarrow 3$  sialyltransferase (EC 2.4.99.-) or  $G_{T1b}$  synthase

aminyl)-galactosyl-glucosylceramide  $\beta 1 \rightarrow 4$  *N*-acetylglactosaminyltransferase (EC 2.4.1.92) or  $G_{M2}$  synthase; UDP-*N*-acetylglactosamine: (*N*-acetylneuraminyl- *N*-acetylneuraminyl)-galactosyl-glucosylceramide  $\beta 1 \rightarrow 4$  *N*-acetylglactosaminyltransferase (EC 2.4.1.-) or  $G_{D2}$  synthase; CMP-*N*-acetylneuramate: (*N*-acetylneuraminyl)-galactosyl-glucosylceramide  $\alpha 2 \rightarrow 8$  sialyltransferase (EC 2.4.99.8) or  $G_{D3}$  synthase; UDP-galactose: *N*-acetylglactosaminyl-galactosyl-glucosylceramide  $\beta 1 \rightarrow 3$  galactosyltransferase (EC 2.4.1.-) or  $G_{A1}$  synthase; UDP-galactose: *N*-acetylglactosaminyl- (*N*-acetylneuraminyl)-galactosyl-glucosylceramide  $\beta 1 \rightarrow 3$  galactosyltransferase (EC 2.4.1.62) or  $G_{M1a}$  synthase; UDP-galactose: *N*-acetylglactosaminyl- (*N*-acetylneuraminyl- *N*-acetylneuraminyl)-galactosyl-glucosylceramide  $\beta 1 \rightarrow 3$  galactosyltransferase (EC 2.4.1.-) or  $G_{D1b}$  synthase; CMP-*N*-acetylneuramate: asialoganglioside ( $G_{A1}$ )  $\alpha 2 \rightarrow 3$  sialyltransferase (EC 2.4.99.-) or  $G_{M1b}$  synthase; CMP-*N*-acetylneuramate: monosialoganglioside ( $G_{M1a}$ )  $\alpha 2 \rightarrow 3$  sialyltransferase (EC 2.4.99.2) or  $G_{D1a}$  synthase; CMP-*N*-acetylneuramate: disialoganglioside ( $G_{D1b}$ )  $\alpha 2 \rightarrow 3$  sialyltransferase (EC 2.4.99.-) or  $G_{T1b}$  synthase

catalyzed by specific glycosyltransferases. Many of these enzymes, which utilize UDP-Gal, UDP-GalNAc or CMP-NeuAc as sugar donors, have been studied and partially characterized in rat liver Golgi apparatus [2-11]. Some of the glycosyltransferases involved in ganglioside biosynthesis catalyze analogous reactions differing only in the identity of the acceptor (e.g.  $G_{A1}$ ,  $G_{M1a}$ ,  $G_{D1b}$  synthase catalyze the transfer of galactose in  $\beta$ 1-4 linkage to the *N*-acetylgalactosamine residue of the respective acceptor molecule). It has been proposed, but not proven, in earlier studies that these enzymes are identical [12,13]. Pohlentz et al. [10] were the first to demonstrate the identical nature of  $G_{A2}$ ,  $G_{M2}$ ,  $G_{D2}$  synthases as well as  $G_{M1b}$ ,  $G_{D1a}$ ,  $G_{T1b}$  synthases in rat liver Golgi [10]. Here, through kinetic and competition experiments, we show that  $G_{A1}$ ,  $G_{M1a}$  and  $G_{D1b}$  synthases are one and the same enzyme in rat liver Golgi.

## 2. MATERIALS AND METHODS

### 2.1. Materials

UDP-D-[U- $^{14}$ C]galactose were purchased from Amersham (Braunschweig) and used after dilution with the appropriate unlabelled nucleotide sugars obtained from Sigma (Deisenhofen). Sephadex G-25 superfine was obtained from Pharmacia (Freiburg). CDP-choline, Triton X-100 and bovine serum albumin were obtained from Sigma (Deisenhofen). Scintillation cocktail Pico Fluor 30 was purchased from Packard (Frankfurt) and sodium cacodylate from Fluka (Buchs, Switzerland). Male rats of the Wistar strain (300-350 g) were procured from Hagemann (Extertal, FRG). Gangliosides used in this study were available in our laboratory.  $G_{D2}$  was a gift from Professor Svennerholm (Göteborg, Sweden). All other reagents and solvents used were of analytical grade quality.

### 2.2. Methods

Golgi-rich vesicles were isolated from rat liver, essentially by the method of Sandberg et al. [14] as detailed in [15,16]. Enrichment of Golgi-specific enzymes (glycosyltransferases) was 50-80-fold. Contamination with other cellular membranes (plasma membrane, lysosomes, endoplasmic reticulum) was less than 5% [15].

### 2.3. Assay of galactosyltransferases ( $G_{A1}$ , $G_{M1a}$ , $G_{D1b}$ synthase)

In a total volume of 50  $\mu$ l, assay solutions contained glycolipid acceptor ( $G_{A2}$ ,  $G_{M2}$ ,  $G_{D2}$ ) up to 200  $\mu$ M, 0.3% (w/v) Triton X-100, 64 mM sodium cacodylate/HCl (pH 7.35), 10 mM CDP-choline, 20 mM  $MnCl_2$ , 500  $\mu$ M UDP-[ $^{14}$ C]Gal (130-330 GBq/mol) and 50  $\mu$ g Golgi protein. Incubation was for 15 min at 37°C.

Gangliosides  $G_{A2}$ ,  $G_{M2}$  and  $G_{D2}$  and the detergent Triton X-100 were in chloroform/methanol (2:1, v/v) solution. The substances in the organic solvents were first dried under a stream of  $N_2$  and sonicated for 45 s in buffer before use in the

enzyme assay. After incubation for 15 min the reaction was stopped by adding 1 ml chloroform/methanol (2:1, v/v). The gangliosides were separated from radioactive nucleotide sugars by Sephadex G-25 gel chromatography [15]. The radioactivity of the products was determined in a liquid scintillation counter. Rates for all the reactions described in this paper were linear with time, for at least 30 min, and linear with protein concentration up to 50-75  $\mu$ g per assay depending on the Golgi preparation. During the 15 min incubation period the decrease in sugar nucleotide concentration was less than 5%.

All experiments were performed at least twice and mean values are presented in the figures. The  $K_m$  values presented are apparent  $K_m$  values determined in detergent-containing assays.

### 2.4. Product identification

Eluted fractions from the Sephadex columns were dried under a stream of  $N_2$ , redissolved in 100  $\mu$ l chloroform/methanol (2:1, v/v) and applied to silica gel G 60 TLC plates. Chromatograms were developed in chloroform/methanol/0.2%  $CaCl_2$  in water (60:35:8, v/v). Radiolabelled material was visualized by fluorography as described [17].

## 3. RESULTS

The conversion of  $G_{A2}$ ,  $G_{M2}$  and  $G_{D2}$  to  $G_{A1}$ ,  $G_{M1a}$  and  $G_{D1b}$ , respectively, could be catalyzed by either the same or three different galactosyltransferases. In order to discriminate between these two possibilities, two of the three substrates ( $G_{A2}$ ,  $G_{M2}$ ,  $G_{D2}$ ) were used in the enzyme assay at the same time at various relative concentrations, keeping the total acceptor concentration at 200  $\mu$ M. For two independent enzymes, each recognizing only one of the substrates a and b, the total reaction velocity  $v_t$  can be calculated as the sum of two partial velocities  $v_a$  and  $v_b$  given by their respective Michaelis equations (eqn 1):

$$v_t = v_a + v_b = \frac{V_a}{1 + \frac{K_a}{[a]}} + \frac{V_b}{1 + \frac{K_b}{[b]}} \quad (1)$$

If both substrates are accepted by the same enzyme (i.e. the same active site) and each substrate acts as a competitive inhibitor of the other, the inhibitor constant ( $K_i$ ) of either substrate would be equal to its  $K_m$  value ( $K_m = K_i$ ). The total velocity is given by eqn 2 as described in [18]:

$$v_t = v_a + v_b = \frac{V_a}{1 + \frac{K_a}{[a]} \left(1 + \frac{[b]}{K_b}\right)} + \frac{V_b}{1 + \frac{K_b}{[b]} \left(1 + \frac{[a]}{K_a}\right)} \quad (2)$$

The  $K_m$  and  $v_{max}$  values were determined simultaneously with the same Golgi preparation. If




Fig. 1. Competition between  $G_{A2}$  and  $G_{M2}$  in the galactosyltransferase assay. As described in section 2,  $G_{A2}$  and  $G_{M2}$  were used as acceptors for galactosyltransferase in various relative concentrations, keeping the total acceptor concentration at  $20 \mu\text{M}$ . Total reaction velocities determined experimentally (●) or calculated for the different models (eqn 1 or 2) are plotted vs substrate concentration. The kinetic constants used were those given in section 3. Upper curve,  $v_t$  as calculated from eqn 1 (two different enzymes); lower curve,  $v_t$  as calculated from eqn 2 (one enzyme).




Fig. 2. Competition between  $G_{M2}$  and  $G_{D2}$  in the galactosyltransferase assay. As described in section 2,  $G_{M2}$  and  $G_{D2}$  were used as acceptors for galactosyltransferase in various relative concentrations, keeping the total acceptor concentration at  $20 \mu\text{M}$ . Total reaction velocities determined experimentally (●) or calculated for the different models (eqn 1 or 2) are plotted vs substrate concentration. The kinetic constants used were those given in section 3. Upper curve,  $v_t$  as calculated from eqn 1 (two different enzymes); lower curve,  $v_t$  as calculated from eqn 2 (one enzyme).




Fig. 3. Competition between  $G_{A2}$  and  $G_{D2}$  in the galactosyltransferase assay. As described in section 2,  $G_{A2}$  and  $G_{D2}$  were used as acceptors for galactosyltransferase in various relative concentrations, keeping the total acceptor concentration at  $20 \mu\text{M}$ . Total reaction velocities determined experimentally (●) or calculated for the different models (eqn 1 or 2) are plotted vs substrate concentration. The kinetic constants used were those given in section 3. Upper curve,  $v_t$  as calculated from eqn 1 (two different enzymes); lower curve,  $v_t$  as calculated from eqn 2 (one enzyme).

$G_{A2}$  and  $G_{M2}$  were used in the galactosyltransferase assay as acceptors ( $G_{A2}$ ,  $K_m = 8 \mu\text{M}$ ,  $v_{\max} = 15 \text{ nmol/h per mg}$ ;  $G_{M2}$ ,  $K_m = 4 \mu\text{M}$ ,  $v_{\max} = 15 \text{ nmol/h per mg}$ ) the total reaction velocities could be calculated from eqns 1 and 2 (fig.1). The measured  $v_t$  values clearly fitted with those calculated using eqn 2; these results show that  $G_{A1}$  and  $G_{M1a}$  are synthesized from their respective precursors at the same active site of a single enzyme. In analogous experiments when  $G_{M2}$  and  $G_{D2}$  ( $G_{M2}$ ,  $K_m = 9 \mu\text{M}$ ;  $v_{\max} = 31 \text{ nmol/h per mg}$ ;  $G_{D2}$ ,  $K_m = 6 \mu\text{M}$ ;  $v_{\max} = 30 \text{ nmol/h per mg}$ ), or  $G_{A2}$  and  $G_{D2}$  ( $G_{A2}$ ,  $K_m = 27 \mu\text{M}$ ;  $v_{\max} = 40 \text{ nmol/h per mg}$ ;  $G_{D2}$ ,  $K_m = 6 \mu\text{M}$ ;  $v_{\max} = 49 \text{ nmol/h per mg}$ ) were used as glycolipid acceptors for synthesis of  $G_{M1a}$  and  $G_{D1b}$  or of  $G_{A1}$  and  $G_{D1b}$ , respectively, similar results were obtained (figs 2,3).

These experiments show that  $G_{A1}$ ,  $G_{M1a}$  and  $G_{D1b}$  synthesis is catalyzed by a single galactosyltransferase in rat liver Golgi.

#### 4. DISCUSSION

The identical nature of the  $G_{M2}$  and  $G_{D2}$  syn-




Fig.4. Suggested scheme for ganglioside biosynthesis.

thases,  $G_{M1a}$  and  $G_{D1b}$  synthases, and  $G_{D1a}$  and  $G_{T1b}$  synthases was suspected in earlier studies [12,13]. Pohlentz et al. [10] were able to demonstrate that the synthesis of  $G_{A2}$ ,  $G_{M2}$  and  $G_{D2}$  from their respective precursors is catalyzed by the same *N*-acetylgalactosaminyltransferase in rat liver Golgi, as is the synthesis of  $G_{M1b}$ ,  $G_{D1a}$  and  $G_{T1b}$  by the same sialyltransferase [10]. By proving the non-identity of  $G_{D3}$  and  $G_{M1a}$  synthase they also demonstrated that this method allows one to distinguish between two different enzymes [10].

The present study demonstrates that  $G_{A1}$ ,  $G_{M1a}$  and  $G_{D1b}$  are synthesized by the same galactosyl-transferase in rat liver Golgi. The synthesis of  $G_{D1b}$  from  $G_{D2}$ , using Golgi membranes isolated from rat liver, is demonstrated for the first time. The results support the modified model of ganglioside biosynthesis proposed previously (fig.4) [10]. In this model, the sialyltransferases I-III seem to determine to which series a certain ganglioside molecule is directed (asialo, a, b or c). Starting with LacCer,  $G_{M3}$  or  $G_{D3}$ , and also possibly with  $G_{T3}$ , further biosynthesis of gangliosides belonging to the three, and possibly four, different series (i.e. asialo, a, b or c) is catalyzed by the same set of enzymes, namely *N*-acetylgalactosaminyltransferase, galactosyltransferase II, sialyltransferase IV and

sialyltransferase V (fig.4). These four enzymes seem to recognize the common neutral carbohydrate 'backbones' of their respective acceptors. The different number of sialic acid residues bound to the inner galactose of the carbohydrate backbone has an influence on the kinetic properties of these glycosyltransferases. The present results also support our hypothesis [10] that a major site for the regulation of ganglioside biosynthesis occurs in the reaction sequence LacCer  $\rightarrow$   $G_{M3}$   $\rightarrow$   $G_{D3}$   $\rightarrow$   $G_{T3}$  (fig.4).

**Acknowledgements:** We are very grateful for the gift of  $G_{D2}$  from Professor Svennerholm (Göteborg). This work was supported financially by the Deutsche Forschungsgemeinschaft and the Fonds der Chemischen Industrie.

## REFERENCES

- [1] Schachter, H. and Roseman, S. (1980) in: *The Biochemistry of Glycoproteins and Proteoglycans* (Lennarz, W.J. ed.) pp. 85-160, Plenum, New York.
- [2] Wilkinson, F.E., Morre, D.J. and Keenan, T.W. (1976) *J. Lipid Res.* 17, 146-153.
- [3] Richardson, C.-L., Keenan, T.W. and Morre, D.J. (1977) *Biochim. Biophys. Acta* 488, 88-96.
- [4] Eppler, C.M., Morre, D.J. and Keenan, T.W. (1980) *Biochim. Biophys. Acta* 619, 318-331.

- [5] Senn, H.J., Cooper, C., Warnke, P.C., Wagner, M. and Decker, K. (1981) *Eur. J. Biochem.* 120, 59-67.
- [6] Senn, H.J., Wagner, M. and Decker, K. (1983) *Eur. J. Biochem.* 135, 231-236.
- [7] Kaplan, F. and Hechtman, P. (1983) *J. Biol. Chem.* 258, 770-776.
- [8] Busam, K. and Decker, K. (1986) *Eur. J. Biochem.* 160, 23-30.
- [9] Klein, D., Pohlentz, G., Schwarzmnn, G. and Sandhoff, K. (1987) *Eur. J. Biochem.* 167, 417-427.
- [10] Pohlentz, G., Klein, D., Schwarzmnn, G., Schmitz, D. and Sandhoff, K. (1988) *Proc. Natl. Acad. Sci. USA* 85, 7044-7048.
- [11] Pohlentz, G., Klein, D., Schwarzmnn, G. and Sandhoff, K. (1988) *Biol. Chem. Hoppe-Seyler* 369, 55-63.
- [12] Cumar, F.A., Tallmann, J.F. and Brady, R.O. (1972) *J. Biol. Chem.* 247, 2322-2327.
- [13] Pacuszka, T., Duffard, R.O., Nishimura, R.N., Brady, R.O. and Fishman, P.H. (1978) *J. Biol. Chem.* 253, 5839-5846.
- [14] Sandberg, P.O., Marzella, L. and Glaumann, H. (1980) *Exp. Cell Res.* 130, 393-400.
- [15] Yusuf, H.K.M., Pohlentz, G., Schwarzmnn, G. and Sandhoff, K. (1983) *Eur. J. Biochem.* 134, 47-54.
- [16] Yusuf, H.K.M., Pohlentz, G. and Sandhoff, K. (1983) *Proc. Natl. Acad. Sci. USA* 80, 7075-7079.
- [17] Sonderfeld, S., Conzelmann, E., Schwarzmnn, G., Burg, J., Hinrichs, U. and Sandhoff, K. (1985) *Eur. J. Biochem.* 149, 247-255.
- [18] Dixon, M. and Webb, E.C. (1979) *Enzymes*, p. 334, Longman, London.
- [19] Svennerholm, L. (1963) *J. Neurochem.* 10, 613-623.