

## Expression in *E. coli* of the catalytic domain of rat poly(ADP-ribose)polymerase

Jacques Thibodeau<sup>1</sup>, Frédéric Simonin<sup>2</sup>, Marisa Favazza<sup>2</sup>, Gérard Gradwohl<sup>2</sup>, Guy Poirier<sup>1</sup> and Gilbert de Murcia<sup>2</sup>

<sup>1</sup>Centre Hospitalier de l'Université Laval, Laboratoire du métabolisme du poly(ADP-ribose), Endocrinologie Moléculaire, 2705 boul. Laurier, Québec, G1V 4G2, Canada and <sup>2</sup>Institut de Biologie Moléculaire et Cellulaire du CNRS and Centre de Recherche de l'Université Louis Pasteur, 15 rue Descartes, 67084 Strasbourg, France

Received 7 March 1990

A 2 kilobase pair cDNA coding for the entire C-terminal catalytic domain of rat poly(ADP-ribose)polymerase has been expressed in *E. coli*. The overproduced 55 kDa polypeptide is active in synthesizing poly(ADP-ribose) and the 4 kDa N-terminal region of this domain is recognized by the monoclonal antibody C I,2 directed against the calf enzyme. Also, the minor  $\alpha$ -chymotrypsin cleavage site found in the human catalytic domain is not present in the rat enzyme as revealed by the absence of the 40 kDa specific degradation product in the *E. coli* cells expressing the rat domain. The expression of this partial rat cDNA should thus permit the rapid purification and subsequent crystallization of the catalytic domain of the enzyme.

Poly(ADP-ribose)polymerase; *E. coli* expression;  $\alpha$ -Chymotrypsin; NAD<sup>+</sup> metabolism; Immunoblotting

### 1. INTRODUCTION

Poly(ADP-ribose)polymerase (PARP) is a nuclear enzyme that catalyzes the covalent attachment and elongation of poly(ADP-ribose) from NAD<sup>+</sup> on nuclear proteins such as histones, topoisomerase I and PARP (automodification) (for review, see [1]). Poly(ADP-ribose) is implicated in nuclear events including the relaxation of chromatin, DNA repair and gene expression. PARP can be divided into three functional domains when treated with papain and  $\alpha$ -chymotrypsin [2–4]. The N-terminal 42 kDa domain contains 2 zinc-finger motifs that bind specifically to DNA single strand breaks [5,6]. The internal 16 kDa is the automodification domain while the C-terminal 55 kDa domain binds NAD<sup>+</sup> and is the catalytic region.

We have recently cloned a partial rat cDNA coding for the entire catalytic domain of the enzyme [7]. The predicted amino acid sequence is greater than 95% homologous to the same domain of the human [3,8–10] and mouse [11] proteins. Most importantly this domain was shown to be the target of autoantibodies against the PARP in several human rheumatic patients [12]. As

a first step toward the purification and crystallization of the catalytic domain, we have expressed in *E. coli* the rat cDNA coding for this region of the PARP.

### 2. MATERIALS AND METHODS

#### 2.1. Expression in *E. coli*

We have used a rat cDNA which is one bp (cytosine) longer at the 5' end than the one described previously [7]. It was inserted in the *EcoRI* site of pTG161 [13] to give the recombinant plasmid, pTGrat55. We have also cloned a *HindIII* fragment of the human PARP cDNA [10] in the plasmid pTG920 [13]. This 2.66 kbp fragment (pTGhum87) starts at the nucleotide position +698 [3] and codes for the catalytic domain, the automodification domain and part of the DNA binding domain. The plasmid pTG161 SH1.4 codes for the C-terminal 51 kDa part of the human catalytic domain, starting from the amino acid 572 corresponding to the *PstI* site [3]. *E. coli* TGE 900 was transformed with the recombinant plasmids, grown at 28°C and shifted to 37°C to induce the transcription of the foreign PARP sequences [13]. Preparation, SDS-PAGE and electrophoretic transfer of the crude extracts on the solid support were as described [14].

#### 2.2. Detection methods

The PARP-specific polypeptides were revealed on the nitrocellulose sheet by 2 methods as described [14]. First, an activity blot using [<sup>32</sup>P]NAD<sup>+</sup> was done in the presence of calf thymus DNA. The immunogenic fragments were then revealed using monoclonal and polyclonal antibodies against the PARP [4] and second antibodies conjugated to alkaline phosphatase [6].

### 3. RESULTS AND DISCUSSION

We have compared the expression in *E. coli* of the

Correspondence address: G. de Murcia, Institut de Biologie Moléculaire et Cellulaire du CNRS, Laboratoire de Biochimie 2, 15 rue Descartes, 67084 Strasbourg Cedex, France

Abbreviations: PARP, poly(ADP-ribose)polymerase; kbp, kilobase pair(s); PAGE, polyacrylamide gel electrophoresis



the overexpressed rat 55 kDa fragment is active in synthesizing poly(ADP-ribose). Taken together these data show that the rat cDNA is the most suitable choice for the overexpression, purification and crystallization attempts of the PARP catalytic domain.

Finally, the difference in size between the rat 55 kDa fragment (which is recognized by the monoclonal antibody C I,2) and the human 51 kDa fragment (which is not (Fig. 4)), represents the epitope of this antibody. This region is part of the 15 kDa  $\alpha$ -chymotrypsin fragment recognized by human autoantibodies [12]. Unlike these antibodies C I,2 does not inhibit the activity of the PARP [4] suggesting that their epitopes are different. However, both map outside the 40 kDa fragment which is still active (Fig. 2B, lane 3) and contains the highly conserved putative nucleotide binding site [7].

**Acknowledgements:** We thank Drs J. Ménissier-de Murcia and James Kirkland for a critical review of the manuscript. The secretarial assistance of J. Poulin and E. Leclerc is greatly acknowledged. J.T. is supported by the Cancer Research Society Inc. and F.F. by the Association pour la Recherche contre le Cancer. M.F. was supported by a grant from EMBO. This work was carried out in Professor G. Dirheimer's laboratory and was supported by the Ligue Nationale contre le Cancer, Comité Départemental du Haut-Rhin, and ARC (Association pour la Recherche contre le Cancer). The human cDNA (from nucleotide 698 to 1840) was kindly provided by Dr M. Schweiger. We thank Transgène (Strasbourg) for the gift of pTG161, pTG920 and *E. coli* TGE900.

## REFERENCES

- [1] Althaus, F.R. and Richter C.R. (1987) *Mol. Biol. Biochem. Biophys.* 37, 1–125.
- [2] Kameshita, I., Matsuda, Z., Taniguchi, T. and Shizuta, Y. (1984) *J. Biol. Chem.* 259, 4770–4776.
- [3] Kurosaki, T., Ushiro, H., Mitsuuchi, Y., Suzuki, S., Matsuda, M., Matsuda, Y., Katunuma, N., Kangawa, K., Matsuo, H., Hirose, T., Inayama, S. and Shizuta, Y. (1987) *J. Biol. Chem.* 262, 15990–15997.
- [4] Lamarre, D., Talbot, B., De Murcia, G., Laplante, C., Leduc, Y., Mazen, A. and Poirier, G.G. (1988) *Biochem. Biophys. Acta* 950, 147–160.
- [5] Ménissier-De Murcia, J., Molinete, M., Gradwohl, G., Simonin, F. and De Murcia, G. (1989) *J. Mol. Biol.* 210, 229–234.
- [6] Mazen, A., Ménissier-De Murcia, J., Molinete, M., Simonin, F., Gradwohl, G., Poirier, G. and De Murcia, G. (1989) *Nucleic Acids Res.* 17, 4689–4698.
- [7] Thibodau, J., Gradwohl, G., Dumas, C., Clairoux-Moreau, S., Brunet, G., Penning, C., Poirier, G.G. and Moreau, P. (1989) *Biochem. Cell Biol.* 67, 653–660.
- [8] Uchida, K., Morita, T., Sato, T., Ogura, T., Yamashita, R., Noguchi, S., Suzuki, H., Nyunoya, H., Miwa, M. and Sugimura, T. (1987) *Biochem. Biophys. Res. Commun.* 148, 617–622.
- [9] Cherney, B.W., McBride, O.W., Chen, O., Alkhatib, H., Bhatia, K., Hensley, P. and Smulson, M.E. (1987) *Proc. Natl. Acad. Sci. USA* 84, 8370–8374.
- [10] Herzog, H., Zadel, B.U., Schneider, R., Auer, B., Hirsch-Kauffman, M. and Schweiger, M. (1989) *Proc. Natl. Acad. Sci. USA* 86, 3514–3518.
- [11] Huppi, K., Bhatia, K., Siwarski, D., Klinman, D., Cherney, B. and Smulson, M. (1989) *Nucleic Acids Res.* 17, 3387–3401.
- [12] Yamanaka, H., Willis, E.H. and Carson, D.A. (1989) *J. Clin. Invest.* 83, 180–186.
- [13] Courtney, M., Buchwalder, A., Tessier, L.H., Jaye, M., Benavente, A., Balland, A., Kohli, V., Lathe, R., Tolstoshev, P. and Lecocq, J.P. (1984) *Proc. Natl. Acad. Sci. USA* 81, 669–673.
- [14] Simonin, F., Ménissier-De Murcia, J., Poch, O., Muller, S., Gradwohl, G., Molinete, M., Penning, G., Keith, G. and De Murcia, G., *J. Biol. Chem.* in press.
- [15] Kameshita, I., Yamamoto, H., Fugimoto, S. and Shizuta, Y. (1985) *FEBS Lett.* 182, 393–397.
- [16] Ikejima, M., Noguchi, S., Yamashita, R., Suzuki, H., Sugimura, T. and Miwa, M. (1989) *Biochem. Biophys. Res. Commun.* 163, 739–745.
- [17] Ushiro, H., Yokayama, Y. and Shizuta, T. (1987) *J. Biol. Chem.* 262, 2352–2357.
- [18] Taniguchi, T., Yamauchi, K., Yamamoto, T., Toyoshima, K., Harada, N., Tanaka, H., Takahashi, S., Yamamoto, H. and Fujimoto, S. (1988) *Eur. J. Biochem.* 171, 571–575.