

Rate-determining steps in penicillopepsin-catalysed reactions

Annie Cunningham¹, Martin I. Hofmann² and Theo Hofmann¹

Departments of ¹Biochemistry and ²Zoology, University of Toronto, Toronto, M5S 1A8 Canada

Received 26 September 1990

The hydrolysis of Ac-(Ala)₂-Lys-Nph-(Ala)₂-amide (II) by penicillopepsin is characterized by a solvent isotope effect of 2.11, whereas the hydrolysis of Ac-Lys-Nph-amide (I) shows no solvent isotope effect. The dependence of the isotope effect on the concentration of D₂O in H₂O for substrate II is not linear and suggests that two or more protons are involved in its rate-determining step. We propose that for substrate I the rate-determining step is the distortion of the scissile bond towards a tetrahedral configuration, and for substrate II a conformational change induced by the occupation of the S₃ pocket in the enzyme.

Penicillopepsin; Aspartic proteinase; Enzyme kinetics; Enzyme mechanism

1. INTRODUCTION

Although several high resolution structures of aspartic proteinases and their complexes with substrate analogue inhibitors are known (see Gilliland et al. for a recent list [2]) and several proposals for their mechanism of action have been made [3-7], many aspects, including the nature of the rate-determining step, are still poorly understood. Information on rate-determining steps in hydrolytic reactions can be obtained from a study of solvent isotope effects. Three such studies were carried out earlier with pig pepsin, but they led to conflicting conclusions. No solvent isotope effects were observed with the substrates Ac-Phe-Tyr-OMe [8] and methyl phenyl sulfite [9], whereas Holland and Fruton [10] observed a solvent isotope effect of about 2 with Gly₃-Nph-Phe-OMe. The latter authors could not find a satisfactory explanation for this discrepancy and felt that further work was needed to resolve this problem. However, to the best of our knowledge no such additional study with any of the aspartic proteinases has been carried out. In the present study we looked at the solvent isotope effect on the hydrolysis by penicillopepsin of the -Lys-Nph- bond of two substrates of the series Ac-Ala_m-Lys-Nph-Ala_n-amide, one with $m = n = 0$, the other with

$m = n = 2$. This substrate series was used previously to determine the effect of different subsites on k_{cat} of the scissile bond [11]. We showed that subsites S₃ and S₂' were primarily responsible for the large increases in k_{cat} with increasing chain length. In a study of the effect of temperature on the reactions [2] we further showed that hydrolysis of Ac-Lys-Nph-amide was not accompanied by any demonstrable conformational changes, whereas hydrolysis of Ac-Ala₂-Lys-Nph-Ala₂-amide was. Both of these earlier studies suggested the possibility that rate-determining steps could be different for the short and the long substrates.

2. MATERIALS AND METHODS

2.1. Materials

Penicillopepsin was prepared as described [12]. The synthesis of Ac-Lys-Nph-amide (I) and Ca-Ala₂-Lys-Nph-Ala₂-amide (II) was described by Hofmann and Hodges [13]. Chemicals used for buffers were of the highest quality available. [¹⁸O]water (97 atom% ¹⁸O) and D₂O (99.9 atom% D) were obtained from MSD Isotopes (Merck Frosst Canada Inc., Montreal).

2.2. Enzyme kinetics

All enzyme assays were carried out at 25°C in 20 mM formate (pH 3.5, pD 3.9), or acetate (pH 4-5.5, pD 4.4-5.9), adjusted to an ionic strength of 20 mM with NaCl, in a Uvikon 820 spectrophotometer (Kontron AG, Zurich, Switzerland) at a wavelength of 335 nm. (This wavelength, rather than the optimal wavelength of 296 nm [13], was chosen because of the high absorbance at the substrate concentrations used.) The buffers contained between 0% and 93% D₂O. The enzyme concentrations were 6.3 μM for peptide I and 0.052 μM for peptide II.

2.3. Analysis of ¹⁸O incorporation

The reaction conditions for the analysis of the incorporation of ¹⁸O into the amino moieties of the substrates were the same as those described above, except that the buffer contained 92.4% [¹⁸O]H₂O. The pH was 5.5, the substrate concentrations were about 2 mM, and the enzyme concentrations were as given above. The reactions were allowed to reach over 90% hydrolysis and were stopped by the addition of 50 μl of 1 N ammonia. The products Ac-Lys and Ac-Ala₂-Lys from peptides I and II respectively, were isolated by HPLC on a C₁₈

Correspondence address: T. Hofmann, Dept. of Biochemistry, King's College Circle, University of Toronto, Toronto, Canada M5S 1A8

Abbreviations: Ac-, acetyl; FAB, fast atom bombardment; HPLC, high-pressure liquid chromatography; Nph, *p*-nitrophenylalanyl; OMe, *O*-methyl ester; TFA, trifluoroacetic acid; the definition of Schechter and Berger [1] for denoting amino acid residues in peptide substrates as P₁ to P_n and P₁' to P_n' and subsites in the enzyme to which the side chains of these residues bind as S₁ to S_n and S₁' to S_n', respectively, is used throughout

Table I
Solvent isotope effect for penicillopepsin catalysed hydrolyses.

pH (pD)	Rate constant k_{obs} (s ⁻¹)		K_m (mM)		k^H/k^D
	H ₂ O	D ₂ O	H ₂ O	D ₂ O	
I = Ac-Lys-Nph-amide					
5.5 (5.9)	0.013 ^a	0.012 ^a	0.2	0.25	1.08
5.0–5.5 (5.4–5.9)	0.0123 ± 0.007 ^b	0.0113 ± 0.006 ^b			1.09 ± 0.04
II = Ac-Ala₂-Lys-Nph-Ala₂-amide					
4.5 (4.9)	46.4 ^a	23.6 ^a	0.08	0.06	1.97
3.5 (3.9)	34.5 ^a	19.7 ^a			1.75
5.5 (5.9)	35.6 ^c	15 ^c			2.33
4.5 (4.9)	40 ^c	18 ^c			2.22
3.5 (3.9)	28.2 ^c	12.9 ^c			2.18
5.25 (5.65)					2.11 ± 0.16 ^d

The conditions were as described in section 2.

^a k_{cat}

^bSubstrate concentration = 10 × K_m , average of 6 experiments

^cSubstrate concentration = 23 × K_m , average of 2–3 experiments

^dfrom the experiments shown in Fig. 1

reverse-phase Vydac 218TP104 column (250 × 4.6 mm) in 0.1% TFA, run at room temperature with a linear gradient from 0% to 50% acetonitrile over 25 min. The peptides were analysed by FAB mass spectrometry on a VG Analytical ZAB SE mass spectrometer (V.G. Analytical, Manchester, UK). The bombarding xenon atom (8 kV, 1.2 mA anode current) was generated with the 'Ion Tech' gun supplied with the instrument; glycerol or *m*-nitrobenzyl alcohol was used as the matrix. Spectra were recorded with a VG 11-250 data system in a multichannel analysis mode in a downfield exponential magnetic scan.

2.4. Analysis of dependence of the rate constant on the D₂O/H₂O ratio

The shape of the plot of the rate constants k_{obs} versus the ratio (n) of D₂O/H₂O was analysed in an attempt to determine the number of protons involved in the rate-determining step [14]. Three different hypotheses, corresponding to the following equations, were tested.

one-proton mechanism: $k_n = k_0 - a n$ (1)

two-proton mechanism: $(k_n/k_0)^{0.5} = b - a n$ (2)

'infinite' proton mechanism: $\ln(k_n/k_0) = -a n + \ln k_0$ (3)

Statistical tests were carried out to determine how well the data fit these 3 linear equations. For each run the data were normalized as follows. First, the mean value for each run was calculated, then the value of each point was divided by the mean value for that run. These normalized data were then combined to produce a single data set that was used in the statistics. The linearity of the curves was tested with the REG procedure from the SAS package [15].

3. RESULTS AND DISCUSSION

Table I shows the kinetic parameters for the hydrolysis of the -Lys-Nph- bond in the two substrates. In the range of pH 5–5.5 (pD 5.4–5.9) the rate constants for peptide I, measured at a substrate concentration about 10 × K_m , are independent of pH and show no significant solvent isotope effect. The K_m s, determined at pH 5.5 (pD 5.9) in separate experiments, were 0.2 mM in H₂O and 0.25 mM in D₂O and not considered significantly different. They agree well with the

previous value in H₂O of 0.22 ± 0.01 mM [16]. The absence of a solvent isotope effect shows that the rate-determining step does not involve the bond breaking step. According to the mechanistic proposals at present generally accepted, the bond breaking is catalysed by a general base mechanism leading to a tetrahedral intermediate, the decay of which involves a proton transfer to the leaving carboxyl moiety [3,4,17]. Both steps involve a proton transfer and hence are likely to show a solvent isotope effect.

If we assume the following scheme as a minimal scheme for the action of penicillopepsin [11].



where the steps E*S to E*P represent the bond breaking steps via the tetrahedral intermediate E*T, the rate-determining step for peptide I lies either before or after these steps. We can eliminate the release of products as the rate-determining step on the following grounds. We show elsewhere (Blum, M., Cunningham, A., Pang, H. and Hofmann, T. submitted for publication) that during transpeptidation reactions release of some products is the rate-determining step. The evidence for this comes from a high degree of incorporation of ¹⁸O into peptide carbonyl and α-carboxyl groups of the products when the reactions are carried out in [¹⁸O]H₂O buffers. Since the rate constants for these reactions are similar to those observed with peptide I product release could also be the rate-determining step for this peptide. We therefore isolated Ac-Lys from a reaction carried out in [¹⁸O]H₂O and found that only one atom of ¹⁸O had been incorporated into the terminal carboxyl group. This shows that the peptide bond cleavage is irreversible, unlike that observed in the transpeptidation reactions and suggests that product release is not the rate-determining step.




Fig. 1. Solvent isotope effects for penicillopepsin catalysed hydrolysis of Ac-(Ala)₂-Lys-Nph-(Ala)₂-amide as a function of D₂O concentration. The buffer used was 20 mM sodium acetate (pH 5.25; pD 5.65); penicillopepsin concentration was 32.5 nM, the substrate concentration was 1.17 mM.

In contrast, the rate constants for peptide **II** show a consistent solvent isotope effect of 2.11 ± 0.16 over the pH range 3.5–5.5 (pD 3.9–5.9) (Table I). In this case also the K_m values are unaffected. The results of a proton inventory, undertaken in order to find out the number of protons actively involved in the rate-determining step [18], are shown in Fig. 1 for the logarithmic relation (Eqn 3), the equation for which the best fit was obtained. We should stress, however, that the fit and the statistical analysis for the relation shown in Eqn 2 is such that it cannot be unambiguously excluded. This suggests that a process involving two or more protons controls the rate-determining step for peptide **II**. Although the interpretation of this finding in molecular terms is not straightforward, the rate-determining step can be placed later than that for substrate **I** and before the product releasing step. The experiments carried out in [¹⁸O]H₂O showed that, like substrate **I**, substrate **II** did not incorporate excess ¹⁸O. It is most likely that the rate-determining step is a conformational change that occurs only when a substrate or other ligand contains an amino acid residue in P₃. We showed previously that conformational changes are associated with the binding of ligands that contain amino acid residues in P₃ and in P₂' [19]. Furthermore, structural studies with inhibitor complexes of endothiapepsin indicate that occupation of the S₃ pocket induces a rotation of a domain comprising residues 190–302 (pig pepsin numbering) which has been described as a 'rigid body movement' [20,21]. The same type of change accounts for half the conformational differences among the aspartic proteinases [21]. On the basis of this the proposal has been made that this type of structural flexibility plays a major part in the function of these enzymes [21]. We also showed that the oc-

cupation of subsite S₃ involves the formation of a specific hydrogen bond between the NH of P₃ and the O_γ of Thr-219 (pepsin numbering) [22] and pointed out that an amino acid with a hydrogen bond accepting O atom in the side chain (Thr, Ser or Asn) is present in the analogous position in all 24 known sequences of aspartic proteinases [11]. There is also evidence that similar unique hydrogen bonds form when the S₂' subsite is occupied [11]. Hence we propose that it is the formation of these hydrogen bonds and possibly other protonic reorganisations involved in concurrent conformational changes [23] that are responsible for the solvent isotope effect and represent the rate-determining step in the hydrolysis of substrate **II**.

In retrospect the results obtained here also provide an explanation of the earlier experiments [8–10]. As in this study the short substrates, dipeptide and sulphite ester, showed no solvent isotope effect [8,9], whereas the longer substrate which had a glycine residue in P₃ showed an effect which was the same as that observed with peptide **II**.

Acknowledgements: We are grateful to Drs M. Blum, R. Kluger and A.J. Kresge for valuable discussions and to Dr Henrianna Pang for the ¹⁸O analyses. This work was supported by the Medical Research Council of Canada (Grant MT-1982).

REFERENCES

- [1] Schechter, I. and Berger, A. (1967) *Biochem. Biophys. Res. Commun.* 27, 157–162.
- [2] Gilliland, G.L., Winborne, E.L., Nachman, J. and Wlodawer, A. (1990) *Proteins Struct. Funct. Genet.* 8, 82–101.
- [3] Pearl, L.H. (1987) *FEBS Lett.* 214, 8–12.
- [4] Polgar, L. (1987) *FEBS Lett.* 219, 1–4.
- [5] Fischer, G. (1987) in: *Enzyme Mechanisms* (Page, M.I. and Williams, A. eds) pp. 229–239, Roy. Soc. Chem., London.
- [6] James, M.N.G. and Sielecki, A. (1984) in: *Biological Macromolecules and Assemblies, Active Sites of Enzymes*, vol. 3 (Jurnak, F.A. and McPherson, A. eds) pp. 415–482, Wiley, New York.
- [7] Hofmann, T., Dunn, B.M. and Fink, A.L. (1984) *Biochemistry* 23, 5253–5256.
- [8] Clement, C.E. and Snyder, S.L. (1966) *J. Am. Chem. Soc.* 88, 5338–5339.
- [9] Reid, T.W. and Fahrney, D. (1967) *J. Am. Chem. Soc.* 89, 3941–3943.
- [10] Hollands, T.R. and Fruton, J.S. (1969) *Proc. Natl. Acad. Sci. USA* 62, 1116–1120.
- [11] Hofmann, T., Allen, B., Bendiner, M., Blum, M. and Cunningham, A. (1988) *Biochemistry* 27, 1140–1146.
- [12] Hofmann, T. (1976) *Methods Enzymol.* 45, 434–452.
- [13] Hofmann, T. and Hodges, R.S. (1982) *Biochem. J.* 203, 603–610.
- [14] Schowen, K.B.J. (1978) in: *Transition States of Biochemical Processes* (Gandour, R.D. and Schowen, R.L. eds) pp. 225–283.
- [15] SAS Inst. (1985) *SAS User's Guide: Statistics*, version 5 Ed., SAS Inst. Inc., Cary, NC, USA.
- [16] Hofmann, T., Hodges, R.S. and James, M.N.G. (1984) *Biochemistry* 23, 635–643.
- [17] James, M.N.G. and Sielecki, A. (1985) *Biochemistry* 24, 3701–3713.

- [18] Schowen, R.L. (1977) in: *Isotope Effects on Enzyme-catalyzed Reactions* (Cleland, W.W., O'Leary, M.H. and Northrop, D.B. eds) pp. 64-99.
- [19] Allen, B., Blum, M., Cunningham, A., Tu, G.C. and Hofmann, T. (1990) *J. Biol. Chem.* 265, 5060-5065.
- [20] Sali, A., Veerapandian, B., Cooper, J.B., Foundling, S.I., Hoover, D.J. and Blundell, T.L. (1989) *EMBO J.* 8, 2179-2188.
- [21] Sali, A., Veerapandian, B., Cooper, J.B., Moss, D.S., Hofmann, T. and Blundell, T.L. (1990) *Proteins Struct. Funct. Genet.* (in press).
- [22] James, M.N.G., Sielecki, A., Salituro, F., Rich, D.H. and Hofmann, T. (1982) *Proc. Natl. Acad. Sci. USA* 79, 6137-6141.
- [23] Venkatasubban, V.S. and Schowen, R.L. (1984) *CRC Crit. Rev. Biochem.* 17, 1-44.