

Molecular cloning and expression of a protein-tyrosine phosphatase showing homology with transcription factors Fos and Jun

G. Swarup, S. Kamatkar, V. Radha and V. Rema

Centre for Cellular and Molecular Biology, Uppal Road, Hyderabad 500 007, India

Received 21 November 1990; revised version received 8 January 1991

A cDNA clone coding for a protein-tyrosine phosphatase (PTPase) was isolated from a rat spleen cDNA library. Nucleotide sequence of the clone showed an open reading frame coding for a polypeptide of 363 amino acids. Expression of this clone in *E. coli* in an expression vector showed PTPase activity. The non-catalytic region of this PTPase located at the carboxy terminus shows homology with the basic domains of transcription factors Fos and Jun. Northern blot analysis showed that a 1.7 kb transcript was present in many tissues and cells, the highest level being in macrophages. This PTPase is a rat homolog of human T-cell PTPase although it shows 3 large deletions in the carboxy terminal non-catalytic region.

Protein-tyrosine phosphatase; cDNA cloning; Gene expression; Transcription factor

1. INTRODUCTION

Protein phosphorylation at tyrosine residues has been implicated as an important regulatory component in cell growth, differentiation, malignant transformation by certain viruses and in signal transduction [1-4]. The level of phosphorylation of substrate proteins is determined by the relative activities of protein kinases and protein phosphatases. The study of protein phosphatases in general has received much less attention as compared to that of protein kinases. This was mainly due to the belief that protein phosphatases simply reverse the effects of protein kinases by dephosphorylating the substrates constitutively. Recent findings suggest that this may not be the case with many protein-serine phosphatases as well as with PTPases [5,6], certain cell cycle genes and a transcriptional regulatory gene code for protein-serine phosphatases [5]. The CD45 molecule which is a receptor-like transmembrane protein, has been found to have PTPase activity [7,8]. These findings suggest that phosphatases themselves might play central and specific roles in cellular physiology. The role of non-receptor-type PTPases, their mode of regulation and the substrates on which they act are also not known. PTPase activity is widely distributed in various tissues [9,10]. Rat spleen and brain are a rich source of this activity [9,10]. The genes for several receptor-type PTPases have been isolated [11-15]. The

non-receptor-type PTPases which have been analysed for their cDNA include placental PTPase 1b [16,17], PTPase 1 from rat brain [18] and a PTPase from T cells [19]. An essential virulence determinant of *Yersinia* (a bacterium which is a causative agent of plague) has been shown to be a PTPase [20]. Here we report the isolation of a cDNA clone coding for a PTPase of 363 amino acids which shows homology in its non-catalytic domain with the basic domains of transcription factors Fos and Jun which are required for binding to DNA.

2. MATERIALS AND METHODS

2.1. Construction and screening of cDNA library

A cDNA library was constructed in λ gt11 from poly (A)⁺RNA using cDNA synthesis and cDNA cloning systems obtained from Amersham. Total RNA was isolated by the acid guanidinium thiocyanate method [21]. Poly (A)⁺RNA was prepared by two cycles of binding to oligo (dT) cellulose [22]. The unamplified cDNA library was screened [22] using nick translated *Eco* RI-*Pst* I fragment of T-cell PTPase cDNA [20]. T-cell PTPase cDNA was kindly provided by Dr D.E. Cool (University of Washington, Seattle, USA). The hybridization for screening the cDNA library was carried out at 60°C with 5 × SSPE (1 × SSPE = 0.15 M NaCl, 10 mM sodium phosphate, pH 7.4, 1 mM EDTA), 0.1% SDS, 5 × Denhardt's (0.1% Ficoll, 0.1% polyvinyl pyrrolidone, 0.1% BSA), 0.1 mg/ml denatured salmon sperm DNA [22]. Filters were washed at 50°C in 1 × SSPE, 0.1% SDS.

2.2. DNA sequencing

Nucleotide sequence of the cDNA inserts was determined using the dideoxy chain-termination method of Sanger [23] as modified for double stranded DNA by Chen and Seeburg [24]. The cDNA inserts released from λ gt11 clones were subcloned in plasmid pGEM3Z for sequencing. The complete nucleotide sequence of the full length cDNA clone was assembled by sequencing various restriction fragments and smaller clones.

Correspondence address: G. Swarup, Centre for Cellular and Molecular Biology, Uppal Road, Hyderabad 500 007, India

Abbreviations: PTPase, Protein-tyrosine phosphatase; PTP-S, PTPase cloned from a rat spleen cDNA library

2.3. Northern blotting

Poly (A)⁺ RNA (5 µg) or total RNA (10 µg) was fractionated under denaturing conditions in formaldehyde gels and transferred to nitrocellulose [22]. Hybridization was carried out with 5 × SSPE, 5 × Denhardt's, 0.1% SDS, 100 µg/ml salmon sperm DNA, 30% formamide at 50°C for 16 h. Blots were washed with 0.5 × SSPE, 0.1% SDS at 50°C and kept for autoradiography at -70°C with intensifying screens.

2.4. Expression of PTPase in *E. coli*

The plasmid PTP-S was digested with *Bam* HI and filled in by using Klenow fragment of DNA polymerase I followed by religation. Transformation of competent *E. coli* cells (DH5α) gave a plasmid without *Bam* HI site and extra 4 nucleotides. This was confirmed by sequence analysis. This plasmid was digested with *Hind* III and the *Hind* III fragment was electrocloned and ligated with the *Hind* III digested expression vector pKK233-2 (Pharmacia). The plasmid containing the insert in correct orientation (pKK-PTP) showed PTPase activity. Orientation of the insert was checked by restriction digestion. Assay for PTPase activity was carried out using ³²P-labelled poly (Glu, Tyr) as substrate as described previously [10].

3. RESULTS AND DISCUSSION

A rat spleen cDNA library (λgt11) was screened with the *Eco* RI-*Pst* I fragment (1.38 kb) of human T-cell PTPase cDNA. This fragment contains the entire coding region for PTPase. The screening of over 200 000 recombinants produced 3 positive clones, one of which (PTP-S) contained a full length cDNA insert of 1.5 kb (Fig. 1). The other two clones were smaller in size and corresponded to nucleotides 254-884 (clone 2) and 750-1287 (clone 3) of the largest clone. The nucleotide sequence showed an open reading frame coding for a polypeptide of 363 amino acids. The first ATG codon may serve as the translation initiation codon at position 26 although the sequence surrounding it does not match perfectly with the proposed translation start consensus sequence A/GNNATGG [25]. The open reading frame is terminated by a stop codon TAA at nucleotide 1115. A polyadenylation signal AATAAA is present at nucleotide 1368.

The PTP-S shows a high level of homology with human T-cell PTPase in the catalytic domain [19]. Out of the first 275 amino acids which include the catalytic domain, 261 are identical (95% homology) in the two proteins. This high level of homology suggests that PTP-S may be a rat homolog of T-cell PTPase. A comparison of the carboxy-terminal sequence of PTP-S in the non-catalytic domain with human T-cell PTPase (Fig. 2) shows that there are 3 large deletions after amino acid 288 (19 amino acids), 308 (5 amino acids) and 363 (28 amino acids). These carboxy-terminal differences are perhaps not due to cloning artefacts since one other independent clone also had the same sequence (clone 3, nucleotides 750-1287). The C-terminal differences between PTP-S and human T-cell PTPase may be due to alternative splicing.

In the carboxy-terminal region outside the catalytic domain, there is a region rich in basic amino acids. This basic region is organised as cluster-spacer-cluster in the

same pattern as found in many sequence specific DNA-binding proteins which function as transcription factors [27]. An alignment of the carboxy-terminal 56 amino acids of the PTP-S with segments of the Fos [28] and Jun [29] proteins containing the basic domain is shown in Fig. 3. Out of 37 residues of Fos protein, 18 are identical with PTP-S and 5 other residues are similar in character. Jun protein shows 15 residues (out of 30) which are identical with PTP-S and 6 other residues are similar. The level of homology was also assessed by comparing the alignment scores obtained by using the program ALIGN as described by Dayhoff et al. [26]. These scores were similar for each of the 3 pairs; PTP-S with Fos (3.34), PTP-S with Jun (4.18) and Fos with Jun (3.49). Thus the level of homology between PTP-S and Fos (or Jun) is the same as between Fos and Jun in the basic regions. The gene coding for PTP-1 (or placental PTPase 1b) does not possess any such basic domains in the non-catalytic region of the molecule as shown in Fig. 2. However this basic region is conserved in human T-cell PTPase (Fig. 2.).

In order to confirm that this cDNA clone codes for enzymatically active PTPase we have expressed it in *E. coli*. The cDNA was cloned after appropriate modification (to align the reading frame) in *E. coli* expression vector pKK 233-2 which uses a hybrid promoter *trc*. The strategy for construction of pKK-PTP is described in section 2. The *E. coli* cell extracts containing control plasmid do not show any detectable PTPase activity whereas cells expressing pKK-PTP showed significant PTPase activity (Fig. 4). Complete dephosphorylation of the substrate could be obtained if time of incubation or the amount of extract was increased (not shown).

Expression of the transcripts for PTP-S was studied by Northern blot analysis of mRNA or total RNA isolated from various rat tissues. A major transcript of about 1.7 kb was observed in spleen, brain and thymus (Fig. 5A) and in some other tissues. The size of this transcript is in agreement with the size of the cDNA insert. When an identical RNA blot was analysed by a full length T-cell PTPase probe, a 1.7-kb transcript was observed; the result was same as that obtained with PTP-S probe (figure not shown). These results suggest that PTP-S is the rat homolog of human T-cell PTPase. The high level of homology between the two supports this conclusion.

The expression of PTP-S transcript was compared in splenic lymphocytes with that in peritoneal macrophages. Surprisingly macrophages showed a much higher level of PTP-S transcripts than lymphocytes (Fig. 5C). The results of expression studies suggest that this PTPase does not show tissue- or cell-type specificity although levels of transcription show variability.

In order to see the effect of an inhibitor of protein synthesis on the expression of PTP-S transcripts, rats were injected with cycloheximide (50 mg/kg body weight) and RNA was isolated from spleen and thymus

G TTC CGC TCG CGC TCC CCC GGC CCC ATG TCG GCT ACC ATC GAG CGG GAG TTC GAG GAA	50
Met Ser Ala Thr Ile Glu Arg Glu Phe Glu Glu	11
CTG GAT GCT CAG TGT CGC TGG CAG CCG TTA TAC TTG GAA ATT CGA AAT GAA TCC CAT GAC	118
Leu Asp Ala Gln Cys Arg Trp Gln Pro Leu Tyr Leu Glu Ile Arg Asn Glu Ser His Asp	31
TAT CCT CAT AGA GTG GCC AAG TTT CCA GAA AAC AGA AAT CGA AAC AGA TAC AGA GAT GTA	178
Tyr Pro His Arg Val Ala Lys Phe Pro Glu Asn Arg Asn Arg Asn Arg Tyr Arg Asp Val	51
AGC CCA TAT GAT CAG AGT CGT GTT AAA CTG CAG AGT GCT GAA AAT GAT TAT ATT AAT GCC	238
Ser Pro Tyr Asp His Ser Arg Val Lys Leu Gln Ser Ala Glu Asn Asp Tyr Ile Asn Ala	71
AGC TTA GTT GAC ATA GAA GAG GCA CAA AGA AGT TAC ATC TTA ACA CAG GGC CCA CTT CCT	298
Ser Leu Val Asp Ile Glu Glu Ala Gln Arg Ser Tyr Ile Leu Thr Gln Gly Pro Leu Pro	91
AAC ACG TGC TGC CAT TTC TGG CTC ATG GTG TGG CAG CAA AAG ACC AGA GCA GTT GTC ATG	358
Asn Thr Cys Cys His Phe Trp Leu Met Val Trp Gln Gln Lys Thr Arg Ala Val Val Met	111
CTA AAC CGA ACT GTA CAG AAA GAA TCG GTT AAA TGT GCA CAG TAC TGG CTA ACG GAT GAC	418
Leu Asn Arg Thr Val Glu Lys Glu Ser Val Lys Cys Ala Gln Tyr Trp Pro Thr Asp Asp	131
CGA GAG ATG GTG TTT AAG GAA ACA GGA TTC AGC GTG AAG CTC TTA TCT GAA GAT GTG AAA	478
Arg Glu Met Val Phe Lys Glu Thr Gly Phe Ser Val Lys Leu Leu Ser Glu Asp Val Lys	151
TCA TAT TAT ACA GTA CAT CTA CTA CAG TTA GAA AAT ATC AAT AGT GGT GAA ACC AGA ACC	538
Ser Tyr Tyr Thr Val His Leu Leu Gln Leu Glu Asn Ile Asn Ser Gly Glu Thr Arg Thr	171
ATA TCT CAC TTT CAT TAT ACC ACC TGG CCA GAT TTT GGC GTT CCG GAG TCA CCA GCT TCA	598
Ile Ser His Phe His Tyr Thr Thr Trp Pro Asp Phe Gly Val Pro Glu Ser Pro Ala Ser	191
TTC CTA AAT TTC TTG TTT AAA GTT AGA GAA TCT GGT TCT TTG AAC CCT GAC CAT GGG CCT	658
Phe Leu Asn Phe Leu Phe Lys Val Arg Glu Ser Gly Ser Leu Asn Pro Asp His Gly Pro	211
GCA GTG ATC CAT TGC AGT GCA GGC ATC GGG CGT TCT GGC ACC TTC TCT CTT GTA GAT ACC	718
Ala Val Ile His Cys Ser Ala Gly Ile Gly Arg Ser Gly Thr Phe Ser Leu Val Asp Thr	231
TGT CTC GTT CTG ATG CAG AAA GGA GAG GAT GTT AAT GTG AAA CAA ATA TTA CTG AGT ATG	778
Cys Leu Val Leu Met Glu Lys Gly Glu Asp Val Asn Val Lys Gln Ile Leu Leu Ser Met	251
AGA AAG TAT CGA ATG GGA CTC ATT CAG ACT CCG GAC CAG CTC AGA TTC TCC TAC ATG GCC	838
Arg Lys Tyr Arg Met Gly Leu Ile Gln Thr Pro Asp Gln Leu Arg Phe Ser Tyr Met Ala	271
ATA ATA GAA GGA GCA AAG TAT ACA AAA GGA GAT TCA AAT ATA CAG AAC AGA ACA ATG ACT	898
Ile Ile Glu Gly Ala Lys Tyr Thr Lys Gly Asp Ser Asn Ile Gln Asn Arg Thr Met Thr	291
GAG AAG TAC AAC GGG AAG AGA ATA GGG TCA GAA GAT GAA AAG TTA ACA GGA CTT TCT TCT	958
Glu Lys Tyr Asn Gly Lys Arg Ile Gly Ser Glu Asp Glu Lys Leu Thr Gly Leu Ser Ser	311
AAG GTT CCA GAT ACT GTG GAA GAG AGC AGT GAG AGT ATT CTC CGG AAA CGC ATT CGA GAG	1018
Lys Val Pro Asp Thr Val Glu Glu Ser Ser Glu Ser Ile Leu Arg Lys Arg Ile Arg Glu	331
GAT AGA AAG GCT ACA ACC GCT CAG AAG GTG CAG CAG ATG AGA CAG AGG CTA AAT GAA ACT	1078
Asp Arg Lys Ala Thr Thr Ala Gln Lys Val Gln Gln Met Arg Gln Arg Leu Asn Glu Thr	351
GAA CGG AAA AGG AAA AGG CCA AGA TTG ACA GAC ACC TAA ATG TTC ATG ACT TGA GAC TAT	1138
Glu Arg Lys Arg Lys Arg Pro Arg Leu Thr Asp Thr	363
TCT GCA GCT ATA AAT TTT GAA CCT TTG ATG TGC AAA GCA AGA CCT GAA GCC CAC TCC GGA	1198
AAC TAA AGT GAG GCT TGC TAA CCC TGT AGA TTG CCT CAC AAG TTG TCT GTT TAC AAA GTA	1258
AGC TTT ACA TCC AGG GGA TGA AGA ACG CCA CCA GCA GAA GAC TTG CAA ACC CTT TAA TTT	1318
GAC GTA TTG TTT TTT AAC ATG TGT ATG AAT TGT AGA AAG ATG TAA AGA <u>AAA TAA</u> AAT TAG	1378
GAG AGA CTA CTT TGT ATT GTA CTG CCA TTC CTA ATG TAT TTT TAT ACT TTT TGG CAG CAT	1438
TAA ATA TTT TTA TTA AAT AGA CAA AAA AAA AAA AAA AAA AAA AAA AAA AAA	1495

Fig. 1 Nucleotide and deduced amino acid sequence of cDNA coding for PTPase isolated from a rat spleen cDNA library (PTP-S).

(PTP-S)	GAKYTKGDSNIQNR-----TMTEKYNGKRIGSEDEKLTG-----LS	310
(PTP-T)	...CI...S...K.WKELSKEDLSPAFDHPSPNKI.....N...L.E.....DRCTG..	334
(PTP-1)	...FIM...SV.DQWKELSHEDLEPPPEHVPPPPRPP.RTLEPHNGKCKELFSNHQWVSE	336
(PTP-S)	SKVPDIVEESSESILRKRIRDRKATTAQKVQQRORLNETERKRRPRLTDT	363
(PTP-T)	..MQ..M..N...A.....K.....H.....HLYWQPIILTKMGF	394
(PTP-1)	ESCE.EDILAR.ESRAPS.AVHSMSSMS.DTEVRKRHVGGGLQSAQASVP.EEELSPTEE	396

Fig. 2 A comparison of amino acid sequence of PTP-S with T-cell PTPase (PTP-T) and rat brain PTPase (PTP-1) in the carboxy-terminal non-catalytic region. Gaps introduced for alignment are indicated by dashes. Dots indicate identical residues.

PTPase activity is present in purified rat liver and brain nuclei (unpublished observations from our laboratory). The strong conservation of clusters of basic amino acids in the non-catalytic region suggests that these residues have an important function.

REFERENCES

- [1] Hunter, T. and Cooper, J.A. (1985) *Annu. Rev. Biochem.* 54, 897-930.
- [2] Ullrich, A. and Schlessinger, J. (1990) *Cell*, 61, 203-212.
- [3] Gould, K.L. and Nurse, P. (1989) *Nature* 342, 39-45.
- [4] Lewin, B. (1990) *Cell* 61, 743-752.
- [5] Cyert, M.S. and Thorner, M.S. (1989) *Cell* 57, 891-893.
- [6] Hunter, T. (1989) *Cell* 58, 1013-1016.
- [7] Charbonneau, H., Tonks, N.K., Kumar, S., Diltz, C.D., Harrylock, M., Cool, D.E., Krebs, E.G., Fisher, E.H. and Walsh, K.A. (1989) *Proc. Natl. Acad. Sci. USA* 86, 5252-5256.
- [8] Tonks, N.K., Charbonneau, H., Diltz, C.D., Fisher, E.H. and Walsh, K.A. (1988) *Biochemistry* 27, 8695-8701.
- [9] Shriner, C.L. and Brautigan, D.L. (1984) *J. Biol. Chem.* 259, 11383-11390.
- [10] Swarup, G. and Subrahmanyam, G. (1989) *J. Biol. Chem.* 264, 7801-7808.
- [11] Ralph, S.J., Thomas, M.L., Morton, C.C. and Trowbridge, I.S. (1987) *EMBO J.* 6, 1251-1257.
- [12] Streuli, M., Krueger, N.X., Hall, L.R., Schlossman, S.F. and Saito, H. (1988) *J. Exp. Med.* 168, 1523-1530.
- [13] Streuli, M., Krueger, N.X., Tsai, A.Y.M. and Saito, H. (1989) *Proc. Natl. Acad. Sci. USA* 86, 8698-8702.
- [14] Sap, J., D'Eustachio, P., Civoli, D. and Schlessinger, J. (1990) *Proc. Natl. Acad. Sci. USA* 87, 6112-6116.
- [15] Kaplan, R., Morse, B., Huebner, K., Croce, C., Howk, R., Ravera, M., Ricca, G., Jaye, M. and Schlessinger, J. (1990) *Proc. Natl. Acad. Sci. USA* 87, 7000-7004.
- [16] Chernoff, J., Schievella, A.R., Jost, C.A., Erikson, R.L. and Neel, B.G. (1990) *Proc. Natl. Acad. Sci. USA* 87, 2735-2739.
- [17] Brown-Shimer, S., Johnson, K.A., Lawrence, J.B., Johnson, C., Bruskin, A., Green, N.R. and Hill, D.E. (1990) *Proc. Natl. Acad. Sci. USA* 87, 5148-5152.
- [18] Guan, K., Haun, R.S., Watson, S.J., Geahlen, R.L. and Dixon, J.E. (1990) *Proc. Natl. Acad. Sci. USA* 87, 1501-1505.
- [19] Cool, D.E., Tonks, N.K., Charbonneau, H., Walsh, K.A., Fisher, E.H. and Krebs, E.G. (1989) *Proc. Natl. Acad. Sci. USA* 86, 5257-5261.
- [20] Guan, K. and Dixon, J.E. (1990) *Science* 249, 553-556.
- [21] Chomczynski, P. and Sacchi, N. (1987) *Anal. Biochem.* 162, 156-159.
- [22] Maniatis, T., Fritsh, E.F., Sambrook, J. (1982) *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Lab., Cold Spring Harbor, New York.
- [23] Sanger, F., Nicklen, S. and Coulson, A.R. (1977) *Proc. Natl. Acad. Sci. USA* 74, 5463-5467.
- [24] Chen, E.Y. and Seeburg, P.H. (1985) *DNA* 4, 165-170.
- [25] Kozak, M. (1986) *Cell* 44, 283-292.
- [26] Dayhoff, M.O., Barker, W.C. and Hunt, L.S. (1983) *Methods Enzymol.* 91, 524-545.
- [27] Busch, S.J. and Sassone-Corsi, P. (1990) *Trends Genet.* 6, 36-40.
- [28] Van Straaten, F., Muller, R., Curran, T., Beveren, C.V. and Verma, I.M. (1983) *Proc. Natl. Acad. Sci. USA* 80, 3183-3187.
- [29] Lamph, M.W., Wamsley, P., Sassone-Corsi, P. and Verma, I.M. (1988) *Nature* 334, 629-631.
- [30] Nath, S.T. and Nayak, D.P. (1990) *Mol. Cell. Biol.* 10, 4139-4145.