

The light-harvesting chlorophyll *a*–*c*-binding protein of dinoflagellates: a putative polyprotein

Roger G. Hiller*, Pamela M. Wrench, Frank P. Sharples

School of Biological Sciences, Macquarie University, Macquarie, NSW 2109, Australia

Received 6 February 1995; revised version received 8 March 1995

Abstract The principle light-harvesting chlorophyll *a*–*c*-binding protein of *Amphidinium carterae* of 19 kDa is encoded as a polyprotein translated from a 6.1 kb mRNA. The cDNA sequences indicate that each derived polypeptide is contiguous with the next and that the mature peptides are formed by cleavage at a C-terminal arginine residue. Comparison of the amino-acid sequences shows the *Amphidinium* protein to be most closely related to the fucoxanthin-chlorophyll-protein (Fcp) of *Phaeodactylum* and less related to the chlorophyll *a*–*b*-binding (Cab) proteins including those from *Euglena*.

Key words: Amino acid sequence; Chlorophyll *c*; Polyprotein; Light-harvesting complex; Dinophyceae; *Amphidinium carterae*

1. Introduction

The intrinsic light-harvesting proteins of higher plants and green algae binding Chl *a* and *b* are encoded by multigene nuclear families [1–3]. Many of the genes are expressed although the specific roles of many of the proteins are by no means clear. In the chromophyte algae the light-harvesting proteins bind Chl *a* and *c* and are also encoded by multigene families [4,5]. All these light-harvesting proteins share conserved motifs although the level of identity is overall rather low. It has, however, been suggested that they share a three transmembrane helix structure. This may indicate a common evolutionary origin involving duplication and fusion of genes encoding a two transmembrane protein followed by elimination of one of the four helices [4].

Peptides comprising the main light-harvesting complex of dinoflagellates have been previously investigated [6,7]. The principle polypeptide of 19 kDa is related at the N-terminal region to the light-harvesting proteins of other chromophytes as well as to the Cab proteins [6]. During PCR experiments designed to confirm synthesis of a cDNA extending to the known N-terminus of the mature 19 kDa peptide, an amplified band of ~1 kb was obtained in addition to that expected at 420 bp. The 1 kb band was found to encode a complete amino acid sequence of a 19 kDa polypeptide flanked by partial sequences encoding two other forms of the 19 kDa polypeptide.

We report here evidence, based on cDNA sequencing and Northern blotting, that a gene for the light-harvesting complex

of dinoflagellates encodes a polyprotein which is posttranslationally cleaved to up to 10 different 19 kDa mature polypeptides.

2. Materials and methods

Amphidinium carterae CS-21 was cultured, its main light-harvesting complex isolated and the 19 kDa principle polypeptide sequenced as previously described [6]. Total RNA was extracted by homogenising exponentially growing cells in Total RNA Isolation Reagent (Advanced Biotechnologies) using 1 ml reagent for 25 µl of packed algal cells. After extraction of the lysed cells with chloroform, RNA was precipitated with isopropanol and washed with 75% ethanol according to the manufacturers' protocol. Precipitated RNA was redissolved in DEPC-treated water and used immediately for poly(A)⁺ mRNA preparation or precipitated by addition of 0.1 vol. of 3 M sodium acetate and 2.5 vol. of ethanol and stored at –20°C until required for Northern blotting.

Poly(A)⁺ mRNA was prepared from total RNA by addition of 0.1 vol. of Oligotex poly-T affinity reagent (Quiagen Inc.) following the manufacturers' protocol. First strand cDNA synthesis was by reverse transcription using the primer 2360 5'-AATTCGCGGCCGC(T)15 and a 5'-Amplifinder RACE kit (Clontech Inc). Fragments of cDNA were amplified by PCR using primers designed from protein sequence data for the 19 kDa principle peptide of the LHC. The forward primer for 1 kb product was 218 5'-TTCGACCCCTCGGTTTCAC corresponding to the amino acid sequence FDLGFT and the reverse primer 682 5'-CAT(ATGC)CC(TGA)AT(TGA)AT(ATGC)GCCATCAT corresponding to the amino acid sequence MMAIHGM. The 3'-end of the cDNA was amplified using as forward primer 2361 5'-CGGTGACTGGGCAACTACA designed to the 3'-end of the 1 kb cDNA product with the reverse primer 2360. Standard PCR conditions were 2 mM Mg, 100 µM dNTP and 2 units each of Taq polymerase and Taq extender (Stratagene) in a total volume of 50 µl. After an initial cycle of 3 min at 94°C, 1 min at 52°C and 2 min at 72°C the PCR reactions were given 34 cycles for 1 min, 1 min and 2 min at the above temperatures. Amplified PCR products were separated on 0.8% agarose gels, excised and purified with GeneClean II (Bio 101) and cloned into the pGEM-T vector (Promega) following the manufacturers protocols. DNA sequencing was by the dideoxy chain termination technique [9] using a Sequenase II kit (USB) and the manufacturer's protocol for double-stranded templates. Both strands were sequenced by a combination of standard subcloning and custom oligonucleotides. DNA and protein sequences were analysed using the Wisconsin program package accessed through the Australian National Genomic Information Service.

For Northern blotting RNA was dissolved in 25 mM EDTA, 0.1% SDS, size-fractionated on a 1.2% agarose gel and probed [8]. The probe for LHC was from the cloned 1 kb PCR fragment (pRGH200) cut with *Sac*I and contained 756 bp of the insert. For PCP the probe was a 1272 bp fragment obtained by genomic PCR encoding 131 C-terminal residues, the intergenic region and the N-terminal leader sequence plus 72 residues of an adjacent PCP (Hiller, Wrench and Sharples, unpublished observations). Probes were labelled using a Gigaprime kit from Biotec.

3. Results

The sequence of the cloned (pRGH200) original 1 kb amplified cDNA band together with a derived amino acid sequence

*Corresponding author. Fax: (61) (2) 850-8245.

Abbreviations: Chl, chlorophyll; LHC, light-harvesting complex; PCP, peridinin chlorophyll *a*-protein; Cab, chlorophyll *a*–*b*-binding; PCR, polymerase chain reaction; Fcp, fucoxanthin-chlorophyll-protein.

The sequences reported here will appear in the EMBL nucleotide data base under numbers Z47562 and Z47563.

FIG 1A

TTCGACCCCCCGGTTTCAAAAGGAGGGCAACGTCGATAACTTCAAGCGGAGGCGAGCT 60
 F D P L G F T K E G N V D N F K R R R A
 ACTGAGATCAAAATGGACGAATTCCTATGCTGCGACCATGGGTACATCACCCCTGAG 120
 T E I K H G R I A M L A T M G Y I T P E
 CTCCTAGGTTTCGATGGCTACCTGTCCGAAGCGCGGTGTGAAGTATGCGGACATTCGA 180
 L F R F D G Y L S R S A G V K Y A D I P
 AATGGCTTGGCTGCTTCTCAAAGGTGCCATTTGTGGGTGGGTGAGATGGTCGCATAC 240
 N G L A A F S K V P F V G W V Q M V A Y
 TGCTTGTGGTTGAGATCTCACAAGATCAGTCGAGGCGCTCCCGCTCAGGGTGACTTT 300
 C L L V E I S Q D Q S Q G A P A Q G D F
 GGCTTTAAGGTCTTGACCTCTTCCGACCCAGCAGAGAAGAGAAGCTCAGAGCTGAG 360
 G F K V L T S S D P A E K E K L R A E
 CTTGCCAACGGGCGCTGGCCATGATGGCCATCATTTGGGATGTCTTCCAGGATGGCTG 420
 L A N G R L A M M A I I G M F F Q D G L
 ACCGGTTCGGCATGGGAGATTGGGCAACTACACGAGTCCCATTTGGGGCGTTTGAG 480
 T G S A W G D W A N Y T E S P L R A F E
 AACGAGCTTGGTGCAAGCACCCACAGGCTTCTTCGACCCCTCGGTCTTTCGTCGGAC 540
 N L V L G V Q A P T G F D P L G L S S D
 GGAAGCGTTGACAACCTTCAAGCGACGTCGTGCAAGCGAGATCAAGCACGCGCGTTGCC 600
 G S V D N F K R R R A S E I K H G R V A
 ATGCTCGCTACTATGGGATACATGACCCCTGAGATAACTGCTAAGTTCCTCGGCTATCTC 660
 M L A T M G Y M T P E I T A K F P G Y L
 TCGTACTCAGAGTAGCAAGTTTTCGGATGTGCCAATGGACTGGCTGCCATGTCCAAG 720
 S Y S Q S S K F A D V P N G L A A M S K
 GTCCCTGTGCTTGGGTGGGCTCAGGTAGCTGCATGTGCTGTCTGCGAGCTGTGCGAA 780
 V P V L G W A Q A Y A G A V C D V S
 GATCAGTCCGAGGACACCTTGGCGCAGCAGGTGACTTCGGCTTCAAGGTGATCACTCA 840
 D Q S A G T P G A A G D F G F K V I T S
 GAAGATGAGAAACCTCAAAGAAAGCTGAATCCGAGCTCGCAACGGCGCGCTGGCA 900
 E D E E T L K R K L N S E L A N G R L A
 ATGATGGCCATCATAGGCTCTTCTTTCAGGATGGTCTTACAGGTGGGCGATACGGTGAC 960
 M M A I I G L F F Q D G L T G G A Y G D
 TGGGCAACTACACGAGCTCTCCATTGCGCGCTTTCGAGAGCGAGCTTGGTGGCGAGCT 1020
 W A N Y T D S P L R A F E S E L G A Q A
 CCAATTGGCTTCTTGTGAAACCGAGGGGCTCGAA
 P V G F F

FIG 1B

CGGTGATGGGCAACTACACTCGCTCTCCCTCGCTGCTTTCAGAGCGAGCTTGGCGT 60
 G D W A N Y T A S P L R A F E S E L G V
 CGCAGCACCCACAGGTTCTGGGACCCATTTGGGCTTTCAGAGCGAGCTTGGCGT 120
 A A P T G F W D P L G L A K D C S M K A
 CTTCAAGAGAAGCGTTCGCTCAGAGATCAAGCATGGTGTATGCAATGTGGCCACTAT 180
 F K R R R A S E I K H G R I A M L A T M
 GGGTACATCACTCTGAGTTGACTGGTAAGTTCACAGGCTACTTGTACCTTCCATGGG 240
 Y I T P E L T G K F P G Y L G S P S M G
 CTTGAAGTACGAGGACATTCCTCAATGGCTTGGGCGCCATCAGCAAGTTCCAGCAGTTGG 300
 L K Y E D I P N G L G A I S K V P A V G
 TTGGGCACAGATCTCGCATACGCCCTTCTACTGTGAGCAGTCCCAAGATCAGTCCGAGGG 360
 W A Q I L A Y A F Y C E Q S Q D Q S E G
 CTCGCGGGTGAGGCGGTGACTTTGGCTTCAAGGTGCTCAGATCTAAGGACGAGGAAGG 420
 S A G E A G D F G F K V L T S K D E E G
 TTTGGAGAGGAAGTTGAATCTGAGATCGCAATGGTGGCTTGGCCATGATGSCAATTAT 480
 L E R K L N S E I A N G R L A M M A I I
 TGGTATGTCTTCCAGGATGGCTTGCAGGCTCAGCCTATGGGAGCTGGGCAACTTCAC 540
 G M F P Q D G L T G S A Y G D W A N F T
 TGCCTCCCACTCCGATAAGCAATCACTTAACCCGAGCTTGGCTTTTTCAGCTCCTGTC 600
 A S P L R *
 GCAAAACAAAAACCGCGCGCTTAA

Fig. 1. Reverse-Transcriptase PCR products encoding Chl *a-c*-binding protein sequences. (A) Product of primers 218/682 (clone pRGH200). (B) Product of primers 2361/2360 (clone pRGH201). Solid lines show the amplifying primer positions and the dashed line the position of primer 2361 in relation to sequence in (A). Arrows show deduced proteolytic cleavage points.

is shown in Fig. 1A. Arrows mark the cleavage points expected from protein sequencing (see Fig. 3 and [6]). We interpret the derived sequence to indicate that a complete mature 19 kDa




Fig. 2. Northern blot of total *Amphidinium* RNA probed with either part of the PCR product shown in Fig. 1A (LHC) or a genomic clone encoding the peridinin-Chl *a*-protein (PCP).

peptide is flanked by part of two other LHC peptides. The untranslated DNA sequence corresponds to the forward PCR primer in reverse and may be a PCR artefact resulting from premature termination of the amplification. To determine the relationship between the sequence of Fig. 1A and poly(A)⁺ tail of the cDNA, we performed a second PCR amplification using in the forward direction an oligonucleotide corresponding to C-terminal region of the complete polypeptide derived from Fig. 1A and in reverse one corresponding to the poly(A)⁺ tail. From this experiment we obtained an amplified product of 630 bp whose sequence is shown in Fig. 1B. This differed in a number of base positions from the corresponding 3' part of the Fig. 1A sequence but the derived amino acid sequence still includes that of a complete mature 19 kDa peptide together with part of the preceding peptide and its C-terminal Arg. We conclude that the results are best explained by the cDNA for the main polypeptide of the light-harvesting complex encoding a polyprotein which is proteolytically cleaved at C-terminal Arg residues to form at least 5 separate mature LHC polypeptides.

To determine the number of potential polypeptides encoded by the gene transcript a Northern blot was performed using as probes a part of the original 1 kb clone and a clone encoding most of the mature PCP, a chloroplast protein which is synthesised as a single precursor with a cleaved N-terminal extension [11]. The results are shown in Fig. 2; the LHC probe hybridises strongly to an mRNA of 6.1 kb whereas the PCP probe hybridises to an mRNA of 1.7 kb. There was weak hybridisation of the LHC probe to species of 2.7 kb and 1.5 kb but no distinct hybridisation to species much greater than 6.1 kb.

The protein-derived sequence of the principle 19 kDa peptide of the purified LHC is shown in Fig. 3 together with its main variants and aligned with the cDNA-derived sequences. The N-terminal X is a modification which does not inhibit the sequential Edman reactions and the resulting PTH-derivative

		S		Q		
a	XFENERGVQD	PVGFFDPLGF	TADGSVENFK	RLAQTEIKHG	RVAMLATMGY	50
b	a-----l--a	-t-----l	ss-----d--	-rras-----	-----	
c	a-s-l--aa	-t--w-----l	ak--mka--	-rras-----	-i-----	
d			-ke-n-d--	-rra-----	-i-----	
		S				
a	ITPEITGKLP	GYLSPSTGVK	YDDIPNGLGA	ISKVPAAGWG	QMIAYAAFSE	100
b	m-----a-f-	-----y-qss-	fa-v-----a-	m-----v1--a-	-va--g-vcd	
c	---l---f-	-----m-l-	-e-----	-----v--a-	-il---fyc-	
d	---l#frfd	---r-a---	-a-----a-	f---fv--v	--v-cllv-	
		I		G	A	
		S	A	A	PI	P
a	LSQDQSAGTP	AAEGDFGFKV	LTSSDPAELE	KKLSAELANG	RLAMMAIIGM	150
b	v-----	g-a-----	i--e-eet-k	r--ns-----	-----l	
c	q-----e-sa	gea-----	---k-eeg--	r--ns-i-----	-----	
d	i-----q-a-	-q#-----	-----k-	-r-----	-----	
a	FFQDGLTGSA	WGDWASYTAS	PL			
b	-----g-	y-----n--d-	--r			
c	-----	y-----nf--	--r*			
d	-----	-----n--e-	--r			

Fig. 3. Amino acid sequences of principle polypeptides of *Amphidinium* LHC. a, protein-derived sequence with major variations in uppercase letters above; b, complete cDNA-derived sequence; c, complete cDNA-derived sequence (3'-end of gene); d, partial cDNA-derived sequence. The masses of peptides b and c are 18,420 and 18,604, respectively. (-, identity; #, a space; *, stop codon).

elutes in the region of PTH-Met. We hypothesise, based on the cDNA derived sequences that X is a modified Ala. (It coelutes with PTH-*n*-methyl-alanine.) The Ser/Asn ambiguity at residue 4 results from approximately equal amounts of the two species and both are specified in the derived sequences. For many of the other variants we have not yet found correspondence in the derived sequences including 3 additional partial DNA sequences obtained by genomic PCR (Hiller and Wrench, unpub-

lished observations). The identity between the protein sequence and the cDNA-derived sequences is 71–78%, which is similar to that between the cDNA-derived sequences themselves.

4. Discussion

As was previously noted [6] comparison of the *Amphidinium* LHC sequence with those of other LHCs (Fig. 4) indicates

	1				50
a
p
eAAGAV
u
t	MASACASSTI	AAVAFSSPSS	RRNGSTVGT	KASFLGGRR	RVSKYSTPT
m
	51				100
aAFENE	RGVQDPVGF	FDPLGFTADG
p*****	I*A*Q*L*Y	W****LV***
e	SKDPFSPLNK	LYGPDVRVKWL	GGFTGFVPEY	LTGELPGDYG	W*TA*LG**P
uYGNRPKWLG	PLSGG*VPEY	LKGEYAGDYG	**TA*LAA*P
t	ARSATTVCVA	ADPDRPLW..	.FPGSTPPPW	LDGSL*GD*G	*****LAS*P
mKSVSTV	VKADIYEPFG	TY*G*GESPI
	101				150
a	SVENFKRLAQ	TEIKHGRVAM	LATMGYITPE	ITGKLP..GY	LSPSTGVKYD
p	DQ*K*D*RY	V*****IC*	**VA**L*Q*	AGIR**..D	IDY*..*TSFE
e	AKLARY*V*	.QVI***W**	*GVL*CWF**	...CINRW*G	VPF.....
u	KLFQRY*D*	.LQN**WA*	*GVL*CLA**	...V*SNVFG	VPYEPVWFK
t	ESLRWNQ*Q*	.LV*C*W**	*GAA*IFI*	...L*TKIGI	*.....
m	IPFGSEKN*E	R*VS***W**	*GVT*AWAA*	NGTGI*WFTA	GTLC*PDDCT
	151				200
a	DIPNGLAIS	KVPAAGWGQM	IAYAAFSELS	QD.....
p	S***FA*L*	A**G**IA*I	**FIG*L*IA	VM.....
eG
u	TGATI*NGG*	IDYL*NPK..	...LIHASN	LLLTLVLELV	FFFAAETWRE
t	NT*SWYT*GE	QEYFTDTT..	...TLFIVE	LVLIGWAEGR	RWADIIPKGC
m	AVADKFPAGAV	APL*PEGSGY	PSFW*VLAIE	VVLVGLAE..
	201				250
aQSAGTPAAE	GDFGFKVL..	...TSSDPAE	LEKKLSAELA
pKDITGGEFV	*DFRNRY*DF	GWD*F*EDKK	*Q*R.AIELN
e	ADLDPL....Y*	*GP.*DP*GL	ADD.PEAF*	LKV*...EVK
u	AGEGFL....GKAQDKSYP	*GV.*DP*GL	SKD.PAFA*	AKV*...EVK
t	VNTDPIFFNN	KLTGTDVGY*	*GLW*DP*GW	GSGSPAKI*	*RT*...EIK
m	AYRTGLTDS	FEELTVGDVS	PGGR*DP*GA	ESG...LE*	*KI*...ELK
	251				300
a	NGRLAMMAII	GMFFQDGLTG	SAWGD.WASY	TASPL.....
p	Q**A*Q*G*L	ALMVHEQ*GV	*ILP.....
e	N*****V*F*	*F*V*GI***	KGPVEN*VDH	LTD*F.VNNI	FQLTPGF...
u	*****L*ML	*LFV*A*VT*	QSPLNLSAH	LAN*G.VNFW	TSYAPTL...
t	N*****L*VM	*AW**HIY**	TGPI*NLFAH	L*D*GHATIF	AAFSFK....
m	HCR*S*F*WL	*CI**GLA*Q	EGPIAN*QAH	V*D*VHANVL	TNAASGFGFY

Fig. 4. Comparison of Cab and Chl *a*-*c*-binding protein sequences aligned using the PILEUP program. a, *Amphidinium* LHC (this paper); p, *Phaeodactylum* Fcp [5]; e and u, *Euglena* LHCPII [9]; t, Tomato LHCPI [11]; m, *Mantoniella* LHCP [12]. (*, identity; ., a space.)

significant identity especially towards the N-terminus. The greatest identity is with *Phaeodactylum* Fcp (39%) whereas it is less than 30% with those of the Cab proteins. The atypical LHC of *Mantoniella* which binds Chl *c* as well as Chl *a* and *b* is only 22% identical and is clearly not part of the chromophyte evolutionary line as previously noted [13]. In a preliminary analysis of the *Amphidinium* sequence [6] we failed to find evidence based on the FDPLG motif for duplication of the sequences which include the postulated first and third transmembrane helices. Such a duplication, although not as strong as for the Cab proteins, is now apparent as shown below:

SVENFKRLAQTEIKHGRVAMLATMGYITPEIT

: : : : : :

AELEKKLSA--ELANGRLAMMAIIGMFFQDGL

A feature of the *Amphidinium* amino acid sequences is a high variability in Arg and Lys residues and the resulting multiplicity of peptides following tryptic digestion accounts in part for the shortcomings of our previous analysis.

There is only one comparable case (*Euglena* [10]) of an LHC being synthesised as a polypeptide and posttranslationally cleaved to multiple forms of the mature peptide. Taking *Euglena* LHCP II as a model [10], *Amphidinium* LHC mRNA of 6.1 kb would be made up of approximately 1 kb of the leader sequence, non-coding 5' and 3' regions plus the poly(A)⁺ tail. The remaining 5.1 kb could comprise up to ten 516/519 bp pseudo-repeats which encode the 172/173 amino acid LHC polypeptides. A major difference exists between the *Amphidinium* polypeptide, which is cleaved at an Arg residue without liberation of spacer peptide and the *Euglena* polypeptide. In *Euglena* the processing protease(s) recognises the consensus sequence XMXAXGXKX and liberates the corresponding decameric peptide. A feature of the gene encoding *Euglena* LHCP II is the occurrence of numerous introns which are flanked by atypical base sequences [10]. If the *Amphidinium* LHC gene contains introns they are likely to be fewer than in *Euglena* as Fig. 2 did not show any hybridisation to distinct species greater than 6.1 kb even after extended exposure. However, it is worth noting that the PCR artefact, which gave our original 1 kb PCR fragment (where the forward 218 oligonucleotide primer also functioned in reverse), has also been observed when the 218/682 primer combination was used for genomic PCR although in this case the site was quite different (Hiller and Wrench, unpublished observations). The last two 3' bases of primer 218 are AC which would hybridise to the GT 5' consensus of typical eukaryotic splice junctions.

Despite the low identity of *Amphidinium* LHC with LHCs containing Chl *b*, alignment of the sequence with the 3-dimensional structure of Kühlbrandt et al. [14] gives some insight into the binding of pigments. Within the sequences comprising the

first and third (B and A in the Kühlbrandt model) helices there are five sites binding Chl *a* which are identical. These are the Arg/Glu combinations which bind Chl *a*₄ and Chl *a*₁, the His binding Chl *a*₅, the Asn binding Chl *a*₂ and the Gln binding Chl *a*₃. It may be expected, due to the highly conserved nature of the inner antennae of the photosystems, that the position of some Chl *a* molecules, especially those at the lowest energy levels, will be common to LHCs in all organisms if the three transmembrane model is universally applicable. There may also be non-conserved Chl *a* binding sites if specific LHC peptides can only participate in energy transfer within the LHC.

Most phylogenetic dendrograms have euglenoids well separated from dinoflagellates [15,16] despite both having three membranes surrounding the chloroplast and we suggest that the unusual transport and processing of their respective LHCs has arisen independently. However, dinoflagellates may well be polyphyletic with respect to their plastids [16] and *Amphidinium carterae* may not be representative of all dinoflagellates.

Acknowledgements: We thank Dr. J. Speirs for assistance with Northern blotting, Dr. A.A. Gooley of Macquarie University Centre for Analytical Biotechnology for protein sequencing and Prof. Tony Larkum for commenting on the manuscript. The work was supported by an Australian Research Council Grant (A19131108) to R.G.H.

References

- [1] Thornber, J.P., Morishige, D.T., Anandan, S. and Peter, G.F. (1991) in: *Chlorophylls* (Scheer, H. ed.) pp. 549–585, CRC Press, Boca Raton, FL, USA.
- [2] White, M.J., Fristensky, B.W., Falconet, D., Childs, L.C., Watson, J.C., Alexander, C., Roe, B.A. and Thompson, W.F. (1992) *Planta* 188, 190–198.
- [3] Di Paolo, M.L., Dal Belin-Peruffo, A. and Bassi, R. (1990) *Planta* 181, 275–286.
- [4] Green, B.R. and Pichersky, E. (1994) *Photosynthesis Res.* 39, 149–162.
- [5] Grossman, A.R., Manodori, A. and Snyder, D. (1990) *Mol. Gen. Genet.* 224, 91–100.
- [6] Hiller, R.G., Wrench, P.M., Gooley, A.A., Shoebridge, G. and Breton, J. (1993) *Photochem. Photobiol.* 57, 125–131.
- [7] Inglesias-Prieto, R., Govind, N.S. and Trench, R.K. (1993) *Phil. Trans. R. Soc. Lond. B* 340, 381–392.
- [8] Fourney, R.M., Miyakoshi, J., Day III, R.S. and Paterson, M.C. (1988) *Focus* 10, 5–7.
- [9] Sanger, F., Nichlen, S. and Coulson, A.R. (1977) *Proc. Natl. Acad. Sci. USA* 74, 5463–5467.
- [10] Muchhal, U.S. and Schwartzbach, S.D. (1992) *Plant Mol. Biol.* 18, 287–299.
- [11] Norris, B.J. and Miller, D.J. (1994) *Plant Mol. Biol.* 24, 673–677.
- [12] Pichersky, E. and Green, B.R. (1990) in: *Progress in Photosynthesis Research*, vol III (Baltscheffsky, M. ed.) pp. 553–556, Kluwer, Dordrecht.
- [13] Rheil, E. and Morschel, E. (1993) *Mol. Gen. Genet.* 240, 403–413.
- [14] Kühlbrandt, W., Wang, D.N. and Fujiyoshi, Y. (1994) *Nature* 367, 614–621.
- [15] Cavalier-Smith, T. (1989) *Nature* 339, 100–101.
- [16] Martin, W., Somerville, C.C. and Loiseaux-de Goer, S. (1992) *J. Mol. Evol.* 35, 385–404.