

## Minireview

## Diverse molecular interactions of the hnRNP K protein

Karol Bomsztyk\*, Isabelle Van Seuningen<sup>1</sup>, Hideaki Suzuki, Oleg Denisenko, Jerzy Ostrowski<sup>2</sup>

Department of Medicine, University of Washington, Seattle, WA 98195, USA

Received 28 December 1996

**Abstract** The hnRNP K protein is a versatile molecule that interacts with RNA, DNA, the proto-oncoprotein Vav, Src-like tyrosine and inducible serine/threonine kinases, the transcription factor TBP and a number of zinc-finger transcriptional repressors. The interaction of K protein with some of its protein partners is modulated by nucleic acids and K protein can alter the in vivo and in vitro rate of transcription. K protein can simultaneously engage several proteins and may facilitate molecular cross-talk. Taken together these diverse interactions suggest that K protein may act as a nucleic acid-regulated docking platform.

© 1997 Federation of European Biochemical Societies.

**Key words:** K protein; Docking platform; KH domain

### 1. K protein primary structure

K protein was first discovered as a component of the heterogeneous nuclear ribonucleoprotein (hnRNP) particle [1] from where it derives its name. K protein is encoded by a gene mapped in humans to chromosome 9 [2]. It produces at least five proteins that are products of alternatively spliced transcripts [2]. The deduced molecular weight of the isoforms is in the range of 50–51 kDa but in SDS-PAGE, the K protein has an apparent molecular weight of 65 kDa. The D isoform of K protein has been cloned from human and mouse cDNA libraries [2,3]. On the amino acid level, this isoform in the two species differs by only one conservative substitution: residue 350 in the mouse is Glu and Asp in the human protein. K protein contains the evolutionarily conserved KH repeats (Fig. 1) that are almost completely conserved between *Xenopus laevis* and mammals [4]. KH-like domains are also found in RNA-binding proteins in species as diverse as *Escherichia coli* and *Saccharomyces cerevisiae* [5]. Mutation in the KH domain of the human FMR1 protein is responsible for the fragile X syndrome, the most common hereditary mental retardation disorder [6,7]. The human and murine K protein sequences contain four GRGG repeats, two of which are conserved in *X. laevis*. Human, mouse and *X. laevis* sequences contain RGRP repeats and a cluster of three proline-rich SH3-binding domains [4,8,9] (Fig. 1).

In addition to these highly evolutionarily conserved features, in the mammalian K protein, the carboxyl end of the

SH3-binding cluster is adjacent to a domain that recruits an interleukin-1 (IL-1)-responsive K protein kinase [9]. The N-terminus of the K protein is highly acidic and has been reported to contain transcriptional activity [10]. The K protein region that encompasses amino acids 206–327 and contains the GRGG box and the SH3-binding domain, binds the transcriptional repressor, Zik1 [11] (Fig. 1).

### 2. Interaction of K protein with nucleic acids

The function of K protein in the hnRNP particle has not yet been defined, but it is easily recovered from both cytoplasmic and nuclear extracts indicating that it has a wide intracellular distribution [2,8] and that a large fraction of K protein is not associated with the hnRNP particle. K protein also binds in a sequence-selective fashion to RNA [1,12], and to a single- [8,13–15] and double-stranded [8,16] DNA. Poly(C) RNA is strongly bound by K protein but other RNA homopolymers are bound poorly or not at all [1,12]. The binding of K protein to RNA might involve the KH domains [6], and binding affinity to RNA is diminished when K protein is phosphorylated [2].

A number of specific interactions of K protein with distinct motifs in DNA have also been observed. For example, K protein binds the homopurine/homopyrimidine (CCCC/GGGG) tract present in the CT motif nested within the *c-myc* promoter P1 [16] and the  $\kappa$ B motif [8]. Like the CT motif, the  $\kappa$ B motif also contains a homopyrimidine/homopurine domain but in the opposite orientation. K protein also binds tenaciously to oligo(dC), which might account for its binding to the late-coding simian virus 40 DNA strand [17].

### 3. Interaction of K protein with factors involved in signal transduction and gene expression

K protein binds and is phosphorylated in vivo and in vitro by an inducible serine/threonine kinase, K protein kinase, KPK [9,12,18]. The identity of KPK remains to be defined. Phosphorylation of K protein by KPK is modulated by its binding to cognate DNA or RNA motifs. In addition, K protein exists in a complex with a number of important signal transduction molecules. For example, K protein has been shown to bind selectively in vitro to the SH3 domains from Src, Fyn and Lyn and it can be co-immunoprecipitated with c-Src from cell extracts, suggesting that K protein-tyrosine kinase complexes may exist in vivo [9]. Even more interestingly, in the context of K protein, c-Src can reactivate the KPK in vitro, suggesting that K protein can facilitate cross-talk between protein tyrosine and serine/threonine kinases. K protein has also been demonstrated to bind in vivo and in vitro the proto-oncoprotein Vav via SH3 interactions [9,19].

\*Corresponding author. Fax: (1) (206) 685-8661.  
E-mail: karolb@u.washington.edu

<sup>1</sup>Present address: Unité INSERM 377, Place de Verdun, 59045 Lille Cedex, France.

<sup>2</sup>Present address: Oncology Center, 02-781 Warsaw, Poland.

Vav is a key element of the apparatus for signaling through both the T- and B-cell antigen receptors [20,21], demonstrating involvement in yet another signaling pathway. Finally, K protein and KPK co-immunoprecipitate from cell extracts with either c-Src or Vav, suggesting that K protein permits multienzyme complex formation *in vivo* [9].

One clue for the involvement of K protein in transcription is observation that K protein co-immunoprecipitates with epitope-tagged TBP from nuclear extracts. [<sup>35</sup>S]TBP translated *in vitro* binds to recombinant K protein, suggesting that *in vivo* K protein can interact with TBP directly [22]. K protein also binds *in vivo* and *in vitro* a zinc-finger transcriptional repressor, Zik1, an interaction that is likewise direct [11]. The interaction of K protein with Zik1 may reflect a general interaction that exists between Kox1-like zinc-finger transcriptional repressors and K protein because Kid-1 [23] and MZF-1 [24], transcriptional repressors structurally related to Zik1, can also be directly engaged by K protein (Denisenko and Bomsztyk, unpublished results). Importantly, the interaction of K protein with the transcriptional repressors appears to be regulated by cognate nucleic acid motifs. For example, poly(C) RNA, which binds K protein, blocks Zik1-K protein interaction, while poly(A) RNA, which does not bind K protein, has no effect on this interaction. K protein-Zik1 interaction can also be blocked by the κB motif, which like poly(C) RNA also binds K protein [11].

There are also functional studies that implicate K protein in the regulation of transcription. For example, K protein has been shown to increase transcription *in vivo* [22,25] as well as *in vitro* [17]. However, the mechanisms by which K protein alters gene transcription are not yet clear because K protein can both activate and repress transcription of reporter gene linked to a heterologous promoter even when that promoter does not contain any obvious K protein-binding sequence-specific DNA motifs [17,22].

#### 4. Perspectives

The most evolutionarily conserved characteristic of K protein is its ability to bind RNA through the KH domains, a motif that occurs in organisms as distant from mammals as bacteria [5]. This extraordinary conservation suggests a very fundamental role of K protein in the processes involving RNA. K protein interacts with a host of protein partners (Table 1) and notably the interaction of K protein with some of these molecular partners is modulated by RNA [8,9,11,12]. Thus, K protein may serve as a RNA linker molecule that would allow RNA to exert effects on protein factors that do not bind nucleic acids directly. As a linker molecule

Table 1  
K protein molecular partners

Molecules involved in signal transduction	
	Tyrosine kinases- Src, Fyn and Lyn
	Proto-oncoprotein Vav
	Serine/threonine inducible kinases
Molecules involved in gene expression	
	TFIID TATA-binding protein (TBP)
	Transcriptional repressors- Zik-1, Kid-1 and MZF-1
	Elongation factor-1α
	Sequence-specific single- and double-stranded DNA
	Sequence-specific RNA




Fig. 1. Diagrammatic illustration of K protein modular domains. Acidic domain is contained in aa 1–40; KH 1 aa 46–98; KH 2 aa 149–197; KH 3 aa 391–439; GRGG box aa 236–273; Src SH3-binding domain aa 289–315; Zik1-binding domain aa 209–337; K protein kinase (KPK)-binding domain aa 337–425.

for nucleic acids, K protein would be expected to interact with tens if not hundreds of different protein factors. Because K protein contains two different dimerization domains (Denisenko, Van Seuning and Bomsztyk, unpublished observations), it may, *in vivo*, form oligomers. If so, K protein could assemble multiple factors on RNA and allow multilateral cross-talk that could be regulated by nucleic acids to which K protein binds. Thus K protein may act as an RNA- and DNA-regulated docking platform. As such, K protein would link sequence-specific nucleic acids to protein-protein interactions. Serving as a nucleic acid-linking docking platform, K protein could act in processes such as transcription and translation. Such a role would be consistent with the observation that K protein alters rates of transcription *in vivo* and *in vitro* [10,14,25]. The effect of K protein on rates of translation has not been described but it does appear to directly interact with the elongation factor-1α (Denisenko and Bomsztyk, unpublished observations), so it may also have a role in translation. Because K protein also binds single- and double-stranded DNA it could also act in the same linker fashion in processes involving DNA. Depending on the circumstances, K protein could assemble on DNA either transcriptional repressors or activators, accounting for the observations that K protein can either increase or decrease rates of transcription [25].

The ability to engage multiple proteins and nucleic acids is not unique to K protein, since, for example, Sam68 which also contains KH domains, binds RNA and interacts with a number of different protein factors involved in signal transduction [26,27]. Sam68 binds to RNA sequences different from those to which K protein binds, and Sam68 engages some protein factors that K protein is not known to bind. As in the case of K protein [9,11], it has been suggested that Sam68 acts as a docking platform or scaffold to facilitate molecular interactions [28]. The class of proteins that act as nucleic acid-regulated docking platforms is likely to include other factors related to K protein and Sam68, each with unique RNA-, DNA- and protein-binding specificities.

**Acknowledgements:** We thank Dr. Carol Sibley for her comments on the manuscript. The work in our laboratory is supported by grants from the NIH, the Northwest Kidney Foundation, and the NATO Collaborative Research Grant.

#### References

- [1] Matunis, M.J., Michael, W.M. and Dreyfuss, G. (1992) *Mol. Cell. Biol.* 12, 164–171.

- [2] Dejgaard, K., Leffers, H., Rasmussen, H.H., Madsen, P., Kruse, T.A., Gesser, B., Nilesen, H. and Celis, J.E. (1994) *J. Mol. Biol.* 236, 33–48.
- [3] Ostrowski, J. and Bomsztyk, K. (1993) *Nucleic Acids Res.* 21, 1045–1046.
- [4] Siomi, H., Matunis, M.J., Michael, W.M. and Dreyfuss, G. (1993) *Nucleic Acids Res.* 21, 1193–1198.
- [5] Gibson, T.J., Thompson, J.D. and Heringa, J. (1993) *FEBS Lett.* 324, 361–366.
- [6] Siomi, H., Choi, M., Siomi, M.C., Nussbaum, R.L. and Dreyfuss, G. (1994) *Cell* 77, 33–39.
- [7] Siomi, H., Siomi, M.C., Nussbaum, R.L. and Dreyfuss, G. (1993) *Cell* 74, 291–298.
- [8] Ostrowski, J., Van Seuningen, I., Seger, R., Rouch, C.T., Sleath, P.R., McMullen, B.A. and Bomsztyk, K. (1994) *J. Biol. Chem.* 269, 17626–17634.
- [9] Van Seuningen, I., Ostrowski, J., Bustelo, X., Sleath, P. and Bomsztyk, K. (1995) *J. Biol. Chem.* 270, 26976–26985.
- [10] Tomonaga, T. and Levens, D. (1995) *J. Biol. Chem.* 270, 4875–4881.
- [11] Denisenko, O.N., O'Neill, B., Ostrowski, J., Van Seuningen, I. and Bomsztyk, K. (1996) *J. Biol. Chem.* 271, 27701–27706.
- [12] Van Seuningen, I., Ostrowski, J. and Bomsztyk, K. (1995) *Biochemistry* 34, 5644–5650.
- [13] Tomonaga, T. and Levens, D. (1996) *Proc. Natl. Acad. Sci. USA* 93, 5830–5835.
- [14] Gaillard, C. and Strauss, F. (1990) *J. Mol. Biol.* 215, 245–255.
- [15] Ito, K., Sato, K. and Endo, H. (1994) *Nucleic Acids Res.* 22, 53–58.
- [16] Takimoto, M., Tomonaga, T., Matunis, M., Avigan, M., Krutzsch, H., Dreyfuss, G. and Levens, D. (1993) *J. Biol. Chem.* 268, 18249–18258.
- [17] Gaillard, C., Weber, M. and Strauss, F. (1988) *Mol. Cell. Biol.* 62, 2380–2385.
- [18] Ostrowski, J., Sims, J.E., Sibley, C.H., Valentine, M.A., Dower, S.K., Meier, K.E. and Bomsztyk, K. (1991) *J. Biol. Chem.* 266, 12722–12733.
- [19] Bustelo, X.R., Suen, K.-I., Michael, W.M., Dreyfuss, G. and Barbacid, M. (1995) *Mol. Cell. Biol.* 15, 1324–1332.
- [20] Tarakhovsky, A., Turner, M., Schaal, S., Mee, P.J., Duddy, L.P., Rajewski, K. and Tybulewicz, V.L.J. (1995) *Nature* 374, 467–470.
- [21] Zhang, R., Alt, F.W., Davidson, L., Orkin, S.H. and Swat, W. (1995) *Nature* 374, 470–473.
- [22] Michelotti, E.F., Michelotti, G.A., Aronsohn, A.I. and Levens, D. (1996) *Mol. Cell. Biol.* 16, 2350–2360.
- [23] Witzgall, R., O'Leary, E., Gessner, R., Ouellette, A.J. and Bonventre, J.V. (1993) *Mol. Cell. Biol.* 13, 1933–1942.
- [24] Morris, J.F., Rauscher III, F.J., Davis, B., Klemsz, M., Xu, D., Tenen, D. and Hromas, R. (1995) *Blood* 86, 3640–3647.
- [25] Lee, M.-H., Mori, S. and Raychaudhuri, P. (1996) *J. Biol. Chem.* 271, 3420–3427.
- [26] Lock, P., Fumagalli, S., Polakis, P., McCormick, F. and Courtneidge, S.A. (1996) *Cell* 84, 23–24.
- [27] Taylor, S.J., Anafi, M., Pawson, T. and Shalloway, D. (1995) *J. Biol. Chem.* 270, 10120–10124.
- [28] Richards, S., Yu, D., Blumer, K.J., Hauslanden, D., Olszowy, M.W., Cennelly, P.A. and Shaw, A.S. (1995) *Mol. Cell. Biol.* 15, 186–197.