

# Extracellular $\text{Ba}^{2+}$ and voltage interact to gate $\text{Ca}^{2+}$ channels at the plasma membrane of stomatal guard cells

David W.A. Hamilton, Adrian Hills<sup>1</sup>, Michael R. Blatt<sup>1,\*</sup>

Laboratory of Plant Physiology and Biophysics, Imperial College at Wye, Wye, Kent TN25 5AH, UK

Received 29 December 2000; accepted 23 January 2001

First published online 6 February 2001

Edited by Richard Cogdell

**Abstract**  $\text{Ca}^{2+}$  channels at the plasma membrane of stomatal guard cells contribute to increases in cytosolic free  $[\text{Ca}^{2+}]_i$  ( $[\text{Ca}^{2+}]_i$ ) that regulate  $\text{K}^+$  and  $\text{Cl}^-$  channels for stomatal closure in higher-plant leaves. Under voltage clamp, the initial rate of increase in  $[\text{Ca}^{2+}]_i$  in guard cells is sensitive to the extracellular divalent concentration, suggesting a close interaction between the permeant ion and channel gating. To test this idea, we recorded single-channel currents across the *Vicia* guard cell plasma membrane using  $\text{Ba}^{2+}$  as a charge carrying ion. Unlike other  $\text{Ca}^{2+}$  channels characterised to date, these channels activate at hyperpolarising voltages. We found that the open probability ( $P_o$ ) increased strongly with external  $\text{Ba}^{2+}$  concentration, consistent with a 4-fold cooperative action of  $\text{Ba}^{2+}$  in which its binding promoted channel opening in the steady state. Dwell time analyses indicated the presence of a single open state and at least three closed states of the channel, and showed that both hyperpolarising voltage and external  $\text{Ba}^{2+}$  concentration prolonged channel residence in the open state. Remarkably, increasing  $\text{Ba}^{2+}$  concentration also enhanced the sensitivity of the open channel to membrane voltage. We propose that  $\text{Ba}^{2+}$  binds at external sites distinct from the permeation pathway and that divalent binding directly influences the voltage gate. © 2001 Federation of European Biochemical Societies. Published by Elsevier Science B.V. All rights reserved.

**Key words:**  $\text{Ca}^{2+}$  channel, hyperpolarisation-activated; Extracellular  $\text{Ba}^{2+}$ ; Gating, ligand-mediated; Cytosolic free  $[\text{Ca}^{2+}]_i$ ; *Vicia*

## 1. Introduction

Changes in cytosolic free  $\text{Ca}^{2+}$  concentration ( $[\text{Ca}^{2+}]_i$ ) are ubiquitous as components of second messenger cascades in plants and have been associated, among others, with mechanical and thermal disturbances [1,2], pathogen attack [3], nodulation [4] and hormonal signal transduction [5–7]. In many cases, the mechanisms for these  $[\text{Ca}^{2+}]_i$  changes are poorly understood, and both  $\text{Ca}^{2+}$  entry and intracellular release have been proposed to play central roles. It seems likely that the early events include activation of  $\text{Ca}^{2+}$  channels at the plasma membrane and  $\text{Ca}^{2+}$  influx that triggers  $\text{Ca}^{2+}$  re-

lease from intracellular stores, much as has been described for  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release (CICR) in animal cells [8,9].

Calcium channels that might fill such a role – exhibiting the canonical, depolarisation-mediated activation of their animal counterparts – have been identified at the plasma membrane of carrot suspension culture cells [10] and wheat roots [11], although in neither case has a physiological role been established. Intriguingly, of the few instances in which plasma membrane  $\text{Ca}^{2+}$  channels are known to be associated with increases in  $[\text{Ca}^{2+}]_i$  in plants, channel activity is potentiated by membrane hyperpolarisation [12–14]. These characteristics indicate gating properties atypical of  $\text{Ca}^{2+}$  channels previously identified in plants or animals.

We previously identified a 12 pS  $\text{Ca}^{2+}$  channel at the plasma membrane of stomatal guard cells [14], following on evidence that membrane hyperpolarisation evokes increases in  $[\text{Ca}^{2+}]_i$  and contributes to  $[\text{Ca}^{2+}]_i$  transients associated with the plant water-stress hormone abscisic acid and related signal cascades [7,15]. These changes in  $[\text{Ca}^{2+}]_i$  depend on  $\text{Ca}^{2+}$  release from intracellular stores [16–19], as well as on  $\text{Ca}^{2+}$  entry across the plasma membrane [15,20]. One intriguing feature of the  $[\text{Ca}^{2+}]_i$  rise in guard cells is its linear dependence on external  $\text{Ca}^{2+}$  concentration [15], including concentrations above those that saturate the open channel conductance of the  $\text{Ca}^{2+}$  channel [14]. This behaviour implies an additional action of the divalent cation on gating. We have examined the gating properties of the guard cell  $\text{Ca}^{2+}$  channel, using  $\text{Ba}^{2+}$  as the permeant cation to avoid  $\text{Ca}^{2+}$  entry and the effects of  $\text{Ca}^{2+}$  on the inner surface of the membrane that lead to channel inactivation [14]. Analysis of single-channel kinetics as a function of  $\text{Ba}^{2+}$  concentration inside and outside the membrane indicates the presence of binding sites associated with the outer surface of the channel that interact cooperatively to stabilise the open state of the channel and enhance its sensitivity to membrane voltage.

## 2. Materials and methods

### 2.1. Plant material

Epidermal strips of *Vicia faba* L., cv. Bunyard Exhibition were obtained and protoplasts prepared as described [21,22]. All operations were carried out on a Zeiss Axiovert microscope with 40×LWD Nomarski D.I.C. optics (Zeiss, Oberkochen, Germany) at 20–22°C. Solution was added ( $\approx 20$  chamber volumes/min) by gravity feed and removed by aspiration.

### 2.2. Electrophysiology

Pipettes were pulled using a Narashige PP-81 puller (Narashige, Tokyo) modified for 3-stage pulls (input resistances, 30–50 M $\Omega$ ) to reduce the number of channels under a patch. Pipettes were coated

\*Corresponding author. Fax: (44)-141-330 4447.  
E-mail: m.blatt@bio.gla.ac.uk

<sup>1</sup> Present address: Laboratory of Plant Physiology and Biophysics, Institute of Biomedical and Life Sciences, University of Glasgow, Hillhead, Glasgow G12 8QQ, UK.

with Sigmacote (Sigma, Poole, UK) to reduce capacitance. Connections to amplifier and bath were via a 0.1 M KCl:Ag–AgCl liquid junctions, and junction potentials were taken into account [23]. Single-channel currents were recorded using an Axopatch 200B patch amplifier (Axon Instruments, Redwood City) after filtering at 5 kHz and sampled at 44 kHz for analysis. Data were filtered at 1 kHz (Kemo, Beckenham, UK) off line and analysed using N-Pro (WyeScience, Kent) and P/V clamp v. 6 (CED, Cambridge) software. Voltages quoted are referenced to the physiological orientation of the membrane, the voltage on the cytosolic side relative to the extracellular side.

### 2.3. Numerical analysis

Channel amplitudes were calculated from point-amplitude histograms estimated from open events  $\geq 5$  ms duration beyond closed levels determined from periods of no channel activity [24]. Channel numbers were estimated from the maximum number of concurrent openings and from binomial distributions of open events [25]. Channel openings were taken as transitions above a threshold of 60% of the single-channel amplitude and  $P_o$  estimated from the open-time fraction corrected for the number of channels. Channel lifetime distributions were determined from dwell-time histograms of open and closed events, and were restricted to patches showing evidence of only a single channel. For analyses of closed lifetimes, the data were limited to the intraburst periods of channel activity and were generally defined by preceding and following closed periods at least 10-fold longer than those within the burst. Open and closed dwell-time histograms were subjected to non-linear, least-squares exponential fittings [24,26] to determine the number of identifiable states and obtain estimates of relaxation constants in each case. Where appropriate results are given as means  $\pm$  S.E.M. of ( $n$ ) independent experiments.

### 2.4. Chemicals and solutions

We use the terms inside and outside with reference to the physiological sidedness of the membrane. Protoplasts were normally bathed in 2, 10 and 30 mM  $\text{Ba}^{2+}$ –HEPES, pH 7.5 (HEPES buffer titrated to its  $\text{pK}_a$  with  $\text{Ba}(\text{OH})_2$  adjusted to 200 mOsm with sorbitol, and pipettes were filled with similar solutions adjusted to 240 mOsm with sorbitol.  $(\text{Mg}^{2+})_2\text{ATP}$  at 1 mM was included in solutions on the inside of the membrane. Buffers and salts were from Sigma Chemicals (Poole, UK).

## 3. Results and discussion

### 3.1. Increasing $[\text{Ba}^{2+}]_o$ strongly promotes channel opening events

A previous study of the guard cell  $\text{Ca}^{2+}$  channel showed that its steady state open probability ( $P_o$ ) was strongly dependent on membrane hyperpolarisation. However, we found that the extent to which hyperpolarisation promoted channel events was also sensitive to the concentration of divalent available to permeate the channel. Fig. 1A shows single-channel records from one membrane patch in the outside-out configuration with  $\text{Ba}^{2+}$  as the permeant ion, and Fig. 1B summarises the results from this, and five additional, independent experiments. In general, increasing the  $\text{Ba}^{2+}$  concentration between 2 and 30 mM resulted in a pronounced rise in the number of channel openings which, on average, gave roughly a 50-fold increase in  $P_o$  at a common clamp voltage of  $-100$  mV. The effect of  $\text{Ba}^{2+}$  also showed a definite sidedness. Similar measurements carried out in the inside-out configuration, varying the the  $\text{Ba}^{2+}$  concentration on the inside of the membrane with 10 mM and 30 mM  $\text{Ba}^{2+}$  outside, showed no comparable change in  $P_o$  [14]. These effects are also clearly different from the effect of  $\text{Ca}^{2+}$  inside, which led to a precipitous drop in channel openings as  $\text{Ca}^{2+}$  concentration rose to micromolar values [14].

Technical difficulties precluded increasing the  $\text{Ba}^{2+}$  on either side of the membrane to obtain estimates of  $P_o$  at con-




Fig. 1. Increasing  $\text{Ba}^{2+}$  concentration outside favours  $\text{Ca}^{2+}$  channel opening in the steady state. A: Single-channel records from one outside-out patch clamped at  $-100$  mV with 30 mM  $\text{Ba}^{2+}$  inside and 2, 10 and 30 mM  $\text{Ba}^{2+}$  outside. Note the bursting characteristics of channel openings. Scale: 1 pA and 100 ms. Point-amplitude histograms (right) for 20 s recordings of the corresponding traces (see Section 2). B: Open probability ( $P_o$ ) as a function of  $\text{Ba}^{2+}$  concentration at  $-100$  mV from six independent experiments. Data fitted to a Hill function with  $P_o$  free (solid line), constrained to a maximum of 1 with  $n$  constrained to 5 (short-dashed line) for comparison. Best fittings gave  $n = 3.6 \pm 0.2$  irrespective of the maximum  $P_o$ . Inset: Best fitting of the same data, joint with measurements at  $-80$  and  $-110$  mV, yielded  $n = 3.7 \pm 0.1$ , an apparent  $K_{1/2}$  of 35 mM at all three voltages and  $P_o$  values of 0.07 (2 mM), 0.38 (10 mM) and 0.97 (30 mM  $\text{Ba}^{2+}$ ).

centrations significantly beyond 30 mM. Nonetheless, these data confirmed a sensitivity to  $\text{Ba}^{2+}$  outside, consistent with a high degree of cooperativity in promoting channel activity. The curve in Fig. 1B shows the results of fitting  $P_o$  to the Hill equation [27]. We found that the fitted value for the Hill coefficient,  $n$ , was well defined and largely insensitive to constraints on the maximum  $P_o$ . Allowing the maximum  $P_o$  to vary freely during the fitting yielded a Hill coefficient of 3.7 and, even with  $P_o$  constrained to a maximum of 1 gave a value only marginally smaller (see Fig. 1B and legend). Constraining  $n$  to values above 4, by contrast, failed to give visually acceptable matches to the data (Fig. 1B). Joint fittings incorporating  $P_o$  values gathered at  $-80$  mV and at  $-110$  mV yielded similar, and statistically more satisfactory results with a Hill coefficient of 3.6, both when values for maximum  $P_o$  and when  $K_{1/2}$  were held in common between data at each voltage (Fig. 1B, inset). Thus, while the data are insufficient to define either of these parameters without points at higher

values for  $[\text{Ba}^{2+}]_o$ , they imply the cooperative binding of four  $\text{Ba}^{2+}$  ions to facilitate opening of the channel.

### 3.2. Voltage and $[\text{Ba}^{2+}]_o$ suppress channel closure

To explore the mechanism of  $\text{Ba}^{2+}$  action and relationship to the voltage-evoked increase in  $P_o$ , we examined the open lifetime distributions of the  $\text{Ca}^{2+}$  channel using recordings in which only a single open level was clearly evident. Recordings from outside-out patches were selected for analysis that included measurements of 100 s duration at each of three  $[\text{Ba}^{2+}]_o$ , 2, 10 and 30 mM and over a range of membrane voltages. Representative lifetime distributions for one patch at selected voltages are shown in Fig. 2A and the results of




Fig. 2. Mean open lifetime of the  $\text{Ca}^{2+}$  channel, and its sensitivity to voltage, depends on external  $\text{Ba}^{2+}$  concentration. A: Open lifetime distributions at voltages between  $-80$  and  $-130$  mV for one outside-out patch with 30 mM  $\text{Ba}^{2+}$  inside and 10 mM  $\text{Ba}^{2+}$  outside. Data were well fitted in every case to a single-exponential function. B: Mean open lifetimes as functions of voltage with 2 (●), 10 (○) and 30 mM  $\text{Ba}^{2+}$  (▼) outside. Data are means  $\pm$  S.E.M., from six independent experiments and show a roughly linear dependence on voltage. Inset: Slope of voltage dependence as a function of  $\text{Ba}^{2+}$  concentration.




Fig. 3. Mean closed lifetime during bursts of  $\text{Ca}^{2+}$  channel activity depends on external  $\text{Ba}^{2+}$  concentration but not on voltage. A: Closed lifetime distributions at voltages between  $-80$  and  $-130$  mV for the same outside-out patch with 30 mM  $\text{Ba}^{2+}$  inside and 10 mM  $\text{Ba}^{2+}$  outside as in Fig. 2. Data were well fitted in every case only to a sum of two exponential components. B: Mean closed lifetimes as functions of voltage (components: fast, slow) with 2 (●, ○), 10 (▲, △) and 30 mM  $\text{Ba}^{2+}$  (▼, ▽) outside. Data are means  $\pm$  S.E.M. from six independent experiments. Inset: Mean closed lifetime dependence of the fast (○) and slow (●) components as a function of  $\text{Ba}^{2+}$  concentration. Data were averaged over all voltages at each concentration.

analyses from six independent experiments, including these data, are summarised in Fig. 2B. In general, open lifetime distributions were well described by a single-exponential function for which the mean open lifetime,  $\tau$ , increased with negative membrane voltage and  $[\text{Ba}^{2+}]_o$ . The analysis also showed that the sensitivity of  $\tau$  to voltage increased with  $[\text{Ba}^{2+}]_o$ . This voltage sensitivity exhibited roughly a 2-fold rise over the order of magnitude increase in  $[\text{Ba}^{2+}]_o$  of these experiments (Fig. 2B, inset). Thus, the  $\text{Ca}^{2+}$  channel could be characterised by the presence of a single open state. Furthermore, increasing negative membrane voltage prolonged the lifetime of the channel once in the open state, and increasing  $[\text{Ba}^{2+}]_o$  potentiated this effect of voltage, as if  $\text{Ba}^{2+}$  itself contributed in part to the voltage sensor.

Analysis of the closed lifetimes of the  $\text{Ca}^{2+}$  channel pointed

to a more complex pattern of non-conducting states. Visual inspection showed a pronounced bursting characteristic to the channel (cf. Fig. 1A) with frequent periods, often of seconds, without opening events. Insufficient data were obtained to carry out a quantitative analysis of this bursting behaviour. Nevertheless, its occurrence indicates at least one long-lived closed state of the  $\text{Ca}^{2+}$  channel. Within the burst periods, the distribution of closed lifetimes showed little evidence of voltage dependence. Fig. 3A shows the closed lifetime distributions from the same experiment as in Fig. 2A and these are included in the summary in Fig. 3B. Closed lifetime distributions required the inclusion of a second exponential component for fitting. We found little evidence of a sensitivity to clamp voltage for either of the exponential components, but increasing  $[\text{Ba}^{2+}]_o$  resulted in a significant decrease in the mean closed lifetime of the slower component (Fig. 3B). The lifetime analyses, in this case, must be treated with some caution. Although we took care to select records that appeared to show only a single channel, the very low values for  $P_o$  made it possible that more than one channel was present even in these circumstances (see also Hamilton, et al. [14]). With two or more channels in a patch, it becomes impossible to know which opens at the end of a closed period [24]. One consequence is a skewing the analysis towards shorter apparent closed lifetimes which, in this case, may mask more profound effects of experimental manipulations. This caveat aside, the kinetic analysis and characteristics of bursting indicates the presence of at least three closed states.

### 3.3. Interpreting the interaction of external $\text{Ba}^{2+}$ with voltage in gating

From Figs. 2 and 3 it is evident that a minimal model for  $\text{Ca}^{2+}$  channel gating requires a single open state and three separate closed states. At present we have few clues as to whether the open state communicates directly with only one, or with more than one of the closed states. However, the data do point to a strong effect both of voltage and  $[\text{Ba}^{2+}]_o$  on the open channel lifetime (Fig. 2), the slower of the two closed lifetimes (Fig. 3) and, as a consequence on  $P_o$  in the steady state (Fig. 1; see also [14]). Furthermore, because  $P_o$  is profoundly dependent on membrane hyperpolarisation [14], we anticipate that the remaining rates of transition between states may also subsume a component of this voltage sensitivity. Overall, the effect of voltage on  $P_o$  on the guard cell  $\text{Ca}^{2+}$  channel is comparable to its influence on  $\text{Ca}^{2+}$  channels now identified from a variety of cell types, including leaf mesophyll [13] and root hairs [28]. Gating processes for plant  $\text{Ca}^{2+}$  channels, in general, have not been well defined. For the guard cell  $\text{Ca}^{2+}$  channel, our results now link voltage to an action on the channel in the open state. In essence, hyperpolarising the membrane stabilises the channel once open, effectively 'locking' the channel in this molecular configuration.

The most striking feature of this  $\text{Ca}^{2+}$  channel, however, is the effect of external  $\text{Ba}^{2+}$  concentration and its interaction with the action of membrane voltage. Not only did increasing  $[\text{Ba}^{2+}]_o$  act to stabilise the open state of the channel, it also enhanced the sensitivity of the open state to membrane hyperpolarisations (Fig. 2B). We are not aware of such a synergy between the chemical and electrical parameters controlling gating in  $\text{Ca}^{2+}$  channels previously characterised either in plants or animals. For neuronal  $\text{Ca}^{2+}$  channels, changes of  $P_o$  with external  $\text{Ca}^{2+}$  are known, but these effects have

been ascribed to surface-charge masking and generally saturate at concentrations near 2 mM [29,30]. In guard cells and yeast, external alkali cations affect voltage-dependent gating of  $\text{K}^+$  channels [31–33], an effect which is mimicked by  $\text{Ba}^{2+}$  in the yeast YKC1  $\text{K}^+$  channel [32]. However, the actions of external cations in these instances are mediated through voltage-dependent block, albeit with cation binding to allosteric sites distinct from the channel pore. Furthermore, both in these  $\text{K}^+$  channels and in the animal  $\text{Ca}^{2+}$  channels the external cations displace the voltage dependence of channel gating, seemingly altering its voltage 'set-point', but without affecting the underlying sensitivity to voltage. In both respects, therefore, the guard cell  $\text{Ca}^{2+}$  channel appears unique.

The fact that increasing  $[\text{Ba}^{2+}]_o$  enhances  $P_o$  and the open channel dwell time as a function of voltage does suggest that the external divalent is an integral component of the voltage gate. Plausibly,  $\text{Ba}^{2+}$  binding to the outer surface of the  $\text{Ca}^{2+}$  channel may influence local energy landscape traversed by the voltage gate. For example, it could become an integral part of the voltage sensor itself, although other physical mechanisms are also possible. Our data do indicate an external location for  $\text{Ba}^{2+}$  binding that is separate from the permeation pathway, since changing internal  $\text{Ba}^{2+}$  concentration did not have an equivalent effect on  $P_o$  (compare also [33]). Therefore, we suggest the presence of four binding sites to account for the high apparent cooperativity of  $[\text{Ba}^{2+}]_o$  action on  $P_o$  (Fig. 1), recognising the 4-fold symmetry in the assembly of subunits of many voltage-sensitive cation channels known to date [34–37], and the implicit distribution of binding sites, one per subunit assembly. Clearly, proof of this idea must now await genetic identification of the  $\text{Ca}^{2+}$  channel.

**Acknowledgements:** We are grateful to Gerhard Thiel, Anna Moroni and Alexander Grabov for comments on the manuscript. This work was possible with the aid of Grants from the BBSRC Grants P09640, C10234, P09561 and British Council Grant PRO897. D.H. was a Sainsbury Ph.D. Student.

### References

- [1] Knight, H., Trewavas, A.J. and Knight, M.R. (1996) *Plant Cell* 8, 489–503.
- [2] Taylor, L.P. and Hepler, P.K. (1997) *Annu. Rev. Plant Physiol. Mol. Biol.* 48, 461–491.
- [3] Hammond-Kosack, K.E. and Jones, J.D.G. (1996) *Plant Cell* 8, 1773–1791.
- [4] Ehrhardt, D.W., Wais, R. and Long, S.R. (1996) *Cell* 85, 673–681.
- [5] Ritchie, S. and Gilroy, S. (1998) *New Phytol.* 140, 363–383.
- [6] Bush, D.S. (1993) *Plant Physiol.* 103, 7–13.
- [7] Blatt, M.R. (2000) *Annu. Rev. Cell Dev. Biol.* 16, 221–241.
- [8] Berridge, M.J. (1998) *Neuron* 21, 13–26.
- [9] Bootman, M.D., Berridge, M.J. and Lipp, P. (1997) *Cell* 91, 367–373.
- [10] Thuleau, P., Ward, J.M., Ranjeva, R. and Schroeder, J.I. (1994) *EMBO J.* 13, 2970–2975.
- [11] Pineres, M. and Tester, M. (1995) *Planta* 195, 478–488.
- [12] Zimmermann, S., Nurnberger, T., Frachisse, J.M., Wirtz, W., Guern, J., Hedrich, R. and Scheel, D. (1997) *Proc. Natl. Acad. Sci. USA* 94, 2751–2755.
- [13] Gelli, A. and Blumwald, E. (1997) *J. Membr. Biol.* 155, 35–45.
- [14] Hamilton, D.W.A., Hills, A., Kohler, B. and Blatt, M.R. (2000) *Proc. Natl. Acad. Sci. USA* 97, 4967–4972.
- [15] Grabov, A. and Blatt, M.R. (1998) *Proc. Natl. Acad. Sci. USA* 95, 4778–4783.
- [16] Grabov, A. and Blatt, M.R. (1999) *Plant Physiol.* 119, 277–287.
- [17] Leckie, C.P., McAinsh, M.R., Allen, G.J., Sanders, D. and

- Hetherington, A.M. (1998) *Proc. Natl. Acad. Sci. USA* 95, 15837–15842.
- [18] Gilroy, S., Fricker, M.D., Read, N.D. and Trewavas, A.J. (1991) *Plant Cell* 3, 333–344.
- [19] Staxen, I., Pical, C., Montgomery, L.T., Gray, J.E., Hetherington, A.M. and McAinsh, M.R. (1999) *Proc. Natl. Acad. Sci. USA* 96, 1779–1784.
- [20] Webb, A.A.R., McAinsh, M.R., Mansfield, T.A. and Hetherington, A.M. (1996) *Plant J.* 9, 297–304.
- [21] Blatt, M.R. and Armstrong, F. (1993) *Planta* 191, 330–341.
- [22] Elzenga, J.T.M., Keller, C.P. and Van Volkenburgh, E. (1991) *Plant Physiol.* 97, 1573–1575.
- [23] Barry, P.H. and Lynch, J.W. (1991) *J. Membr. Biol.* 121, 101–117.
- [24] Colquhoun, D. and Sigworth, F.J. (1995) in: *Practical Analysis of Single Channel Records* (Sakmann, B. and Neher, E., Eds.), 2(18), pp. 397–482.
- [25] Horn, R. (1991) *Biophys. J.* 60, 433–439.
- [26] Marquardt, D. (1963) *J. Soc. Ind. Appl. Math.* 11, 431–441.
- [27] Hill, A.V. (1910) *J. Physiol.* 40, 4–7.
- [28] Kiegle, E., Gilliam, M., Haseloff, J. and Tester, M. (2000) *Plant J.* 21, 225–229.
- [29] Zamponi, G.W. and Snutch, T.P. (1996) *Pflug. Arch. Eur. J. Physiol.* 431, 470–472.
- [30] Hille, B. (1992) *Ionic Channels of Excitable Membranes*, pp. 1–607, Sinauer, Sunderland, MA.
- [31] Blatt, M.R. and Gradmann, D. (1997) *J. Membr. Biol.* 158, 241–256.
- [32] Vergani, P., Miosga, T., Jarvis, S.M. and Blatt, M.R. (1997) *FEBS Lett.* 405, 337–344.
- [33] Vergani, P., Hamilton, D., Jarvis, S. and Blatt, M.R. (1998) *EMBO J.* 17, 7190–7198.
- [34] Catterall, W.A. (1995) *Annu. Rev. Biochem.* 64, 493–531.
- [35] MacKinnon, R. (1995) *Neuron* 14, 889–892.
- [36] Doyle, D.A., Cabral, J.M., Pfuetzner, R.A., Kuo, A.L., Gulbis, J.M., Cohen, S.L., Chait, B.T. and MacKinnon, R. (1998) *Science* 280, 69–77.
- [37] MacKinnon, R., Cohen, S.L., Kuo, A.L., Lee, A. and Chait, B.T. (1998) *Science* 280, 106–109.