

Synthetic and structural studies on *Pyrularia pubera* thionin: a single-residue mutation enhances activity against Gram-negative bacteria

Miquel Vila-Perelló^a, Andrea Sánchez-Vallet^b, Francisco García-Olmedo^b, Antonio Molina^b, David Andreu^{a,*}

^aDepartment of Experimental and Health Sciences, Pompeu Fabra University, Dr. Aiguader 80, E-08003 Barcelona, Spain

^bLaboratory of Biochemistry, Department of Biotechnology, UPM-ETSIA, Avda. Complutense, E-28040 Madrid, Spain

Received 11 November 2002; revised 10 January 2003; accepted 14 January 2003

First published online 23 January 2003

Edited by Gunnar von Heijne

Abstract The thionin from *Pyrularia pubera* (Pp-TH), a 47-residue peptide with four internal disulfide bonds, was efficiently produced by chemical synthesis. Its antimicrobial activity in vitro against several representative pathogens ($EC_{50} = 0.3\text{--}3.0\ \mu\text{M}$) was identical to that of natural Pp-TH. This peptide has a unique Asp³² instead of the consensus Arg found in other thionins of the same family. In order to evaluate the effect of this mutation, the Arg³² analogue (Pp-TH(D32R)) was also synthesized and showed a significant increase in antibiotic activity against several Gram-negative bacteria, whereas it retained the same activity against other pathogens. The overall structure of Pp-TH(D32R) was maintained, though a slight decrease in the helical content of the peptide was observed.

© 2003 Published by Elsevier Science B.V. on behalf of the Federation of European Biochemical Societies.

Key words: Thionin; Antimicrobial peptide; Oxidative folding; Plant

1. Introduction

All organisms, from plants and insects to humans, produce gene-encoded cationic antimicrobial peptides [1–3] as constituents of their innate immune systems. In recent years these peptides have been proposed as a potential solution to the widespread appearance of resistance to classical antibiotics in clinically relevant bacteria [4–6]. Such expectations are based on the fact that these peptides exert their antimicrobial action through fairly simple mechanisms of membrane disruption [7–9] which are unlikely to promote the appearance of resistance [3].

In the course of our studies on antimicrobial peptides we have focused our attention on the thionins, the first family of plant peptides [10,11] for which activity against plant pathogens was demonstrated in vitro [12]. Thionins are basic pep-

tides (minimum net charge of +2), usually 45–47 amino acids in length, with a high rate of sequence homology [10,11]. They have been classified into five structural types (I–V) based on their amino acid sequences and disulfide bridge patterns. The secondary structure of thionins is well conserved and consists of one β -sheet and two antiparallel α -helices, with hydrophobic and hydrophilic residues segregated at the outer and inner helix surfaces, respectively. Such an amphipathic distribution gives thionins their ability to disrupt microbial membranes and phospholipid liposomes [13]. Despite sequence differences, the 3D structure of thionins of types I–IV is almost identical, resembling a capital gamma (Γ) [14].

The thionin from *Pyrularia pubera* (Pp-TH), a cytotoxic and hemolytic type II thionin [15], is remarkable for the presence of an Asp residue at position 32 instead of the consensus Arg found in most family I and II thionins with the same disulfide pattern (Fig. 1) [10,11]. In order to evaluate the effect of Asp³², and the ensuing loss of cationic character, on the activity of this peptide, we decided to synthesize both the native sequence and its Arg³² (D32R) analogue, with a view to expanding the spectrum of antimicrobial activity of this thionin.

We report here the chemical synthesis of Pp-TH and its D32R analogue, their study by circular dichroism (CD) and the in vitro antimicrobial activities against bacterial and fungal plant pathogens. The antibiotic activities of the synthetic and natural Pp-TH were found to be identical against the pathogens tested. In contrast, the D32R mutation selectively increased the activity against the Gram-negative species analyzed, whereas it did not affect the activity against other pathogens. The D32R substitution has only minor effects in the overall structure of the thionin.

2. Materials and methods

2.1. Peptide synthesis

The 47-amino acid sequence of Pp-TH (KSCCRNTWARN-CYNVCRPLPGTISREICAKKCDCKIISGTTCPSPDYPK) and its D32R analogue were assembled on a 9-fluorenylmethoxycarbonyl (Fmoc)-Lys(*tert*-butoxycarbonyl (Boc))-polyethyleneglycol polystyrene resin (0.18 mmol/g, Applied Biosystems) by Fmoc solid-phase synthesis protocols using an Applied Biosystems 433A instrument. Side chain protection was as follows: Lys(Boc), Ser(*tert*-butyl ('Bu)), Arg(2,2,5,7,8-pentamethylchroman-6-sulfonyl), Asn(trityl (Trt)), Trp(Boc), Tyr('Bu), Thr('Bu), Glu(*tert*-butyl ester), Asp(*tert*-butyl ester) and Cys(Trt). Couplings (10 equivalent of Fmoc-amino acid) were mediated by 2-(1-*H*-benzotriazol-1-yl)-1,1,3,3-tetramethyl-

*Corresponding author. Fax: (34)-93-542 28 02.

E-mail address: david.andreu@cexs.upf.es (D. Andreu).

Abbreviations: Boc, *tert*-butoxycarbonyl; 'Bu, *tert*-butyl; Trt, trityl; Fmoc, 9-fluorenylmethoxycarbonyl; DMF, dimethylformamide; HPLC, high performance liquid chromatography; MALDI-TOF MS, matrix-assisted laser desorption ionization time of flight mass spectrometry; TFA, trifluoroacetic acid; GSH, reduced glutathione; GSSG, oxidized glutathione; CD, circular dichroism

uronium tetrafluoroborate (10 equivalent) and diisopropylethylamine (20 equivalent) in dimethylformamide (DMF). After chain assembly was completed, the N^α Fmoc group was removed with 20% piperidine in DMF and the peptide was fully deprotected and cleaved from the resin by treatment with trifluoroacetic acid (TFA)/thioanisole/water/phenol/ethanedithiol (82.5:5:5:5:2.5) for 3.5 h. The peptides were isolated by precipitation with cold *tert*-butyl methyl ether and centrifugation, then taken up in 10% AcOH and lyophilized. The crude octathiol peptides thus obtained were purified by preparative reverse-phase high-performance liquid chromatography (HPLC; Kromasil C8, 2×25 cm column, 10 μm particle size, 100 Å pore size, Akzo Nobel), using a 10–40% linear gradient of 0.1% TFA/acetonitrile into 0.1% TFA/water over 60 min at 25 ml/min. Fractions judged to be homogeneous by analytical HPLC were pooled and their identity was further confirmed by amino acid analysis and matrix-assisted laser desorption ionization time of flight (MALDI-TOF) mass spectra, both of them consistent with the expected compositions.

2.2. Oxidative folding of octathiol precursors

Several oxidative folding conditions (see Section 3 and Table 1) were tested. In the optimal protocol, the purified octathiol was dissolved in 0.1 M NH₄OAc, pH 7.8, at 5 μM concentration, under Ar atmosphere, in the presence of 1 M Gn·HCl and reduced and oxidized glutathione (GSH, GSSG) to give a peptide:GSH:GSSG ratio of 1:100:10. After gentle stirring for 48 h at 25°C, and since the Ellman test [16] of the major HPLC product was negative, the reaction mixture was quenched by TFA addition to pH 1.5–2, then directly loaded onto a preparative reverse-phase HPLC system (see above). After eluting the salts with 0.1% TFA/water, the folded peptide was purified by a linear 10–40% gradient of 0.1% TFA/acetonitrile into 0.1% TFA/water over 60 min. The Pp-TH(D32R) analogue was synthesized and purified by similar methods. The folded synthetic peptide coeluted with an authentic sample of Pp-TH under the above HPLC conditions, as well as in isocratic elution at 20% acetonitrile in 0.1% TFA/water.

2.3. Circular dichroism

CD spectra were collected on a JASCO J-715 spectropolarimeter using a quartz cuvette of 0.1 cm pathlength. The system was purged with N₂ (25 ml/min) and thermostated by means of a water bath. Peptide samples (15 μM; 25 μM for thermal unfolding curves) were prepared in 1 mM citrate, 1 mM phosphate, 1 mM borate buffer. For each spectrum two 190–260-nm scans were accumulated, at 10 nm/min, with 0.2 nm resolution, 2 s response and 1.0 nm band width. Thermal unfolding curves between 5 and 80°C were obtained by measuring molar ellipticity at the 207 nm minimum, using an 0.5°C resolution and a 20°C/h ramp.

2.4. Modeling studies

The sequences of Pp-TH and Pp-TH(D32R) were submitted to the SWISS-MODEL (www.expasy.org/swissmod/SWISS-MODEL.html) automatic server for protein modeling, and the PDB files thus ob-

tained were used to generate the input file for molecular dynamics studies with the package MOLARIS (v. 9.05). Molecular dynamics were run for 500 000 steps of 0.002 ps at 300 K with a 22 Å water grid radius, and 24 Å Langevin grid radius. A pH titration with POLARIS was previously performed in order to fix side chain charges in the folded peptides.

2.5. Antimicrobial activity

Inhibition tests were carried out in sterile microtiter plates by mixing different amounts of the peptide dissolved in 66.7 μl of sterile water with either 33.3 μl of bacterial suspensions (final concentration 10⁴ cfu/ml) in nutrient broth (Oxoid, Basingstoke, UK) or TY medium (*Rhizobium meliloti* [17]), or 33.3 μl of fungal spore suspensions (final concentration 10⁴ spores/ml) in potato dextrose (Difco, Detroit, MI, USA). Growth of microorganisms was recorded 24–48 h after incubation at 28°C, by measuring absorbance at 490 nm in an ELISA plate reader. Spores used in the inhibition experiments were harvested and stored as previously described [18]. The bacterial pathogens *Clavibacter michiganensis* subsp. *sepedonicus*, strain C5, *Xanthomonas campestris* pv. *campestris*, *X. campestris* pv. *translucens* and *R. meliloti*, as well as the fungal pathogen *Botrytis cinerea* (strain 1) were from the ETSIA collection (Madrid, Spain). The fungi *Plectosphaerella cucumerina* and *Fusarium oxysporum* f. sp. *conglutinans* were the gift of Dr. B. Mauch-Mani (University of Fribourg, Switzerland) and Dr. M.I.G. Roncero (University of Cordoba, Spain), respectively. Purified *P. pubera* thionin was the gift of Dr. John D. Bell (Brigham Young University, Provo, UT, USA).

3. Results

3.1. Synthetic thionins

Pp-TH and Pp-TH(D32R) were readily obtained by Fmoc/^tBu solid-phase peptide synthesis protocols [19,20], with all eight Cys residues protected with the Trt group. HPLC purification of the products resulting from TFA treatment gave the fully reduced octathiol precursors in highly pure form, in 15–20% yield (relative to the original resin substitution). MALDI-TOF mass spectrometry (MS) analysis gave molecular masses of 5288.15 and 5330.64 Da for the Pp-TH and Pp-TH(D32R) precursors, respectively, in full agreement with the theoretical values (5288.24 and 5330.34). Different temperature and denaturant conditions (Table 1) were tested for the folding [21,22] of these precursors, the optimal preparative protocol consisting in anaerobic (Ar atmosphere) oxidation of a 5 μM reduced peptide solution in 0.1 M NH₄OAc buffer (pH 7.8) in the presence of GSH (500 μM), GSSG (50 μM) and Gn·HCl (1 M) for 48 h at 25°C. The reactions were




Fig. 1. Amino acid sequence alignment (A) and disulfide bridge pattern (B) of thionins, types II and I. Conserved positions are shaded in black. The Asp residue of native Pp-TH (SwissProt accession number P07504) mutated to Arg in Pp-TH(D32R) is indicated by an arrow. The following sequences (SwissProt accession numbers in parentheses) were used in the alignment algorithm: Hv-THD (P08772), Hv-THB (P09618), Ta-THβ (P01543), Hv-THβ (P21742) and Hv-THα (P01545).




Fig. 2. HPLC analysis of the oxidative folding of Pp-TH after 0 (a), 2 (b) and 48 h reaction (c) on a Nucleosil C18 (0.4×25 cm) column, eluted with a 15–45% linear gradient of acetonitrile/0.1% TFA into 0.1% TFA over 30 min. UV detection at 220 nm was used.

remarkably clean (Fig. 2), with combined yields in the 35–40% range consistently obtained for the folding and subsequent HPLC purification of the folded peptides, which had the expected amino acid compositions and were shown by MALDI-TOF MS to have molecular masses exactly 8.0 Da lower than the precursor peptides: 5280.28 (theory, 5280.17) and 5322.54 (theory 5322.27) for Pp-TH and Pp-TH(D32R), respectively (Fig. 3). Synthetic Pp-TH coeluted with an authentic sample of the natural peptide under several HPLC conditions (Fig. 3).

3.2. Structural studies

The CD spectra of Pp-TH (Fig. 4) in aqueous solution are consistent with a mainly α -helical conformation. The natural and synthetic forms of the peptide gave essentially superimposable signatures, providing evidence for the correct folding of the synthetic material (see Section 4). The spectrum of the D32R analogue had similar features, but minima at 222 and 207 nm were less intense, indicating lower helical content. Helicities were estimated by the Yang equation [23,24] (which, being devised for linear helical peptides, provides only a rough approximation in this case) to be 70% and 43% for Pp-TH and Pp-TH(D32R), respectively. Thermal denaturation experiments showed the helical pattern to be fully preserved between 5 and 80°C (data not shown). Only slight variations in CD curve shapes were observed in pH titration experiments between pH 3 and 7, with the overall structure being maintained throughout the interval. Interestingly, for Pp-TH an increase of helicity with pH was found, while Pp-TH(D32R) exhibited the opposite effect.




Fig. 3. MALDI-TOF mass spectra of Pp-TH and Pp-TH(D32R) (left and right panels, respectively). Spectra of the reduced (dotted line) and oxidized (full line) forms of each peptide are superimposed. Left panel inset: HPLC coelution of equivalent amounts of native and synthetic Pp-TH. HPLC conditions as in Fig. 2.

Assuming these changes to be related to the presence in the former peptide of the Asp³² residue, which in the latter is replaced by Arg, we obtained 3D structures for both peptides by molecular modeling, using the SWISS-MODEL homology server [25]. The modeled structures were generated from 11 available Protein Data Bank templates, with an homology range of 50–70%, and were refined by energy minimization with Gromos 96. The resulting structures were further evaluated by molecular dynamics using the package MOLARIS [26]. Results from the computational study (Fig. 5) were in agreement with the CD observation of decrease in helical content upon Arg-for-Asp replacement.

3.3. Antimicrobial activity

The antibiotic activities of natural and synthetic *P. pubera* thionin [(Pp-TH(nat) and (Pp-TH(synt), respectively], and the synthetic D32R analogue [Pp-TH(D32R)] were evaluated in vitro against bacterial and fungal plant pathogens (Table 2). Pp-TH(synt) and Pp-TH(nat) were similarly active against all the microorganisms tested (EC_{50} = 0.33–3.1 μ M), which further reinforced the previous chemical (HPLC) and structural (CD) evidence for the correct folding of the synthetic thionin. The D32R analogue was significantly more active than Pp-TH against several Gram-negative bacteria (*R. meliloti*, *X. campestris* pv. *translucens* and *X. campestris* pv. *campestris*) but comparable to Pp-TH against the Gram-positive bacterium *C. michiganensis* and the fungal plant pathogens *P. cucumerina*, *F. oxysporum* and *B. cinerea* (Table 2). As previously described [18,27,28], inhibition of microbial growth was abolished by salt (1 mM $CaCl_2$ +50 mM KCl; data not shown).

Table 1
Conditions for oxidative folding of Pp-TH and Pp-TH(D32R) precursors^a

	1	2	3	4	5
Denaturant	1 M HCl-Gn	1 M HCl-Gn	2 M HCl-Gn 1 mM EDTA	2 M HCl-Gn 1 mM EDTA	1 mM EDTA
Temperature	25°C	5°C	25°C	5°C	25°C
Time ^b	48 h	96 h	48 h	72 h	48 h
Extent of reaction ^c	74%	68%	70%	56%	72%

^aReduced octathiol precursor at 5 μ M in 0.1 M NH_4OAc , pH 7.8.

^bTime required for maximum extent of oxidative reaction.

^cEstimated by HPLC peak integration of main oxidation product.




Fig. 4. CD spectra of natural (Pp-TH(nat)) and synthetic Pp-TH (Pp-TH(synt)) and of the D32R analogue (Pp-TH(D32R)). See conditions in text.

4. Discussion

Thionins were the first plant antimicrobial peptides to show activity against plant pathogens *in vitro* and the first eukaryotic peptides for which a defense role was postulated [12]. The present study had a two-fold purpose: (i) to provide useful amounts of *P. pubera* thionin for a complete study of its antimicrobial spectrum, and (ii) to investigate the role of Asp³², an unusual substitution at one of the best conserved positions in the thionins (Fig. 1), and in particular in those of subfamilies I and II, which share an identical disulfide pattern.

The efficient synthesis of both Pp-TH and its D32R analogue described above has been crucial for the success of our approach. The linear precursor, assembled in high purity by stepwise Fmoc solid-phase synthesis, could be cleanly oxidized under highly optimized conditions to provide the folded peptides in reasonable yields. Despite the resistance of the peptide (synthetic and natural) to most proteases even under denaturing conditions, which prevented us from determining its disulfide connectivity by proteolysis and mass spectrometry, we are sure of having achieved the native folding in both synthetic Pp-TH and its D32R analogue, based on several criteria:




Fig. 5. 3D structure of the Arg²⁴–Lys³³ segment of Pp-TH (a) and Pp-TH(D32R) (b) after 1 ns of molecular dynamics simulations with MOLARIS. The side chains of Asp/Arg³² and Lys^{29,30} are displayed.

(i) synthetic Pp-TH coelutes with the natural peptide under several HPLC conditions; (ii) CD spectra of both peptides are superimposable, and (iii) their antimicrobial activities against several representative plant pathogens are indistinguishable.

Interestingly, the Asp to Arg replacement in the D32R analogue significantly enhanced activity against several Gram-negative bacteria, while no relevant effects against Gram-positives or fungi were found. This fact can be plausibly related to the presence of two additional positive charges in the mutant peptide, cationic character being well known to favor antimicrobial activity of peptides [29,30]. The different effect of this mutation on the activity of Pp-TH against Gram-negative and -positive bacteria is probably related to differences in membrane composition. In line with this hypothesis, it has been shown that the permeabilization of model membranes by thionins was dependent on their phospholipid composition [13], and thionin-sensitive mutants of the Gram-negative bacterium *Ralstonia solanacearum* have been described to display alterations in their membrane lipopolysaccharide structure [31].

On the other hand, the mutant D32R shows a surprising decrease in helicity, unpredicted by Chou–Fasman algorithms [32,33], which may be explained (as suggested by molecular modeling, Fig. 5) by the newly created repulsion between Arg³² and Lys²⁹ (*i, i+3* position), compounded by the possible loss of the salt bridge between Asp³² and Lys²⁹, a potentially

Table 2
Inhibition of bacterial and fungal plant pathogens by Pp-TH(nat), Pp-TH(synt) and Pp-TH(D32R)

Pathogen	EC ₅₀ (μM) ^a		
	Pp-TH(nat)	Pp-TH(synt)	Pp-TH(D32R)
Bacteria			
Gram-negative			
<i>R. meliloti</i>	> 20	> 20	3.3 ± 1 ^b
<i>X. campestris</i> pv. <i>campestris</i>	3.1 ± 0.1	2.9 ± 0.1	1.2 ± 0.1 ^b
<i>X. campestris</i> pv. <i>translucens</i>	1.7 ± 0.4	1.4 ± 0.1	0.8 ± 0.03 ^b
Gram-positive			
<i>C. michiganensis</i>	0.48 ± 0.1	0.48 ± 0.03	0.38 ± 0.03
Fungi			
<i>F. oxysporum</i> f. sp. <i>conglutinans</i>	0.43 ± 0.15	0.33 ± 0.06	0.73 ± 0.21
<i>P. cucumerina</i>	0.7 ± 0.17	0.8 ± 0.2	1.25 ± 0.25
<i>B. cinerea</i>	2.2 ± 0.1	nd ^c	2.5 ± 0.3

^aEffective concentration for 50% inhibition (mean of two/three experiments ± S.D.).

^bEC₅₀ values of Pp-TH(D32R), Pp-TH(nat) and Pp-TH(synt) are statistically different (*P* = 0.05, *t*-test).

^cnd, not done.

helix-stabilizing feature of the native form. Despite this slight difference in helicity, both Pp-TH and Pp-TH(D32R) showed remarkable conformational imperviousness to different environments (pH, temperature), as well as resistance to proteolysis, all in all suggesting a very compact folding highly stabilized by the four disulfide bridges. Taken together, these results clearly show that even compact and complex structures such as thionins are susceptible to considerable improvements in their activity by judicious choice of structural modifications.

Acknowledgements: This work was supported by funds from the Generalitat de Catalunya, Spain (Centre de Referència en Biotecnologia, to D.A.) and the Ministerio de Ciencia y Tecnología, Spain (BIO2000-1308, to A.M.). M.V.-P. thanks the Department of Universities and Research of the Generalitat de Catalunya for a predoctoral fellowship. We also acknowledge bioinformatic support by Dr. J. Villà (Grup de Recerca en Bioinformàtica, Universitat Pompeu Fabra) and technical help from G. López (ETSIA, Madrid).

References

- [1] Andreu, D. and Rivas, L. (1998) *Pept. Sci.* 47, 415–433.
- [2] Boman, H.G. (2000) *Immunol. Rev.* 173, 5–16.
- [3] Zasloff, M. (2002) *Nature* 415, 389–395.
- [4] Ganz, T. and Lehrer, R.I. (1999) *Mol. Med. Today* 5, 292–297.
- [5] Cohen, L.M. (1992) *Science* 257, 1050–1054.
- [6] Walsh, C. (2000) *Nature* 406, 775–781.
- [7] Dathe, M., Meyer, J., Beyerman, M., Maul, B., Hoischen, C. and Bienert, M. (2002) *Biochim. Biophys. Acta* 1558, 171–186.
- [8] Ganz, T. and Lehrer, R.I. (1995) *Pharmacol. Ther.* 66, 191–205.
- [9] Coulon, A., Berkane, E., Sautereau, A.-M., Urech, K., Rougé, P. and Lopez, A. (2002) *Biochim. Biophys. Acta* 1559, 145–159.
- [10] García-Olmedo, F., Molina, A., Alamillo, J.M. and Rodríguez-Palenzuela, P. (1998) *Pept. Sci.* 47, 479–491.
- [11] García-Olmedo, F., López-Fando, J.J., Castagnaro, A., Molina, A., Hernández-Lucas, C. and Carbonero, P. (1992) in: *Genes Involved in Plant Defense* (Boller, T. and Meins, F., Eds.), pp. 283–302, Springer-Verlag, Vienna.
- [12] Fernandez de Caleyra, R., González-Pasqual, B., García-Olmedo, F. and Carbonero, P. (1972) *Appl. Microbiol.* 23, 998–1000.
- [13] Caaveiro, J.M.M., Molina, A., González-Mañas, J.M., Rodríguez-Palenzuela, P., García-Olmedo, F. and Goñi, F.M. (1997) *FEBS Lett.* 410, 338–342.
- [14] Hendrickson, W.A. and Teeter, A.M. (1981) *Nature* 290, 107–113.
- [15] Vernon, L.P., Evett, G.E., Zeikus, R.D. and Gray, W.R. (1985) *Arch. Biochem. Biophys.* 238, 18–29.
- [16] Ellman, G.L. (1958) *Arch. Biochem. Biophys.* 74, 443–450.
- [17] Beringer, J.E. (1973) *J. Gen. Microbiol.* 84, 188–198.
- [18] Molina, A., Goy, P.A., Fraile, A., Sanchez-Monge, R. and García-Olmedo, F. (1993) *Plant Sci.* 92, 169–177.
- [19] Merrifield, R.B. (1963) *J. Am. Chem. Soc.* 85, 2149–2154.
- [20] Yokum, S.T. and Barany, G. (2000) in: *Solid-Phase Synthesis* (Kates, S.A. and Albericio, F., Eds.), pp. 79–102, Marcel Dekker, New York.
- [21] Nishiuchi, Y., Nishio, H., Inui, T., Bódi, J. and Kimura, T. (2000) *J. Pept. Sci.* 6, 84–93.
- [22] Andreu, D., Albericio, F., Solé, N.A., Munson, M.C., Ferrer, M. and Barany, G. (1994) *Methods Mol. Biol.* 35, 91–169.
- [23] Chen, Y.H., Yang, J.T. and Chau, K.U. (1974) *Biochemistry* 13, 3350–3359.
- [24] Yang, J.T., Wu, C.S. and Martinez, H.M. (1986) *Methods Enzymol.* 130, 208–269.
- [25] Schwede, T., Diemand, A., Guex, N. and Peitsch, M.C. (2000) *Res. Microbiol.* 151, 107–112.
- [26] Lee, F.S., Chu, Z.T. and Warshel, A. (1993) *J. Comp. Chem.* 14, 161.
- [27] De Samblanx, G.W., Goderis, I.J., Thevesien, K., Raemarekers, R., Fant, F., Borremans, F., Acland, D.P., Osborn, R.W., Patel, S. and Broekaert, W.F. (1997) *J. Biol. Chem.* 272, 1171–1179.
- [28] Berrocal-Lobo, S.A., Moreno, M., López, G., García-Olmedo, F. and Molina, A. (2002) *Plant Physiol.* 128, 951–961.
- [29] Shai, Y. (1999) *Biochim. Biophys. Acta* 1462, 55–70.
- [30] Sitaram, N. and Nagaraj, R. (1999) *Biochim. Biophys. Acta* 1462, 29–54.
- [31] Titarenko, E., Lopez-Solanilla, E., García-Olmedo, F. and Rodríguez-Palenzuela, P. (1997) *J. Bacteriol.* 179, 6699–6704.
- [32] Chou, P.Y. and Fasman, G.D. (1978) *Annu. Rev. Biochem.* 47, 254–276.
- [33] Chakraborty, A. and Baldwin, R.L. (1993) in: *Protein Folding in Vivo and in Vitro* (Cleland, J.L., Ed.), pp. 166–177, American Chemical Society, Washington, DC.