

## Critical Casimir forces and anomalous wetting

S BALIBAR and R ISHIGURO

Laboratoire de Physique Statistique de l'Ecole Normale Supérieure, associé au CNRS et aux Universités Paris 6 et 7, 24 rue Lhomond, 75231 Paris Cedex 05, France  
E-mail: balibar@lps.ens.fr

**Abstract.** We present a review of critical Casimir forces in connection with successive experiments on wetting near the critical point of helium mixtures.

**Keywords.** Wetting; critical; Casimir.

**PACS Nos** 68.35.Rh; 64.60.Fr; 67.60.-g

### 1. Introduction

In 1978, Fisher and de Gennes [1] considered a critical system whose fluctuations are confined between two plates a distance  $L$  apart. They noticed that, with two identical plates, there is a singular contribution of order  $k_B T/L^2$  to the free energy per unit area of the system so that there is an attractive force between the plates which should be of order  $-2k_B T/L^3$  near the critical point at the temperature  $T_c$ . In analogy with the standard Casimir effect, which originates in the confinement of electromagnetic fluctuations between two electrodes, the phenomenon considered by Fisher and de Gennes is now known as the ‘critical Casimir effect’. It has been studied by several groups and recently reviewed by Kardar and Golestanian [2] and by Krech [3] (among others). Despite all the work already done, it seems to us that it is not yet fully understood: the amplitude of the critical Casimir force has not yet been calculated with the boundary conditions corresponding to experimental situations; furthermore, Garcia and Chan demonstrated the existence of this effect with a series of two remarkable experiments [4,5], but some of their quantitative results appear somewhat puzzling to us.

In order to interpret their experiments on wetting by helium mixtures, Ueno *et al* [6] related the critical Casimir effect to another critical phenomenon, which is known as ‘critical point wetting’ which was first predicted by Cahn [7]. This relation had first been proposed by Nightingale and Indekeu [8,9] who had already noticed the potential interest of liquid helium as a model system for the whole issue. Consider a binary liquid mixture below its critical temperature  $T_c$ : it is separated in two phases with an interface in between. The contact angle  $\theta$  of this interface may be non-zero away from  $T_c$ , but Cahn predicted that, as  $T$  tends to  $T_c$ ,  $\theta$  should vanish and the

wall be completely wet by one of the two critical phases. Several theoretical studies have confirmed that critical point wetting is very general and it has been observed in several experiments with different systems [10–12]. However, de Gennes pointed out the importance of long-range forces in Cahn’s situation and opened the possibility of exceptions to critical point wetting if such long range forces are present [13].

Ueno *et al* studied  $^3\text{He}$ – $^4\text{He}$  liquid mixtures in contact with a wall in a series of two experiments: the first one in Kyoto [14] where magnetic resonance imaging (MRI) was used, and the second one in Paris [15] where the contact angle was measured optically. These two experiments showed that, apparently, an exception to critical point wetting had been found in the physics of helium mixtures. In their third article, Ueno *et al* [6] proposed that critical Casimir forces were the long-range forces responsible for this exception. By using Garcia’s measurements, Ueno *et al* found that the critical Casimir forces had the right sign and the right magnitude to explain the non-wetting behavior found in the Paris experiment.

Together with further developments of the theory, these interesting findings urged us to extend and confirm our experimental results. One of the important questions raised by Kardar and Golestanian [2] concerned a new contribution to the critical Casimir force. Indeed, they explained that such a force could exist even if the medium was not close to a critical point. In the case of superfluids, the order parameter has a phase, so that the so-called ‘Goldstone modes’ fluctuate whatever the temperature, in the whole temperature domain where long-range correlations exist. A similar effect was predicted for liquid crystals [16]. For liquid helium, a force originating in Goldstone modes should thus exist in the whole temperature region where it is superfluid. As a consequence, a liquid helium film adsorbed on a wall would be thinner if it is superfluid than if it is normal, even if  $T$  is much lower than  $T_\lambda$ , the superfluid transition temperature. In our context of wetting by helium mixtures, the Goldstone mode contribution to the Casimir force could have led to a non-zero contact angle of the  $^3\text{He}$ – $^4\text{He}$  interface with a wall at low temperature.

In order to test Kardar’s prediction, we changed the geometry of our experimental setup, so that measurements could be done at lower temperature without too much difficulties with optical refraction effects. The results of this ongoing experiment are not yet published [17]. Its preliminary results indicate that the contact angle is in fact zero at low temperature (complete wetting). As we shall see, it does not mean that Kardar’s prediction is wrong, only that the magnitude of this effect is too small to be observed in an indirect measurement such as ours. Furthermore, we have tried to reproduce Ueno’s former results near the critical point of helium mixtures, and, this time, we have found that the angle is very likely to be zero. This now means that there was probably an experimental artefact in Ueno’s series of experiments. It also means that the amplitude of the critical Casimir force is probably smaller than what was deduced from Garcia’s measurements. We thus hope that Garcia’s experiment can also be reproduced, the comparison of its results with theory extended to the region below  $T_c$ , and its relation with the critical point wetting by helium mixtures more critically discussed.

The somewhat difficult goal of this review is to clarify the present status of this confusing situation, particularly the questions concerning the magnitude of the critical Casimir force and its connexion with critical point wetting.

## 2. Critical Casimir forces


Fisher and de Gennes only gave an order of magnitude estimate for the amplitude of the critical Casimir effect at the critical temperature  $T_c$ . One usually writes the critical Casimir force as

$$F(L, T) = \frac{k_B T}{L^3} \vartheta(L/\xi). \quad (1)$$

The ‘scaling function’  $\vartheta(L/\xi)$  depends on the temperature and thickness  $L$  through its ratio to the bulk correlation length  $\xi$ . Near  $T_c$ ,  $\xi$  diverges proportionally to  $t^{-\nu}$  where  $t = (T/T_c - 1)$  is the reduced temperature and  $\nu = 0.67$  for ordinary critical points ( $\nu = 1$  for tri-critical points). Following the seminal paper by Fisher and de Gennes, several theoretical works have brought out important information about the critical Casimir force:

- (1) The sign of the force is given by the sign of the scaling function, and it depends on the symmetry of the boundary conditions. If they are symmetric,  $\vartheta$  is negative and the force is attractive; in the opposite case, if the boundary conditions are antisymmetric,  $\vartheta$  is positive and the force is repulsive.
- (2) The magnitude of the force is generally considered as universal, in particular, at the bulk critical temperature  $T_c$ , where its value is twice the ‘Casimir amplitude’  $\Delta$ , which is the universal value of  $\Theta$  (similar scaling function appearing in the singular contribution to the free energy) at  $T_c$ . Furthermore, the Casimir amplitude depends on the dimension  $N$  of the order parameter. From the work of Nightingale and Indekeu [9,18] and Krech and Dietrich [19], it appears that  $\Delta$  is roughly proportional to  $N$ . For example, it is expected to be twice as large for a superfluid transition ( $N = 2$ ) as for the phase separation of a usual liquid mixture ( $N = 1$ ). It also depends on the boundary conditions, more precisely on their nature, not on the exact details of surfaces [19]. These conditions can be periodic, or the order parameter can vanish at the boundary (‘Dirichlet’ boundary conditions) or its derivative can vanish (‘von Neumann’ conditions).
- (3) With Dirichlet boundary conditions, the critical temperature in the film is significantly displaced with respect to the bulk critical temperature  $T_c$ , and the maximum of the scaling function  $\Theta(L/\xi)$  is expected to be rather different from the Casimir amplitude  $\Delta$ . In fact, as far as we know, there exists no calculation of the scaling functions both below and above  $T_c$  for Dirichlet boundary conditions. According to Krech and Dietrich [19],  $\Delta$  is much smaller for Dirichlet boundary conditions than for periodic ones, but it does not mean that the maximum amplitude of  $\vartheta$  is also much smaller, mainly that the temperature at which this maximum is reached is displaced (as far as we understand).

The calculation of  $\Theta(L/\xi)$  has been done above  $T_c$  by Krech and Dietrich [19], using an  $\epsilon$ -expansion method. For periodic boundary conditions and below  $T_c$ , it has been more recently calculated by Williams in the frame of his vortex loop-model for liquid helium [20]. According to Williams, his calculation below  $T_c$  matches nicely with Krech’s calculation above  $T_c$ , the Casimir amplitude being about  $-0.15$ , close to the maximum amplitude of the scaling function  $\Theta(L/\xi)$ .



**Figure 1.** The scaling function  $\vartheta(x)$  as measured by Garcia and Chan. The horizontal coordinate is  $x = tL^{1/\nu}$  where  $t$  is the reduced temperature and  $L$  the film thickness,  $\nu$  the critical exponent of the bulk correlation length  $\xi$ ; it is measured in  $\text{\AA}^{1/\nu}$  units. These results could only be compared with Krech's theory above the critical temperature  $T_\lambda$ , i.e. for  $x > 0$ , where  $\vartheta(x)$  has a very small tail. Different symbols correspond to different film thicknesses.

In their first experiment, Garcia and Chan observed the thinning of a pure liquid helium film near the superfluid transition at  $T_\lambda$ . This film was adsorbed on a copper electrode and most of the thinning occurred in a small temperature region near  $T_\lambda$ . They analyzed it in terms of the critical Casimir effect and extracted a scaling function  $\vartheta(L/\xi)$  which was very similar in shape with calculations (see figure 1), for example the recent ones by Dantchev and Krech [21]. Garcia's scaling function displays important features which deserve several comments:

- (1) The maximum amplitude of  $\vartheta$  does not occur at the bulk critical temperature  $T_c$ . This was expected because the order parameter for superfluidity vanishes on both sides of the superfluid film, so that the superfluid transition temperature is depressed in the film:  $T_c^{\text{film}} < T_c^{\text{bulk}}$ . It occurs significantly below  $T_c$ , for  $x = tL^{1/\nu} \approx -10$  (the reduced temperature is taken negative below  $T_c$ . Note also that, in both articles by Garcia and Chan, the horizontal coordinate  $x = tL^{1/\nu} = (L\xi_0/\xi)^{1/\nu}$  is not dimensionless, but close to  $(L/\xi)^{1/\nu}$  since  $L$  is taken in  $\text{\AA}$  and the quantity  $\xi_0$  is about 1  $\text{\AA}$  [22]). The magnitude of the scaling function above  $T_c$  is very small, as predicted by Krech and Dietrich [19]. Garcia and Chan claim that their measurement of  $\vartheta$  agrees with the calculation, but this only concerns the small tail at  $T > T_c$ , where the signal/noise ratio is poor, while most of the observed effect occurs below  $T_c$ .
- (2) Garcia and Chan found that  $\vartheta(L/\xi)$  reaches maximum negative values which vary from  $-1.5$  to  $-2$  as a function of the film thickness  $L$ . This is doubly surprising, because (a) dependence on  $L$  seems to contradict the predicted universality and (b) no calculation has ever found such large amplitudes for  $\vartheta$ . In the various situations which have been calculated, the theoretical results

are 5 to 50 times smaller. This is a serious problem which needs further studies: new experiments should identify the origin of the  $L$ -dependence, and  $\vartheta$  should be calculated below  $T_c$  with Dirichlet boundary conditions.

- (3) They also found indications that the scaling function does not tend to zero in the low-temperature limit, away from  $T_c$ . One possible explanation for this is the confinement of Goldstone modes invoked by Kardar and Golestanian [2]. The amplitude of the Goldstone mode contribution looked too small to explain the rather large negative value of  $\vartheta(T \rightarrow 0)$  found by Garcia and Chan, but a more recent calculation by Zandi *et al* proposes that, the film surface being mobile, a contribution from third sound modes at the surface of a superfluid film should be added to the one coming from Goldstone modes, so that the total attractive force acting on the film surface is larger than first calculated [23]. One could even imagine that phonons also contribute to the force at low temperature [24].

### 3. Critical point wetting and Ueno's experiments


Following Cahn's argument [7], let us consider a binary liquid mixture near its critical temperature  $T_c$ . Below  $T_c$ , the mixture is separated into a concentrated 'c-phase' and a diluted 'd-phase'. In the case of partial wetting, the contact angle  $\theta$  of the  $cd$ -interface against a substrate 's' obeys the Young–Dupré relation [25]:

$$\cos \theta = \frac{\sigma_{sc} - \sigma_{sd}}{\sigma_i} = \frac{\delta\sigma}{\sigma_i}, \quad (2)$$


where  $\sigma_{sc}$ ,  $\sigma_{sd}$ , and  $\sigma_i$  are interfacial free energies between  $s$ - and  $c$ -phases,  $s$ - and  $d$ -phases and  $c$ - and  $d$ -phases respectively. If one assumes that  $\delta\sigma$  is proportional to the difference in concentration ( $\delta X = X_c - X_d$ ), one finds that, for ordinary critical points,  $\delta\sigma \propto \delta X \propto t^{0.33}$ . Since the interfacial tension  $\sigma_i \propto t^{1.28}$ , one finds that the numerator in eq. (2) vanishes more slowly than the denominator, so that the cosine increases as  $T$  approaches  $T_c$ , and the contact angle reaches zero at some temperature below  $T_c$ : a 'critical point wetting' transition occurs. In the case of a tri-critical point, as is the case for liquid helium mixtures because phase separation occurs at the same temperature  $T_t$  as superfluidity (see figure 2), the (mean field) exponents are respectively 2 and 1 [22,26], so that the same reasoning should apply and complete wetting should occur as  $T$  approaches  $T_t$ . The above argument is too simple but more careful analysis have proved that this is qualitatively correct in the absence of long-range forces [8].

In their two successive experiments, Ueno *et al* did not find Cahn's critical point wetting when studying liquid helium mixtures. Below their tri-critical temperature  $T_t$ , these mixtures are separated in a 'c-phase' which is concentrated in  $^3\text{He}$ , and a 'd-phase' which is diluted in  $^3\text{He}$ , consequently rich in  $^4\text{He}$ . In the Kyoto experiment [14], Ueno *et al* measured the profile of the  $cd$ -interface near a wall made of epoxy glue. They used a magnetic resonance imaging technique (MRI) and found a non-zero contact angle (see figure 3). In the Paris experiment [15], Ueno *et al* measured the interface profile with an optical interferometric technique. They also found that the contact angle  $\theta$  was non-zero below  $T_t$ ; moreover, they found that  $\theta$  increased as  $T$  approached  $T_t$  (see figure 4). This was not compatible with critical


point wetting, so that, in their theoretical article [6], Ueno *et al* reconsidered Cahn's argument after identifying three long-range forces which are present in their experimental situation.



**Figure 2.** The phase diagram of liquid helium mixtures (left) and a schematic representation of the partial wetting of a wall by the phase-separated mixture (right).



**Figure 3.** The contact angle  $\theta$  of the  $^3\text{He}$ - $^4\text{He}$  interface with a wall was found non-zero in Ueno's experiments in Kyoto (crosses). They used a magnetic resonance imaging (MRI) technique whose accuracy was not good near the tri-critical point, because of the large scattering of data points in this temperature region. Agreement with this anomalous behavior was found in a later experiment by Ueno *et al* in Paris. However, in their more recent experiment, Ishiguro and Balibar found  $\theta = 0$  both at low temperature and near the critical point.



**Figure 4.** The contact angle near the tri-critical point. This is an enlargement of the graph shown in the previous figure. The experiment by Ueno *et al* in Paris (2003) showed that  $\theta$  increased instead of tending to zero as  $T$  approached the tri-critical temperature  $T_c$ . However, this anomalous behavior was not confirmed in the more recent experiment done by Ishiguro and Balibar, who found  $\theta = 0$ .


The van der Waals force is attractive on atoms and  $^4\text{He}$  atoms occupy a smaller volume than  $^3\text{He}$  atoms because their quantum fluctuations are weaker (their mass is larger). As a result, the van der Waals attraction on the  $d$ -phase is stronger than on the  $c$ -phase, and a  $c$ -phase is always separated from a solid wall (in figure 2, a sapphire window) by a film of  $d$ -phase. Being attractive on atoms, the van der Waals field induces an effective force which is *repulsive* on the film surface. In the absence of other long-range forces, a film of finite thickness would only exist off-equilibrium, but as equilibrium is approached the film thickness would diverge and complete wetting by the  $d$ -phase would occur. Romagnan *et al* [27,28] found some experimental evidence for this, but their measurements were limited to thicknesses up to about 80 Å only. The sketch in figure 2 corresponds to a situation where another long-range force acts on the film. Being attractive, this other force is able to counterbalance the van der Waals field and to limit the film thickness, so that the macroscopic contact angle is non-zero.

In the presence of such long-range forces, the way to calculate the surface energies, and consequently the contact angle, is to integrate the so-called ‘disjoining pressure’  $\Pi(L)$  which is nothing but the sum of the forces acting on the film surface [29]:

$$\cos(\theta) = \frac{\sigma_{sc} - \sigma_{sd}}{\sigma_i} = 1 + \frac{\int_{l_{eq}}^{\infty} \Pi(l) dl}{\sigma_i}. \quad (3)$$


In addition to the van der Waals force which is known, Ueno *et al* explained that there is a critical Casimir force. This is because the  $d$ -phase film is superfluid, while the  $c$ -phase is normal. The order parameter of superfluidity is non-zero inside the

film but it has to vanish on both sides. This symmetric vanishing should produce an attractive Casimir force on the film surface. Since there exists no calculation with such Dirichlet boundary conditions yet, Ueno *et al* [6] used Garcia's measurement of  $\vartheta(L/\xi)$  to calculate the contribution of the critical Casimir effect to the disjoining pressure. They also included the Helfrich force, which is repulsive, due to the cutoff of capillary modes at long wavelength by the presence of the nearby wall [30]. At a reduced temperature  $t = -0.01$  below  $T_t$ , they found that the critical Casimir force was stronger than the other two in the thickness range from 0 to 400 Å (see figure 5). According to this calculation, the equilibrium film thickness was thus 400 Å, and Ueno *et al* could calculate the contact angle by integration of the disjoining pressure (eq. (3)). They found the contact angle to be  $45^\circ$ , which is in good agreement with their measurement (see figure 4). Moreover, they argued that the Casimir force could be twice as large in their experiment as in Garcia's experiment, because it is a tri-critical point instead of an ordinary critical point such as the lambda point of pure liquid  $^4\text{He}$ . According to this argument, they found that the contact angle was  $60^\circ$  at  $t = 0.01$ , which is in even better agreement with their experimental results. Furthermore, Ueno *et al* also extracted values for the interfacial tension  $\sigma_i$  and found good agreement with previous measurements (see figure 6). Although the support of experiments by theory and vice versa looked rather strong, we decided to repeat the experiment in a different geometry and with more careful analysis of the interferometric images. As we shall see, our new results show that there was probably an artefact in Ueno's experiment, and also that the amplitude of the Casimir force might be smaller than what Ueno *et al* deduced directly from Garcia's measurements.



**Figure 5.** The 2003 calculation by Ueno *et al* of the various forces acting on the film surface. The amplitude of the Casimir force was taken directly from the measurement by Garcia and Chan in 1999. It was apparently stronger than the van der Waals effective force which is positive, meaning repulsive on the film surface. After adding the Helfrich force, Ueno *et al* found a total disjoining pressure crossing zero at  $l = 400$  Å, the equilibrium film thickness. A finite film thickness means partial wetting and quantitative agreement was found with the measurements by Ueno *et al* in 2003.






**Figure 6.** Various measurements of the interfacial tension  $\sigma_i$  between the  $c$ - and the  $d$ -phases in the vicinity of the tri-critical temperature  $T_t = 870$  mK. Within the error bars, all data agree with the  $t^2$  critical behavior measured by Leiderer *et al* in 1977.

#### 4. The new experiment in Paris


In Ueno's experiment, the angle of incidence of the laser beam was large on the  $cd$ -interface, that is far from normal. As a consequence, as soon as the difference in optical index between the  $c$ - and the  $d$ -phases was large, refraction at the  $cd$ -interface induced a substantial difference in orientation between the incoming and the outgoing beams. The fringe pattern was distorted and the calculation of the interface profile too difficult. This is the reason why Ueno *et al* could not measure the contact angle below about 0.8 K. In order to do this, and be able to look for a possible effect of Goldstone modes on the wetting by helium mixtures, we rotated the cell by nearly  $90^\circ$ , so that the incidence was now close to normal. We also tried to adjust the optical cavity more carefully, in order to obtain straight fringes. The interface profile is obtained by subtracting the optical path without the interface from the optical path with the interface (see figure 7). This is rather delicate and needs a very accurate knowledge of the fringe pattern in the absence of an interface.

Close to  $T_t$  where the index difference is very small, any error in the subtracted background produces large errors in the calculated profile. We believe that this is the origin of artefacts in the profiles found by Ueno *et al*, whose fringe patterns were severely bent, due to inhomogeneous stresses on the two cell windows. In our new experiment, we have found zero contact angle in the whole temperature range. Two examples of profiles are shown in figure 8. At low temperature (271 mK) and also at 840 mK, near  $T_t$ , the interface bends upwards and meets the window tangentially. If Ueno *et al* had been right, the interface would have bent downwards



**Figure 7.** The profile of the  $cd$ -interface between phase-separated liquid helium mixtures is calculated from the difference in optical paths between a pattern with an interface (left) and a pattern without interface (right). These two fringe patterns were recorded at  $T = 636$  mK.

near  $T_t$  (note that, in the two figures, the vertical scale is not the same as the horizontal one, so that the window looks more tilted than in reality ( $20^\circ$ )). A critical discussion of the profile extraction and a full presentation of results will be presented in a later publication where we shall explain more precisely why we now believe that there was an artefact in Ueno's work. For these conference proceedings, we only briefly discuss the meaning of our new measurements.



**Figure 8.** The profile of the  $cd$ -interface between phase-separated liquid helium mixtures and a sapphire window at 271 mK (left) and 840 mK (right). The arrows indicate the magnitude of the capillary length and the broken lines are fits from Laplace's equation. The sapphire window is tilted by  $20^\circ$  with respect to the horizontal. We have found complete wetting by the  $^4\text{He}$ -rich  $d$ -phase:  $\theta = 0$ .

## 5. Discussion


The van der Waals effective force acting on a  $cd$ -interface is repulsive, and is proportional to the difference in the average volume per atom in each phase  $V_c - V_d$ :

$$\Pi_{\text{vdW}} = \frac{A_0}{L^3} \left( \frac{1}{V_d} - \frac{1}{V_c} \right) \quad (4)$$

with  $A_0 \approx 1000 \text{ K} \cdot \text{\AA}^{-3}$ . At low temperature,  $V_c = 61.15 \text{ \AA}^3$  and  $V_d = 46.56 \text{ \AA}^3$  [31], so that, far below  $T_t = 0.87 \text{ K}$ , the effective van der Waals force is about  $5/L^3$  in  $\text{K} \cdot \text{\AA}^{-3}$  units. For partial wetting to occur in the low-temperature limit, the contribution of the Goldstone modes to the Casimir force would need to be more negative than  $-5/L^3$ . In their review article [2], Kardar and Golestanian propose  $-0.048 k_B T / L^3$  which looks much too small. Following the recent work of Zandi *et al* [23], one should also account for the existence of fluctuations at the film surface, i.e. third sound modes. Their contribution should add to that of Goldstone modes and lead to a total Casimir force which is three times more negative than previously thought, about  $-0.15 k_B T / L^3$ . However, this looks still too small compared to the van der Waals field. In a sense, it is not surprising that we found complete wetting at low temperature, but in the first experiment done by Ueno *et al* in Kyoto, partial wetting had been found and, since Garcia's measured value of the Casimir force is much larger than in available calculations, it was worth checking that complete wetting occurred at low  $T$ .

As for the vicinity of  $T_t$ , we have recalculated what should be the disjoining pressure if one assumed that, in our case, the magnitude of the Casimir force was only one-fifth of what had been measured by Garcia and Chan [4]. There could be several reasons for this. Firstly, Garcia's results are indeed at least five times larger than in any available calculation. Secondly, their measurement was done with a pure liquid  $^4\text{He}$  film, while we are dealing with a mixture. Note that in a later experiment, Garcia and Chan also studied mixture films [5] but they measured the force acting on a liquid-gas interface, not on the  $cd$ -interface as we do. In our case, a rigorous estimate should account for the existence of two competing effects. The confinement of superfluidity leads to an attractive force because the order parameter (the amplitude of the macroscopic wave function) vanishes symmetrically on both boundaries of the film. However, there are also fluctuations of concentration near  $T_t$ . Due to the van der Waals field, the film is more diluted in  $^3\text{He}$  near the solid wall than near the  $cd$ -interface, so that, for concentration fluctuations, the boundary conditions are anti-symmetric. This is why, for an ordinary liquid mixture with no superfluidity, the Casimir force would add to the van der Waals field to favor wetting by the  $d$ -phase. Of course, superfluidity and concentration fluctuations are coupled near  $T_t$ , and the effect of superfluidity should dominate because the dimension of its order parameter is  $N = 2$  instead of  $N = 1$  for the concentration. Still, we expect a rigorous calculation to find that the effect of concentration fluctuations decreases the effect of superfluidity fluctuations.

As shown in figure 9, when estimating the amplitude of the Casimir force as only one-fifth of what was measured by Garcia and Chan, we find that complete wetting is restored. Compared to the results shown in figure 5, the disjoining pressure is now positive for all thicknesses. Given our most recent measurements [17], we now



**Figure 9.** A modified calculation of the disjoining pressure acting on the film surface, where the amplitude of the critical Casimir force has been taken as one-fifth only of the experimental result obtained by Garcia and Chan. Contrary to what was shown in figure 5, one now finds that the van der Waals force always dominates over the Casimir force, so that the disjoining pressure is always positive, leading to a macroscopic film thickness at equilibrium, and complete wetting.

believe that, in our experiment, that is near a sapphire window, the critical Casimir force is too weak compared to the van der Waals field, so that complete wetting occurs, as usual. To find an exception to critical point wetting, one should try using a substrate exerting a much weaker van der Waals field than ordinary insulating materials, and this does not look easy to us.

## Acknowledgements

The authors are grateful to E Rolley and F Caupin for their help in the experiments, to R Garcia and M Chan for several discussions of their experiments and allowing us to reproduce one of their figures, also to S Dietrich and M Krech for numerous discussions on the theoretical aspects of the whole issue. R Ishiguro acknowledges support from the Japan Society for the Promotion of Science.

## References

- [1] M Fisher and P G de Gennes, *C. R. Acad. Sci. Paris* **B287**, 209 (1978)
- [2] M Kardar and R Golestanian, *Rev. Mod. Phys.* **71**, 1233 (1999)
- [3] M Krech, *J. Phys. Condens. Matter* **11**, R391 (1999)
- [4] R Garcia and M H W Chan, *Phys. Rev. Lett.* **83**, 1187 (1999)
- [5] R Garcia and M H W Chan, *Phys. Rev. Lett.* **88**, 086101 (2002)
- [6] T Ueno, S Balibar, F Caupin, T Mizusaki and E Rolley, *Phys. Rev. Lett.* **90**, 116102 (2003)
- [7] J W Cahn, *J. Chem. Phys.* **66**, 3667 (1977)

- [8] M P Nightingale and J O Indekeu, *Phys. Rev.* **B32**, 3364 (1985)
- [9] J Indekeu, *Faraday Transactions II*, **82**, 1835 (1986)
- [10] R B Heady and J W Cahn, *J. Chem. Phys.* **58**, 896 (1973)
- [11] See, for instance, the review articles by J O Indekeu, *Acta Phys. Pol.* **B26**, 1065 (1995) and by D Bonn and D Ross, *Rep. Prog. Phys.* **64**, 1085 (2001)
- [12] D Bonn, H Kellay and G H Wegdam, *Phys. Rev. Lett.* **69**, 1975 (1992)
- [13] P G de Gennes, *J. Phys. Lett.* **42**, L-377 (1981)
- [14] T Ueno, M Fujisawa, K Fukuda, Y Sasaki and T Mizusaki, *Physica* **B284–288**, 2057 (2000)
- [15] T Ueno, S Balibar, F Caupin, T Mizusaki and E Rolley, *J. Low Temp. Phys.* **130**, 543 (2003)
- [16] A Ajdari, L Peliti and J Prost, *Phys. Rev. Lett.* **66**, 1482 (1991)
- [17] R Ishiguro and S Balibar, unpublished
- [18] M P Nightingale and J O Indekeu, *Phys. Rev. Lett.* **54**, 1824 (1985)
- [19] M Krech and S Dietrich, *Phys. Rev. Lett.* **66**, 345 (1991)
- [20] G Williams, *Phys. Rev. Lett.* **92**, 197003 (2004)
- [21] D Dantchev and M Krech, *Phys. Rev.* **E69**, 046119 (2004)
- [22] P Leiderer, H Poisel and M Wanner, *J. Low Temp. Phys.* **28**, 167 (1977)
- [22] P Leiderer, D R Watts and W W Webb, *Phys. Rev. Lett.* **33**, 483 (1974)
- [23] R Zandi, J Rudnick and M Kardar, preprint and communication at the APS March Meeting (Montreal, 2004)
- [24] M H Nahale and M W Cole, *Surf. Sci.* **172**, 311 (1986)
- [25] T Young, *Philos. Trans. R. Soc. London* **95**, 65 (1805)
- [26] B Widom, in *Phase transitions and critical phenomena* edited by C Domb and S Green (Academic, New York, 1972) vol. 2, chapter III
- [27] J P Romagan, J P Laheurte, J C Noiray and W F Saam, *J. Low Temp. Phys.* **30**, 425 (1978)
- [28] D Sornette and J P Laheurte, *J. Phys. (Paris)* **47**, 1951 (1986)
- [29] D Ross, D Bonn and J Meunier, *Nature* **400**, 737 (1999)
- [30] W Helfrich and R M Servuss, *Nuovo Cimento* **D3**, 137 (1984)
- [31] A Sato, K Ohishi and M Suzuki, *J. Low Temp. Phys.* **107**, 165 (1997)