

ON NUMERICAL RANGES OF GENERALIZED DERIVATIONS AND RELATED PROPERTIES

SEN-YEN SHAW

(Received 28 February 1982; revised 11 November 1982)

Communicated by G. Brown

Abstract

This paper is concerned with the numerical range and some related properties of the operator $\Delta|\mathfrak{S}: T \rightarrow AT - TB$ ($T \in \mathfrak{S}$), where A, B are (bounded linear) operators on the normed linear spaces \mathbf{X} and \mathbf{Y} , respectively, and \mathfrak{S} is a linear subspace of the space $\mathcal{L}(\mathbf{Y}, \mathbf{X})$ of all operators from \mathbf{Y} to \mathbf{X} . \mathfrak{S} is assumed to contain all finite operators, to be invariant under Δ , and to be suitably normed (not necessarily with the operator norm). Then the algebra numerical range of $\Delta|\mathfrak{S}$ is equal to the difference of the algebra numerical ranges of A and B . When $\mathbf{X} = \mathbf{Y}$ and $\mathfrak{S} = \mathcal{L}(\mathbf{X})$, Δ is Hermitian (resp. normal) in $\mathcal{L}(\mathcal{L}(\mathbf{X}))$ if and only if $A - \lambda$ and $B - \lambda$ are Hermitian (resp. normal) in $\mathcal{L}(\mathbf{X})$ for some scalar λ ; if $\mathbf{X} := \mathbf{H}$ is a Hilbert space and if \mathfrak{S} is a C^* -algebra or a minimal norm ideal in $\mathcal{L}(\mathbf{H})$, then any Hermitian (resp. normal) operator on \mathfrak{S} is of the form $\Delta|\mathfrak{S}$ for some Hermitian (resp. normal) operators A and B . $AT = TB$ implies $A^*T = TB^*$, provided that A and B^* are hyponormal operators on the Hilbert spaces \mathbf{H}_1 and \mathbf{H}_2 , respectively, and T is a Hilbert-Schmidt operator from \mathbf{H}_2 to \mathbf{H}_1 .

1980 *Mathematics subject classification* (*Amer. Math. Soc.*): 47 A 12, 47 B 15, 47 B 47.

Keywords and phrases: algebra numerical range, derivation, Hermitian and normal operators, hyponormal operator, normed operator ideal, Schatten p -class, C^* -algebra, Hilbert-Schmidt operator.

0. Introduction

Let \mathbf{X} be a complex normed linear space and let $\mathcal{L}(\mathbf{X})$ be the algebra of all (bounded linear) operators on \mathbf{X} . The spatial numerical range of an operator A on \mathbf{X} is given by $W(A; \mathcal{L}(\mathbf{X})) := \{f(Ax); (x, f) \in \pi(\mathbf{X})\}$, where $\pi(\mathbf{X})$ denotes the set of all pairs $(x, f) \in \mathbf{X} \times \mathbf{X}'$ such that $\|x\| = \|f\| = f(x) = 1$. The algebra numerical range of A in $\mathcal{L}(\mathbf{X})$ is given by $V(A; \mathcal{L}(\mathbf{X})) := \{F(A); (I, F) \in \pi(\mathcal{L}(\mathbf{X}))\}$, where I is the identity operator. It is known that $V(A; \mathcal{L}(\mathbf{X}))$ is compact

and is the closed convex hull of $W(A; \mathcal{L}(\mathbf{X}))$. When $\mathbf{X} = \mathbf{H}$ is a Hilbert space with inner product $\langle \cdot, \cdot \rangle$, $W(A; \mathcal{L}(\mathbf{X}))$ reduces to the classical numerical range $W(A) := \{ \langle Ax, x \rangle; x \in \mathbf{H}, \|x\| = 1 \}$ (see [4] for details).

If B is an operator on another normed linear space Y , then the generalized derivation $\Delta = \Delta_{A,B}$, defined by $\Delta(T) := AT - TB$, is an operator on the space $\mathcal{L}(Y, X)$ of all operators from Y to X . In this paper, we consider the numerical ranges of restrictions of Δ to certain invariant subspaces \mathcal{S} and some consequences. First, the algebra numerical range of $\Delta|_{\mathcal{S}}$ is shown to be the difference of $V(A; \mathcal{L}(X))$ and $V(B; \mathcal{L}(Y))$, provided that \mathcal{S} contains all finite-rank operators and is suitably normed. Then it is applied to determine when Δ or $\Delta|_{\mathcal{S}}$ is Hermitian or normal, and to derive a Fuglede-Putnam theorem for hyponormal operators. The results will extend some theorems of Kyle [8], Sourour [13] and Berberian [3], respectively.

1. The numerical range

We will assume that \mathcal{S} is a linear subspace of $\mathcal{L}(Y, X)$ equipped with a norm $\|\cdot\|$ (possibly different from the operator norm $\|\cdot\|$) such that the following conditions are satisfied:

(1) $A\mathcal{S} \subset \mathcal{S}$ and $\mathcal{S}B \subset \mathcal{S}$; (2) If $D \in \mathcal{L}(X)$, $T \in \mathcal{S}$, $E \in \mathcal{L}(Y)$ and $DTE \in \mathcal{S}$, then $\|DTE\| \leq \|D\| \|T\| \|E\|$; (3) $\|T\| \leq \|\|T\|\|$ for all T in \mathcal{S} , and the equality holds whenever T has rank one; (4) \mathcal{S} contains all finite rank operators from Y to X .

It follows from (1) that \mathcal{S} is an invariant subspace of $\Delta = \Delta_{A,B}$, and from (2) that the restriction $\Delta|_{\mathcal{S}}$ of Δ is a bounded linear operator on $(\mathcal{S}, \|\cdot\|)$. We consider the numerical range of $\Delta|_{\mathcal{S}}$.

THEOREM 1.1. *For operators $A \in \mathcal{L}(X)$ and $B \in \mathcal{L}(Y)$ let $(\mathcal{S}, \|\cdot\|)$ be the normed linear space as mentioned above. Then*

$$(*) \quad V(\Delta|_{\mathcal{S}}; \mathcal{L}(\mathcal{S})) = V(A; \mathcal{L}(X)) - V(B; \mathcal{L}(Y)).$$

PROOF. We first prove that the left side is contained in the right side. So, let λ be an arbitrary element of $V(\Delta|_{\mathcal{S}}; \mathcal{L}(\mathcal{S}))$. Then $\lambda = f(\Delta|_{\mathcal{S}})$ for some $f \in (\mathcal{L}(\mathcal{S}))'$ such that $\|f\| = f(I_{\mathcal{L}(\mathcal{S})}) = 1$, where $I_{\mathcal{L}(\mathcal{S})}$ denotes the identity operator in $\mathcal{L}(\mathcal{S})$. It is clear that the set \mathfrak{N} (resp. \mathfrak{U}) of all $D \in \mathcal{L}(X)$ (resp. $E \in \mathcal{L}(Y)$) such that $D\mathcal{S} \subset \mathcal{S}$ (resp. $\mathcal{S}E \subset \mathcal{S}$) is a linear subspace of $\mathcal{L}(X)$ (resp. $\mathcal{L}(Y)$) containing $I_{\mathcal{L}(X)}$ and A (resp. $I_{\mathcal{L}(Y)}$ and B). Define a linear functional F on \mathfrak{N} by $F(D) = f(L_D|_{\mathcal{S}})$ ($D \in \mathfrak{N}$) and a linear functional G on \mathfrak{U} by $G(E) = f(R_E|_{\mathcal{S}})$ ($E \in \mathfrak{U}$), where L_D and R_E stand for the left multiplication by D and the right multiplication by

E , respectively. Now the Hahn-Banach theorem guarantees the existence of \hat{F} in $(\mathcal{L}(\mathbf{X}))'$ and \hat{G} in $(\mathcal{L}(\mathbf{Y}))'$ such that $\hat{F}|_{\mathfrak{N}} = F$, $\|\hat{F}\| = \|F\|$, $\hat{G}|_{\mathfrak{N}} = G$ and $\|\hat{G}\| = \|G\|$. Since $\hat{F}(I_{\mathcal{L}(\mathbf{X})}) = F(I_{\mathcal{L}(\mathbf{X})}) = f(I_{\mathcal{L}(\mathcal{S})}) = 1$ and since $|F(D)| \leq \|f\| \|L_D|_{\mathcal{S}}\| = \sup\{\|DT\|; T \in \mathcal{S}, \|T\| = 1\} \leq \|D\|$ for all D in \mathfrak{N} , we have that $\|\hat{F}\| = \|F\| \leq 1 = \hat{F}(I_{\mathcal{L}(\mathbf{X})}) \leq \|\hat{F}\|$, that is, $(I_{\mathcal{L}(\mathbf{X})}, \hat{F})$ belongs to $\pi(\mathcal{L}(\mathbf{X}))$. Similarly, we have $(I_{\mathcal{L}(\mathbf{Y})}, \hat{G}) \in \pi(\mathcal{L}(\mathbf{Y}))$. Hence

$$\begin{aligned} \lambda &= f(\Delta|_{\mathcal{S}}) = f(L_A|_{\mathcal{S}}) - f(R_B|_{\mathcal{S}}) \\ &= F(A) - G(B) = \hat{F}(A) - \hat{G}(B) \\ &\in V(A; \mathcal{L}(\mathbf{X})) - V(B; \mathcal{L}(\mathbf{Y})). \end{aligned}$$

To prove the other inclusion, it suffices to show that

$$V(\Delta|_{\mathcal{S}}; \mathcal{L}(\mathcal{S})) \supset W(A; \mathcal{L}(\mathbf{X})) - W(B; \mathcal{L}(\mathbf{Y}))$$

since the closed convex hull of the set on the right side is $V(A; \mathcal{L}(\mathbf{X})) - V(B; \mathcal{L}(\mathbf{Y}))$, by an elementary proof. So, let $\alpha = g(Ax)$ with $(x, g) \in \pi(\mathbf{X})$ and let $\beta = h(By)$ with $(y, h) \in \pi(\mathbf{Y})$. Using the usual notation $x \otimes h$ for the rank-one operator: $z \mapsto h(z)x$ ($z \in \mathbf{Y}$), we define the linear functional P on $\mathcal{L}(\mathcal{S})$ by

$$P(\Omega) := g([\Omega(x \otimes h)]y) \quad (\Omega \in \mathcal{L}(\mathcal{S})).$$

Clearly we have $P(I_{\mathcal{L}(\mathcal{S})}) = 1$ and, by (3) and (4),

$$\begin{aligned} |P(\Omega)| &\leq \|g\| \|\Omega(x \otimes h)\| \|y\| \leq \|\Omega(x \otimes h)\| \leq \|\Omega\| \|x \otimes h\| \\ &= \|\Omega\| \|x \otimes h\| = \|\Omega\|, \end{aligned}$$

that is, $(I_{\mathcal{L}(\mathcal{S})}, P) \in \pi(\mathcal{L}(\mathcal{S}))$. Hence $V(\Delta|_{\mathcal{S}}; \mathcal{L}(\mathcal{S}))$ contains the number

$$\begin{aligned} P(\Delta|_{\mathcal{S}}) &= g(A(x \otimes h)y - (x \otimes h)By) \\ &= g(Ax)h(y) - g(x)h(By) = \alpha - \beta. \end{aligned}$$

The proof is complete.

REMARK. Conditions (3) and (4) are used only in proving the direction “ \supset ”, therefore the inclusion “ \subset ” will hold for any subspace satisfying (1) and (2). That (4) is essential for the direction “ \supset ” is easily seen from the example where $\mathbf{X} = \mathbf{Y} = C^2$, $A = B = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$, $\mathcal{S} = \text{span}\{A\}$ and $\|\cdot\| = \|\cdot\|$.

The following are some examples of qualified subspaces $(\mathcal{S}, \|\cdot\|)$:

(a) Components in $\mathcal{L}(\mathbf{Y}, \mathbf{X})$ of all the operator ideals on Banach spaces, as studied in Chapter 1 of [9], such as the classes of finite operators, approximate operators, compact operators, weakly compact operators, completely continuous

operators, unconditionally summing operators, separable operators, Kato operators and Pelczynski operators.

(b) Components in $\mathcal{L}(\mathbf{Y}, \mathbf{X})$ of all the normed operator ideals on Banach spaces, as studied in Chapter 6 of [9], such as nuclear operators, integral operators, absolutely summing operators and Hilbert operators. (Compare the assumptions on $(\mathfrak{S}, \|\cdot\|)$ with Definitions 6.1.1, 6.2.2 and Propositions 6.1.4, 6.1.5 of [9].)

(c) The Schatten p -class $C_p(\mathbf{H}_2, \mathbf{H}_1)$ ($1 \leq p < \infty$) or approximable operators from a Hilbert space \mathbf{H}_2 to another Hilbert space \mathbf{H}_1 , that is, operators T such that $\|T\|_p := [\text{trace}(T^*T)^{p/2}]^{1/p} < \infty$, (see [9, page 216]). In the case where $p = 1$ these are the operators of trace class (nuclear operators), and $p = 2$ yields the Hilbert space of Hilbert-Schmidt operators (see [2, Chapter 12]).

COROLLARY 1.2. *For any $A \in \mathcal{L}(\mathbf{X})$ and $B \in \mathcal{L}(\mathbf{Y})$,*

$$V(\Delta; \mathcal{L}(\mathcal{L}(\mathbf{Y}, \mathbf{X}))) = V(A; \mathcal{L}(\mathbf{X})) - V(B; \mathcal{L}(\mathbf{Y})).$$

This contains Kyle’s result [8] (for the case $\mathbf{X} = \mathbf{Y}$) as a special case.

COROLLARY 1.3. *Let A and B be any operators on Hilbert spaces \mathbf{H}_1 and \mathbf{H}_2 , respectively, and let $C_p(\mathbf{H}_2, \mathbf{H}_1)$ be normed with $\|\cdot\|$ or $\|\cdot\|_p$. Then*

$$V(\Delta|C_p(\mathbf{H}_2, \mathbf{H}_1); \mathcal{L}(C_p(\mathbf{H}_2, \mathbf{H}_1))) = W(A)^- - W(B)^-.$$

Thus Corollary 1.3 becomes a numerical range analogue of Fialkow’s [5] formula for spectra: $\sigma(\Delta|C_p(\mathbf{H})) = \sigma(A) - \sigma(B)$.

We end this section by deriving from Theorem 1.1 the following known property, which will be of use in Section 2.

COROLLARY 1.4. *If $AT = TB$ holds for all rank-one operators T in $\mathcal{L}(\mathbf{Y}, \mathbf{X})$, then $A = \lambda I_{\mathcal{L}(\mathbf{X})}$ and $B = \lambda I_{\mathcal{L}(\mathbf{Y})}$ for some scalar λ .*

PROOF. Take \mathfrak{S} to be the space of all finite rank operators. Then $\Delta|\mathfrak{S} = 0$ and so $V(A; \mathcal{L}(\mathbf{X})) - V(B; \mathcal{L}(\mathbf{Y})) = V(\Delta|\mathcal{L}(\mathfrak{S})) = \{0\}$, or equivalently, $V(A; \mathcal{L}(\mathbf{X})) = V(B; \mathcal{L}(\mathbf{Y})) = \{\lambda\}$ for some scalar λ . It follows that $V(A - \lambda I_{\mathcal{L}(\mathbf{X})}; \mathcal{L}(\mathbf{X})) = \{0\}$ and

$$\|A - \lambda I_{\mathcal{L}(\mathbf{X})}\| \leq e \max\{|\mu|; \mu \in V(A - \lambda I_{\mathcal{L}(\mathbf{X})}; \mathcal{L}(\mathbf{X}))\} = 0$$

(see [4, page 34]). Hence $A = \lambda I_{\mathcal{L}(\mathbf{X})}$, and similarly $B = \lambda I_{\mathcal{L}(\mathbf{Y})}$.

2. Hermitian and normal derivations

An operator A on a normed linear space X is said to be Hermitian if its numerical range is contained in the real line and it is normal if $A = H + iK$ for some commuting Hermitian operators H and K . In this section we try to answer partly the question about when the operator $\Delta|_{\mathfrak{S}}$ is Hermitian or normal.

First, from the formula (*) comes immediately the following

COROLLARY 2.1. *Let $(\mathfrak{S}, |||\cdot|||)$ be as assumed in Theorem 1.1. Then $\Delta|_{\mathfrak{S}}$ is Hermitian in $\mathcal{L}(\mathcal{L}(Y,))$ if and only if $A - \lambda I \in \mathcal{L}(X)$ and $B - \lambda I \in \mathcal{L}(Y)$ are Hermitian for some scalar λ .*

Kyle [8] has proved that when $X = Y$ is a Banach space and when $A = B$, Δ is normal if and only if A is normal in $\mathcal{L}(X)$. We shall extend this result under various situations. The statement for the most general situation is as follows.

THEOREM 2.2. *Let $A \in \mathcal{L}(X)$ and $B \in \mathcal{L}(Y)$ be of the forms $A = H + iK$ and $B = M + iN$, where H, K, M and N are Hermitian operators. Suppose that $(\mathfrak{S}, |||\cdot|||)$ satisfies $H\mathfrak{S} \subset \mathfrak{S}$, $K\mathfrak{S} \subset \mathfrak{S}$, $\mathfrak{S}M \subset \mathfrak{S}$, $\mathfrak{S}N \subset \mathfrak{S}$ and conditions (2), (3), (4). Then $\Delta|_{\Sigma}$ is normal if and only if both A and B are normal.*

PROOF. $\Delta_{A,B}$ can be written as $\Delta_{H,M} + i\Delta_{K,N}$, where $\Delta_{H,M}$ and $\Delta_{K,N}$ and their restrictions to \mathfrak{S} are Hermitian, by Corollary 2.1. Now, from the easily verified identity

$$\Delta_{H,M}\Delta_{K,N} - \Delta_{K,N}\Delta_{H,M} = \Delta_{HK-KH, MN-NM}$$

we see that $\Delta|_{\mathfrak{S}}$ is normal if and only if $(HK - KH)T = T(MN - NM)$ for all T in \mathfrak{S} . Since \mathfrak{S} contains all finite rank operators, the latter condition is, by Corollary 1.4, equivalent to that $HK - KH = \lambda I_{\mathcal{L}(X)}$ and $MN - NM = \lambda I_{\mathcal{L}(Y)}$ for some scalar λ . But this is possible only when $\lambda = 0$, that is, A and B are normal (see [10, page 332]).

It follows that for $\Delta|_{\mathfrak{S}}$ to be normal it is sufficient that A and B are normal. That this is also necessary in case X and Y are Hilbert spaces is already contained in the above theorem.

COROLLARY 2.3. *Let H_1 and H_2 be Hilbert spaces. For $A \in \mathcal{L}(H_1)$ and $B \in \mathcal{L}(H_2)$ let $\mathfrak{S} \subset \mathcal{L}(H_2, H_1)$ be a subspace satisfying conditions (1)–(4) (for example, $C_p(H_2, H_1)$ with norm $\|\cdot\|_p$ or operator norm $\|\cdot\|$). Then $\Delta|_{\mathfrak{S}}$ is normal if and only if both A and B are normal.*

REMARK. When A and B are normal operators on a Hilbert space \mathbf{H} , Δ and $\Delta|_{C_2(\mathbf{H})}$ become normal operators on the Banach space $\mathcal{L}(\mathbf{H})$ and the Hilbert space $(C_2(\mathbf{H}), \|\cdot\|_2)$, respectively. It follows (see [6, Theorem A] or [1]) that the null space $N(\Delta)$ is orthogonal to the range $R(\Delta)$ of Δ . Hence we have

$$R(\Delta)^- \oplus N(\Delta) \supset (R(\Delta|_{C_2(\mathbf{H})})^- \oplus N(\Delta|_{C_2(\mathbf{H})}))^- = C_2(\mathbf{H})^-,$$

where the superscripts “ $-$ ” and “ $=$ ” denote the closure relative to $\|\cdot\|$ and $\|\cdot\|_2$, respectively. Thus $R(\Delta)^- \oplus N(\Delta)$ contains all compact operators while it is in general strictly less than $\mathcal{L}(\mathbf{H})$ [1].

Since a general operator on a normed linear space is not necessarily of the form $H + iK$, with H and K Hermitian, it is not known from Theorem 2.2 whether a normal $\Delta_{A,B} | \mathfrak{S}$ ($A \in \mathcal{L}(\mathbf{X}), B \in \mathcal{L}(\mathbf{Y})$) must be made of two normal A and B . But, at least when \mathbf{X} is equal to \mathbf{Y} and when $\mathfrak{S} = \mathcal{L}(\mathbf{X})$, this is true, as is shown by the following extension of Kyle’s result.

THEOREM 2.4. *Let A and B be operators on a normed linear space X . Then $\Delta_{A,B}$ is normal in $\mathcal{L}(\mathcal{L}(\mathbf{X}))$ if and only if both A and B are normal in $\mathcal{L}(\mathbf{X})$.*

This will follow from Theorem 2.2 (with $\mathfrak{S} = \mathcal{L}(\mathbf{X})$) and the next

LEMMA 2.5. *$\Delta_{A,B} = \Phi + i\Psi$ for some Hermitian operators Φ and Ψ in $\mathcal{L}(\mathcal{L}(\mathbf{X}))$ if and only if $A = H + iK$ and $B = M + iN$ for some Hermitian operators H, K, M and N in $\mathcal{L}(\mathbf{X})$.*

PROOF. Suppose $\Delta = \Phi + i\Psi$ where Φ and Ψ are Hermitian. Fix a pair (x_0, f) in $\pi(\mathbf{X})$ and define operators H_1, K_1, M_1 and N_1 by $H_1x := (\Phi(x \otimes f))x_0$, $K_1x := (\Psi(x \otimes f))x_0$ ($x \in X$), $M_1 := H_1 - \Phi(I)$ and $N_1 := K_1 - \Psi(I)$, respectively, where I is the identity operator on \mathbf{X} .

We first show that these operators are Hermitian. To show that H_1 is Hermitian, we will prove that $g(H_1x)$ is real for any pair (x, g) in $\pi(\mathbf{X})$. Indeed, for a fixed (x, g) in $\pi(\mathbf{X})$ there corresponds the linear functional $G: T \rightarrow g(Tx_0)$ ($T \in \mathcal{L}(\mathbf{X})$) on $\mathcal{L}(\mathbf{X})$ which satisfies: $\|G\| = G(x \otimes f) = \|x \otimes f\| = 1$, that is, $(x \otimes f, G) \in \pi(\Lambda(\mathbf{X}))$. This implies that

$$g(H_1x) = g((\Phi(x \otimes f))x_0) = G(\Phi(x \otimes f)) \in W(\Phi; \mathcal{L}(\mathcal{L}(\mathbf{X}))) \subset R.$$

By a similar way one can show that K_1 is also Hermitian. To claim that $\Phi(I)$ is Hermitian we observe that $F(\Phi(I))$ belongs to $W(\Phi, \mathcal{L}(\mathcal{L}(\mathbf{X})))$ for every (I, F) in $\pi(\mathcal{L}(\mathbf{X}))$, or equivalently, $V(\Phi(I); \mathcal{L}(\mathbf{X})) \subset (\Phi; \mathcal{L}(\mathcal{L}(\mathbf{X}))) \subset R$. Similarly, $\Psi(I)$ is Hermitian.

Now we have, for $D \in \mathcal{L}(\mathbf{X})$ and $x \in \mathbf{X}$,

$$\begin{aligned} (\Delta_{H_1, M_1}(D))x &= (H_1D - DM_1)x = H_1Dx - DH_1x + D\Phi(I)x \\ &= (\Phi(Dx \otimes f))_{x_0} - D(\Phi(x \otimes f))_{x_0} + D\Phi(I)x, \end{aligned}$$

and similarly

$$(\Delta_{K_1, N_1}(D))x = (\Psi(Dx \otimes f))_{x_0} - D(\Psi(x \otimes f))_{x_0} + D\Psi(I)x.$$

Thus

$$\begin{aligned} ((\Delta_{H_1, M_1} + i\Delta_{K_1, N_1})D)x &= (\Delta_{A, B}(Dx \otimes f))_{x_0} - D(\Delta_{A, B}(x \otimes f))_{x_0} + D(A - B)x \\ &= A(Dx \otimes f)_{x_0} - (Dx \otimes f)Bx_0 \\ &\quad - D(A(x \otimes f) - (x \otimes f)B)x_0 + D(A - B)x \\ &= ADx - DAx + D(A - B)x \\ &= \Delta_{A, B}(D)x. \end{aligned}$$

That is, $(A - H_1 - iK_1)D = D(B - M_1 - iN_1)$ holds for every D in $\mathcal{L}(\mathbf{X})$. It follows from Corollary 1.4 that $A = H_1 + iK_1 + \lambda I$ and $B = M_1 + iN_1 + \lambda I$ for some scalar λ . Now we can take $H = H_1 + (\operatorname{Re} \lambda)I$, $K = K_1 + (\operatorname{Im} \lambda)I$, $M = M_1 + (\operatorname{Re} \lambda)I$ and $N = N_1 + (\operatorname{Im} \lambda)I$ as the desired Hermitian operators.

So far, the question about when $\Delta_{A, B}|_{\mathfrak{S}}$ is Hermitian in $\mathcal{L}(\mathfrak{S})$ ($\mathfrak{S} \subset \mathcal{L}(\mathbf{Y}, \mathbf{X})$) has been answered by Corollary 2.1, and the question about when $\Delta|_{\mathfrak{S}}$ is normal has been answered by Theorem 2.2 for special operators A and B on normed linear spaces \mathbf{X} and \mathbf{Y} , by Corollary 2.3 for the case where \mathbf{X} and \mathbf{Y} are Hilbert spaces, and by Theorem 2.4 for the case where \mathbf{X} is the same normed linear space as \mathbf{Y} and \mathfrak{S} is $\mathcal{L}(\mathbf{X})$. But the latter question for the more general case where $\mathbf{X} \neq \mathbf{Y}$ or where $\mathbf{X} = \mathbf{Y}$ and $\mathfrak{S} \neq \mathcal{L}(\mathbf{X})$ remain unanswered. It is unknown whether there exist nonnormal operators A, B such that $\Delta_{A, B}|_{\mathfrak{S}}$ is normal.

On the other hand, when $\mathbf{X} = \mathbf{Y} = \mathbf{H}$ is a Hilbert space, one can deduce a stronger result than Corollaries 2.1 and 2.3. Indeed, a result of Sinclair [12, page 213] states that a Hermitian operator on a C^* -algebra (with identity) is the sum of a left multiplication by a Hermitian element in the algebra and a $*$ -derivation, and a result of Kadison [7] and Sakai [11] asserts that every derivation of a C^* -algebra acting on H is spatial (that is, of the form $\Delta_{A, A}$, with A and element in the weak operator closure of the algebra). These facts together with Corollary 2.1 imply that an operator on a C^* -algebra \mathfrak{S} in $\mathcal{L}(\mathbf{H})$ is Hermitian if and only if it is of the form $\Delta_{H, M}|_{\mathfrak{S}}$ for some Hermitian operators H and M (in the weak operator closure of \mathfrak{S}). Recently, Sourour [13] has proved the same assertion for the case where \mathfrak{S} is a minimal norm ideal (including the $C_p(\mathbf{H})$ ideals, $p \neq 2$). Thus every

normal operator on such \mathfrak{S} has to be of the form $\Delta_{A,B}|\mathfrak{S}$, with $A = H + iK$ and $B = M + iN$ for some Hermitian H, K, M and N . Using Theorem 2.2 we obtain the following

THEOREM 2.6. *Let \mathfrak{S} be a minimal norm ideal or a C^* -algebra in $\mathfrak{L}(\mathbf{H})$, which contains all finite rank operators. Then an operator U on \mathfrak{S} is Hermitian (resp. normal) in $\mathfrak{L}(\mathfrak{S})$ if and only if $U = \Delta_{A,B}|\mathfrak{S}$ for some Hermitian (resp. normal) operators A and B .*

3. Berberian's theorem

As another application of Theorem 1.1, we shall derive an extension of the Fuglede-Putnam theorem to hyponormal operators A and B^* on Hilbert spaces \mathbf{H}_1 and \mathbf{H}_2 , respectively. It is also a slight extension of a theorem of Berberian [3] who proved in a different way the special case where $\mathbf{H}_1 = \mathbf{H}_2$.

A natural and consistent definition for a hyponormal operator A on a normed linear space \mathbf{X} is that it can be written as $A = H + iK$ for some Hermitian operators H and K such that $A^*A - AA^* = 2i(HK - KH)$ is positive (that is, has nonnegative numerical range). Suppose $B^* = M - iN$ is a hyponormal operator on another normed linear space \mathbf{Y} . Then the operator $\Delta_{A,B}|\mathfrak{S}$ is also hyponormal. Indeed, from the easily verified identity:

$$\begin{aligned} \Delta^*\Delta - \Delta\Delta^* &= 2i(\Delta_{H,M}\Delta_{K,N} - \Delta_{K,N}\Delta_{H,M}) = 2i\Delta_{HK-KH, MN-NM} \\ &= \Delta_{A^*A-AA^*, B^*B-BB^*} \end{aligned}$$

we see that the numerical range of $(\Delta|\mathfrak{S})^*(\Delta|\mathfrak{S}) - (\Delta|\mathfrak{S})(\Delta|\mathfrak{S})^*$, as the sum of the numerical ranges of the two positive operators $A^*A - AA^*$ and $BB^* - B^*B$ is nonnegative.

In particular, if $\mathbf{X} = \mathbf{H}_1$ and $\mathbf{Y} = \mathbf{H}_2$ are two Hilbert spaces, then $\Delta|C_2(\mathbf{H}_2, \mathbf{H}_1)$ is a hyponormal operator on the Hilbert space $(C_2(\mathbf{H}_2, \mathbf{H}_1), \|\cdot\|_2)$. Hence we have $\|\Delta T\|_2 \geq \|\Delta^*T\|_2$ for all T in $C_2(\mathbf{H}_2, \mathbf{H}_1)$. Since $(\Delta_{A,B})^* = \Delta_{A^*,B^*}$, we have proved the following

THEOREM 3.1. *Let A and B^* be hyponormal operators on the Hilbert spaces \mathbf{H}_1 and \mathbf{H}_2 , respectively. If T is a Hilbert-Schmidt operator from \mathbf{H}_2 to \mathbf{H}_1 such that $AT = TB$, then $A^*T = TB^*$.*

References

- [1] J. Anderson and C. Foias, 'Properties which normal operators share with normal derivations and related operators', *Pacific J. Math.* **61** (1975), 313–325.
- [2] J. P. Aubin, *Applied functional analysis* (Wiley-Interscience, New York, 1979).
- [3] S. K. Berberian, 'Extensions of a theorem of Fuglede and Putnam', *Proc. Amer. Math. Soc.* **71** (1978), 113–114.
- [4] F. F. Bonsall and J. Duncan, *Numerical ranges of operators on normed spaces and of elements of normed algebras* (London Math. Soc. Lecture Note Series 2, Cambridge Univ. Press, 1971).
- [5] L. Fialkow, 'A note on the operator $X \rightarrow AX - XB$ ', *Trans. Amer. Math. Soc.* **243** (1978), 147–168.
- [6] C. K. Fong, 'Normal operators on Banach spaces', *Glasgow Math. J.* **20** (1979), 163–168.
- [7] R. V. Kadison, 'Derivations of operator algebras', *Ann. of Math.* **83** (1966), 280–293.
- [8] J. Kyle, 'Numerical ranges of derivations', *Proc. Edinburgh Math. Soc.* **21** (1978), 33–39.
- [9] A. Pietsch, *Operator ideals* (North-Holland Mathematical library vol. 20, 1980).
- [10] W. Rudin, *Functional analysis* (McGraw-Hill, New York, 1973).
- [11] S. Sakai, 'Derivations of W^* -algebras', *Ann. of Math.* **83** (1966), 273–279.
- [12] A. M. Sinclair, 'Jordan homomorphisms and derivations on semi-simple Banach algebras', *Proc. Amer. Math. Soc.* **24** (1970), 209–214.
- [13] A. R. Sourour, 'Isometries of norm ideals of compact operators', preprint.

Department of Mathematics
National Central University
Chung-Li, Taiwan 320
Republic of China