

SUBSETS CHARACTERIZING THE CLOSURE OF THE NUMERICAL RANGE

K. C. DAS, S. MAJUMDAR and BRAILEY SIMS

(Received 28 May 1984)

Communicated by J. H. Chabrowski

Abstract

For an operator on a Hilbert space, points in the *closure* of its numerical range are characterized as either extreme, non-extreme boundary, or interior in terms of various associated sets of bounded sequences of vectors. These generalize similar results due to Embry, for points in the numerical range.

1980 *Mathematics subject classification* (*Amer. Math. Soc.*): 47 A 12

1. Introduction

Let T be an operator (that is, a bounded linear transformation) on a complex Hilbert space H with inner product $\langle \cdot, \cdot \rangle$ and associated norm $\|\cdot\|$. It is well known that the numerical range

$$W(T) = \{ \langle Tx, x \rangle : \|x\| = 1, x \in H \}$$

is a convex subset of the complex plane. Denote the closure of $W(T)$ by $W(T)^-$. Theorem 1 of M. R. Embry (1970) characterizes every point z of $W(T)$ as either an extreme point, a non-extreme boundary point or an interior point in terms of the subset $M_z(T)$ and its linear span, where

$$M_z(T) = \{ x \in H : \langle Tx, x \rangle - z\|x\|^2 = 0 \} \quad (z \in W(T)).$$

This theorem, though very interesting, does not characterize the unattained boundary points of the numerical range. In this note we attempt to fill this gap by

a generalization which can be applied to every point of $W(T)^-$. For any $z \in W(T)^-$, let

$$N_z(T) = \{(x_n) \in l_\infty(H) : \langle Tx_n, x_n \rangle - z\|x_n\|^2 \rightarrow 0\},$$

$$N'_z(T) = \{(x_n) \in l_\infty(H) : \langle Tx_n, x_n \rangle / \|x_n\|^2 \rightarrow z\},$$

$$N^L(T) = \bigcup_z \{N_z(T) : z \in L \cap W(T)^-\}$$

and

$$N_L(T) = \{(x_n) \in l_\infty(H) : \inf_{z \in L} |\langle Tx_n, x_n \rangle - z\|x_n\|^2| \rightarrow 0\}$$

where $l_\infty(H)$ is the set of all bounded sequences of vectors from H and L is a line of support for $W(T)^-$. Let $\gamma N_z(T)$ be the linear span of $N_z(T)$. Since $N_z(T)$ is homogeneous, $\gamma N_z(T) = N_z(T) + N_z(T)$. It is readily seen that $N_L(T)$ is a subspace (Majumdar and Sims (to appear)).

2. Basic lemmas

In order to establish our characterization for points of $W(T)^-$ we need the following two lemmas. The first, stated without proof, is an easy corollary to Lemma 3 of Majumdar and Sims (to appear).

LEMMA 1. *If b is an extreme point of $W(T)^-$ and L is a line of support for $W(T)$ passing through b , then $\lim \langle (T - b)x_n, y_n \rangle = 0$ and $\lim \langle (T - b)y_n, x_n \rangle = 0$ for all $(x_n) \in N_b(T)$ and $(y_n) \in N_L(T)$.*

LEMMA 2. *Let z be in the interior of a line segment lying in $W(T)^-$ with end points a and b . Then $N'_a(T) \subset \gamma N_z(T)$.*

PROOF. Without loss of generality we may take $a = 1$, $b = 0$ and $(x_n) \in N'_1(T)$ to have $\|x_n\| = 1$. Let $(y_n) \in N_0(T)$ be such that $\|y_n\| = 1$ and $\operatorname{Re} \langle \operatorname{Im} Tx_n, y_n \rangle = 0$. For any bounded sequence (r_n) , let $h_n = r_n x_n + y_n$; then we have $\langle \operatorname{Im} Th_n, h_n \rangle \rightarrow 0$. We show the existence of two such distinct sequences (r_n) for which

$$(1) \quad \langle \operatorname{Re} Th_n, h_n \rangle - z\|h_n\|^2 = 0$$

for all sufficiently large n . The equations in r_n given by (1) are equivalent to

$$r_n^2(1 - z + \epsilon_n) + 2r_n \operatorname{Re} \langle (\operatorname{Re} T - z)x_n, y_n \rangle + (\epsilon'_n - z) = 0$$

where $\epsilon_n = \langle \operatorname{Re} Tx_n, x_n \rangle - 1$ and $\epsilon'_n = \langle \operatorname{Re} Ty_n, y_n \rangle$, both of which tend to zero. Thus the equations in (1) are of the form $A_n r_n^2 + B_n r_n + C_n = 0$ where A_n, B_n, C_n are real numbers independent of r_n .

Let $D_n = B_n^2 - 4A_nC_n$, then

$$D_n = 4[\operatorname{Re}\langle \operatorname{Re} T - z, x_n, y_n \rangle]^2 + 4z(1 - z) + \delta_n$$

where $\delta_n \rightarrow 0$. Hence there are positive constants α, β such that for all sufficiently large n , $\alpha \leq A_n$, $D_n \leq \beta$ and $|B_n| \leq \beta$. This shows the existence of two distinct sequences solving (1) both of which are bounded and whose differences $d_n = \sqrt{D_n}/A_n$ are eventually bounded away from zero. Thus we have for both these sequences that $h_n \in N_z(T)$. Subtraction and the fact that d_n is uniformly bounded away from zero gives $(x_n) \in \gamma N_z(T)$.

REMARK. A simplified version of the above argument applied to a pair of points a, b lying in a line segment in $W(T)$ shows the existence of a real number r and a vector y such that $a = \langle Tx, x \rangle$, $b = \langle Ty, y \rangle$, $\|x\| = \|y\| = 1$ and $\langle T(rx + y), rx + y \rangle / \|rx + y\|^2 = ta + (1 - t)b$, $0 < t < 1$, yielding the convexity of $W(T)$. In contrast with the proof of convexity given by Halmos (1967), this argument gives two explicit values of r .

3. Characterization of $W(T)^-$

THEOREM 3. *Every element z of $W(T)^-$ can be characterized as follows.*

- (i) z is an extreme point of $W(T)^-$ if and only if $N_z(T)$ is a subspace.
- (ii) If z is a nonextreme boundary point of $W(T)^-$ and L the line of support for $W(T)$ passing through z , then (a) $\gamma N_z(T) = N_L(T) + N_z(T)$ and (b) $N_L(T) = l_\infty(H)$ if and only if $W(T)^- \subset L$.
- (iii) If $W(T)^-$ is not a straight line segment, then z is an interior point of $W(T)^-$ if and only if $N'_a(T) \subset \gamma N_z(T)$ for all $z \in W(T)^-$.

PROOF. (i) See Das and Craven (1983) and also Majumdar and Sims (to appear). Also note that the result $N_z(T)$ is a subspace when z is an extreme point of $W(T)^-$ can be deduced as a corollary to Lemma 1. Homogeneity being obvious, we prove linearity. Let $(x_n^{(1)}), (x_n^{(2)}) \in N_z(T)$. Thus $(x_n^{(1)}), (x_n^{(2)}) \in N_L(T)$ where L is a line of support for $W(T)$ passing through z . But $N_L(T)$ is a subspace. So $(x_n^{(1)} + x_n^{(2)}) \in N_L(T)$. Now since $(x_n^{(i)}) \in N_z(T)$, $i = 1, 2$ and $(x_n^{(1)} + x_n^{(2)}) \in N_L(T)$, by Lemma 1 we have $\lim \langle (T - z)x_n^{(1)}, x_n^{(1)} + x_n^{(2)} \rangle = 0$ for $i = 1, 2$ and hence $\lim \langle (T - z)(x_n^{(1)} + x_n^{(2)}), x_n^{(1)} + x_n^{(2)} \rangle = 0$ as required.

(ii) (a) We first show $N_a(T) \subset \gamma N_z(T)$ for each $a \in L \cap W(T)^-$. Without loss of generality we may take L as the real axis and $\operatorname{Im} W(T) \geq 0$. Let $(x_n) \in N_a(T)$ and $(y_n) \in N_b(T)$, $\|y_n\| = 1$ where $b \in L$ is the extreme point of $W(T)^-$ such that $(a - z)/(z - b) \geq 0$. Then (y_n) can be chosen so that $\operatorname{Re}\langle y_n, x_n \rangle = 0$ and

Lemma 1 gives $\operatorname{Re}\langle Ty_n, x_n \rangle \rightarrow 0$. Also $\operatorname{Im} W(T) \geq 0$ implies $\operatorname{Im} Ty_n \rightarrow 0$. Let $r_n = [(a - z)/(z - b)]^{1/2} \|x_n\|$. Then easy calculations show that with our assumptions $\langle T(x_n \pm r_n y_n), x_n \pm r_n y_n \rangle - z \|x_n \pm r_n y_n\|^2 \rightarrow 0$. That is $(x_n \pm r_n y_n) \in N_z(T)$. As in the proof of Lemma 2, adding these two sequences and using the homogeneity of $N_z(T)$ we have $(x_n) \in \gamma N_z(T)$. Thus $N_a(T) \subset \gamma N_z(T)$ for all $a \in L \cap W(T)^-$ and so we have $N^L(T) \subset \gamma N_z(T)$. Since $N_z(T) \subset N^L(T) \subset \gamma N_z(T)$, by taking the vector sum of $N_z(T)$ with each of these subsets we obtain $\gamma N_z(T) = N^L(T) + N_z(T)$.

(b) As before, if we take L as the real axis, we have $N_L(T) = \{(x_n) \in l_\infty(H) : \operatorname{Im}\langle Tx_n, x_n \rangle \rightarrow 0\}$. Now if $W(T)^- \subset L$, $(x_n) \in l_\infty(H)$ implies $\operatorname{Im}\langle Tx_n, x_n \rangle = 0$ and so $(x_n) \in N_L(T)$. Hence $N_L(T) = l_\infty(H)$. Conversely if $W(T)^-$ is not a subset of L , there exists $(x_n) \in l_\infty(H)$, $\|x_n\| = 1$ such that $\operatorname{Im}\langle Tx_n, x_n \rangle$ does not tend to zero, or equivalently, $(x_n) \notin N_L(T)$. Hence $N_L(T) \neq l_\infty(H)$.

(iii) If z is an interior point of $W(T)^-$, by Lemma 2, $N'_a(T) \subset \gamma N_z(T)$ whenever $a \in W(T)^-$. On the other hand, if z is a boundary point of $W(T)^-$, without loss of generality we may take L , the line of support for $W(T)$ passing through z , as the real axis, in which case, $N_L(T) = \{(x_n) \in l_\infty(H) : \operatorname{Im}\langle Tx_n, x_n \rangle \rightarrow 0\}$. Thus $\gamma N_z(T) \subset N_L(T)$ since $N_L(T)$ is a subspace, but as $W(T)^-$ does not lie in L , there exists an $a \in W(T)$ such that $\operatorname{Im} a \neq 0$. Hence $N'_a(T)$ is not a subset of $\gamma N_z(T)$.

References

- K. C. Das and B. D. Craven (1983), 'Linearity and weak convergence on the boundary of numerical range', *J. Austral. Math. Soc. Ser. A* **35**, 221–226.
- M. R. Embry (1970), 'The numerical range of an operator', *Pacific J. Math.* **32**, 647–650.
- M. R. Embry (1975), 'Orthogonality and the numerical range', *J. Math. Soc. Japan* **27**, 405–411.
- P. R. Halmos (1967), *Hilbert space problem book* (Van Nostrand, New York).
- S. Majumdar and B. Sims (to appear), 'Subspaces associated with boundary points of the numerical range', *J. Austral. Math. Soc. Ser. A*.
- J. G. Stampfil (1966), 'Extreme points of the numerical range of a hyponormal operator', *Michigan Math. J.* **13**, 87–89.

K. C. Das
Indian Institute of Technology
Karagpur 721302
India

S. Majumdar and Brailey Sims
University of New England
Armidale, N.S.W. 2351
Australia