

ON THE SPECTRUM OF C_1 AS AN OPERATOR ON bv_0

J. I. OKUTOYI

(Received 28 March 1988; revised 3 March 1989)

Communicated by S. Yamamuro

Abstract

In 1985 John Reade determined the spectrum of C_1 regarded as an operator on the space c_0 of all null sequences normed by $\|x\| = \sup_{n \geq 0} |x_n|$. It is the purpose of this paper to determine the spectrum of C_1 regarded as an operator on the space bv_0 of all sequences x such that $x_k \rightarrow 0$ as $k \rightarrow \infty$ and $\|x\| = \sum_{k=0}^{\infty} |x_{k+1} - x_k| < \infty$.

1980 *Mathematics subject classification* (*Amer. Math. Soc.*) (1985 *Revision*): 40 F 05, 47 A 10.

NOTATION. s ; c_0 ; l_1 ; bv_0 ; bs ; T^* ; $X^*B(X)$; A^t ; $\sigma(T)$; $O(1)$; $o(1)$; \asymp ; $\text{Re}(z)$; will denote the set of all sequences; convergent to zero sequences, that is, null sequences; sequences such that $\sum_{k=0}^{\infty} |x_k| < \infty$; sequences such that $\sum_{k=0}^{\infty} |x_{k+1} - x_k| < \infty$; bounded series, that is, sequences x such that $\sup_{n \geq 0} |\sum_{k=0}^n x_k| < \infty$; the adjoint operator of T ; the space of all continuous linear functionals on X , that is, the continuous dual of X ; the linear space of all bounded linear operators, say, T on X into itself; the transposed matrix of A ; the spectrum of T ; capital order, that is, $x_n = O(1)$ if there exists $M \in \mathbf{R}^+$ such that $|x_n| \leq M$ for all n ; small order, that is, $x_n = o(1)$ as $n \rightarrow \infty$, that is, $\lim_{n \rightarrow \infty} x_n = 0$; lies between two positive constant multiples, for example $a_n \asymp b_n$ means that there exist $m, M \in \mathbf{R}^+$ such that $mb_n \leq a_n \leq Mb_n$; the real part of the complex number z , respectively.

1. Introduction

In his 1985 paper Reade considers the operator which converts a sequence $(x_n)_0^\infty$ into its sequence of averages

$$\left(\frac{x_0 + x_1 + \cdots + x_n}{n+1} \right)_{n=0}^\infty.$$

He shows it is a bounded operator on c_0 . We shall denote this operator by $C_1 = (C, 1)$ and call it the Cesàro operator. It can be represented by the matrix

$$\begin{pmatrix} 1 & 0 & 0 & 0 & \cdots \\ \frac{1}{2} & \frac{1}{2} & 0 & 0 & \cdots \\ \frac{1}{3} & \frac{1}{3} & \frac{1}{3} & 0 & \cdots \\ \cdots & \cdots & \cdots & \cdots & \cdots \end{pmatrix}.$$

The key to determining the spectrum $\sigma(C_1)$ of a bounded linear operator $C_1: bv_0 \rightarrow bv_0$ on a Banach space bv_0 is the determination of all eigenvalues of $C_1^* \in B(bv_0^*)$, that is, the determination of all $\lambda \in \mathbb{C}$ such that $(C_1 - \lambda I)^{-1} \in B(bv_0)$.

1.1 THEOREM. *Let $T \in B(X)$, where X is any Banach space. Then the spectrum of T^* is identical with the spectrum of T . Furthermore, $R_\lambda(T^*) = (R_\lambda(T))^*$ for $\lambda \in \rho(T) = \rho(T^*)$, where $R_\lambda(T) = (T - \lambda I)^{-1}$ and $\rho(T) = \{\lambda \in \mathbb{C}: (T - \lambda I)^{-1} \text{ exists}\}$.*

PROOF. See [2, page 568] and [3, page 71].

1.2. LEMMA. $C_1: bv_0 \rightarrow bv_0$ and $C_1 \in B(bv_0)$ with $\|C_1\|_{bv_0} = 1$.

PROOF. Since $C_1: bv_0 \rightarrow bv_0$, write $y_n = C_1 x$ and define $x_n = a_0 + a_1 + \cdots + a_n$. Then $y_n \rightarrow 0$ as $n \rightarrow \infty$ and

$$\begin{aligned} \sum_{n=0}^{\infty} |y_n - y_{n+1}| &= \sum_{n=0}^{\infty} \frac{1}{(n+1)(n+2)} \left| \sum_{\nu=1}^{n+1} \nu a_\nu \right| \\ &\leq \sum_{n=0}^{\infty} \frac{1}{(n+1)(n+2)} \sum_{\nu=1}^n \nu |a_\nu| \\ &= \sum_{\nu=1}^{\infty} \nu |a_\nu| \sum_{n=\nu-1}^{\infty} \frac{1}{(n+1)(n+2)} \\ &\leq \sum_{\nu=1}^{\infty} |a_\nu| = \sum_{\nu=0}^{\infty} |x_\nu - x_{\nu-1}|. \end{aligned}$$

Direct manipulation gives

$$\|C_1\|_{(bv_0, bv_0)} = \sup\{1, 0, 0, \dots\} = 1.$$

Clearly $\lim_{n \rightarrow \infty} \frac{1}{1+n} = 0$ and hence $C_1 \in B(bv_0)$.

1.3. **LEMMA.** *Each bounded linear operator $T: X \rightarrow Y$, where $X = c_0, l_p$ and $Y = c_0, l_p$ ($1 \leq p < \infty$), l_∞ (where l_p denotes sequences x such that $\sum_{k=0}^\infty |x_k|^p < \infty$ and l_∞ denotes bounded sequences) determines and is determined by an infinite matrix of complex numbers.*

PROOF. See Taylor [13, pages 221–223].

1.4. **LEMMA.** *Let $C_1: bv_0 \rightarrow bv_0$. Then $C_1^*: bv_0^* \rightarrow bv_0^*$ is given by C_1^* and $C_1^t \in B(bs)$.*

PROOF. Since bv_0 has AK and bv_0^* is isomorphic to bs under the map $h: bv_0^* \rightarrow bs$, $h(f) = (t_0, t_1, t_2, \dots)$, where $t_n = f(\delta^n)$, $n \geq 0$, $f \in bv_0^*$, we have (see Lemma 1.3)

$$C_1^* = C_1^t = \begin{pmatrix} 1 & \frac{1}{2} & \frac{1}{3} & \frac{1}{4} & \dots \\ 0 & \frac{1}{2} & \frac{1}{3} & \frac{1}{4} & \dots \\ 0 & 0 & \frac{1}{3} & \frac{1}{4} & \dots \\ \dots & \dots & \dots & \dots & \dots \end{pmatrix}.$$

But for any operator T on a normed linear space X , $\|T\|_X = \|T^*\|_{X^*}$ (see [2, page 478], [3, page 54] and [7, page 232]), so

$$\|C_1\|_{bv_0} = \|C_1^*\|_{bv_0^*} = \|C_1^t\|_{bs} = 1.$$

Thus $C_1^* \in B(bv_0^*)$, that is, $C_1^t \in B(bs)$ since it is also clear that each column of C_1^t is null (C_1 being a normal matrix).

1.5. **COROLLARY.** $C_1 \in B(bv_0)$ has not eigenvalues.

PROOF. The proof follows from the fact that $C_1 \in B(c_0)$ has no eigenvalues (see [10]) since $bv_0 \subset c_0$.

1.6. **LEMMA.** *Let*

$$Z_n = \prod_{\nu=0}^n \left(1 - \frac{1}{\lambda(\nu+1)}\right), \quad \lambda \neq 0, \lambda \in \mathbb{C}.$$

Then the partial sums of $\sum_{n=1}^\infty Z_n$ are bounded if and only if $\operatorname{Re}(1/\lambda) \geq 1$.

PROOF. When $\lambda = 1$, $Z_n = 0$ for all n and so the partial sums of $\sum_{n=0}^\infty Z_n$ are certainly bounded.

Let C be a constant depending only on λ which may be different at each occurrence and A a non-zero constant. We have that

$$(1.1) \quad \log_e(1-u) = -\sum_{n=1}^{\infty} \frac{1}{n} u^n = -u + O(u^2)$$

uniformly in $|u| \leq \frac{1}{2}$, $u \in \mathbb{C}$. Now given $\lambda \neq 0$ there is ν_0 such that $|\lambda|(\nu+1) > 2$ for $\nu \geq \nu_0$,

$$(1.2) \quad \begin{aligned} \log_e Z_n &= \sum_{\nu=0}^n \log \left(1 - \frac{1}{\lambda(\nu+1)} \right) \\ &= C - \frac{1}{\lambda} \sum_{\nu=\nu_0} \frac{1}{1+\nu} + \sum_{\nu=\nu_0}^n t_{\nu} \end{aligned}$$

where $t_{\nu} = O(1/\nu^2)$, and

$$(1.3) \quad \sum_{\nu=\nu_0}^n t_{\nu} = \sum_{\nu=\nu_0}^{\infty} t_{\nu} - \sum_{\nu=n+1}^{\infty} t_{\nu} = C + O\left(\frac{1}{n}\right)$$

Also

$$(1.4) \quad \sum_{\nu=\nu_0}^n \frac{1}{\nu+1} = C + \log n + O\left(\frac{1}{n}\right)$$

since if $C = \sum_{\nu=0}^n \frac{1}{\nu+1} - \log n$, then

$$C_{n+1} - C_n = \frac{1}{2+n} - \log \left(\frac{n+1}{n} \right) = O\left(\frac{1}{n^2}\right)$$

Therefore

$$\begin{aligned} C_{n+1} &= C + \sum_{\nu=0}^n (C_{\nu+1} - C_{\nu}) \\ &= C + \sum_{\nu=0}^{\infty} (C_{\nu+1} - C_{\nu}) - \sum_{\nu=n+1}^{\infty} (C_{\nu+1} - C_{\nu}), \quad C_0 = 0, \end{aligned}$$

that is,

$$(1.5) \quad C_{n+1} = C - \sum_{\nu=n+1}^{\infty} (C_{\nu+1} - C_{\nu}) = C + O\left(\frac{1}{n}\right)$$

Hence as $n \rightarrow \infty$, $\log Z_n = C - 1/\lambda \log n + O(\frac{1}{n})$ so $Z_n = An^{-1/\lambda}(1 + O(\frac{1}{n})) = An^{-1/\lambda} + O(n^{-\operatorname{Re}(1/\lambda)-1})$. If $\operatorname{Re}(1/\lambda) \geq 1$, $\lambda = 1$, the partial sums of $\sum_{n=1}^{\infty} n^{-1/\lambda}$ are bounded and $\sum_{n=1}^{\infty} n^{-\operatorname{Re}(1/\lambda)-1} < \infty$, so the partial sums of $\sum_{n=1}^{\infty} Z_n$ are bounded. If $0 < \operatorname{Re}(1/\lambda) < 1$ or $\lambda = 1$ then the partial sums

of $\sum_{n=1}^{\infty} n^{-1/\lambda}$ are unbounded, but we still have $\sum_{n=1}^{\infty} n^{-\operatorname{Re}(1/\lambda)-1} < \infty$. If $\operatorname{Re}(1/\lambda) \leq 0$ then

$$(1.6) \quad \sum_{n=1}^N n^{-1/\lambda} \asymp N^{1-1/\lambda} \left(1 - \frac{1}{\lambda}\right)$$

Now

$$\sum_{n=1}^N n^{-\operatorname{Re}(1/\lambda)-1} = \begin{cases} O(N^{-\operatorname{Re}(1/\lambda)}), & \text{if } \operatorname{Re}(1/\lambda) < 0, \\ O(\log N), & \text{if } \operatorname{Re}(1/\lambda) = 0. \end{cases}$$

Using (1.6) we see that the partial sums of $\sum_{n=1}^{\infty} n^{-1/\lambda}$ are unbounded although $\sum_{n=1}^{\infty} n^{-\operatorname{Re}(1/\lambda)-1} < \infty$, and hence we conclude that the partial sums of $\sum_{n=1}^{\infty} Z_n$ are bounded if and only if $\operatorname{Re}(1/\lambda) \geq 1$.

2. Determination of the spectrum of C_1 on bv_0

2.1. THEOREM. *The eigenvalues of $C_1^* \in B(bv_0^*)$, that is, $C_1^t \in B(bs)$, are all $\lambda \in \mathbb{C}$ satisfying $|\lambda - \frac{1}{2}| \leq \frac{1}{2}$.*

PROOF. Suppose $C_1^t x = \lambda x$, $x \in bs$, $x \neq \theta$ where θ is the zero sequence. Then as in Lemma 1.4,

$$C_1^* = C_1^t = \begin{pmatrix} 1 & \frac{1}{2} & \frac{1}{3} & \frac{1}{4} & \cdots & \\ 0 & \frac{1}{2} & \frac{1}{3} & \frac{1}{4} & \cdots & \\ 0 & 0 & \frac{1}{3} & \frac{1}{4} & \cdots & \\ \cdots & \cdots & \cdots & \cdots & \cdots & \end{pmatrix}$$

and solving the system of equations

$$x_0 + \frac{1}{2}x_1 + \frac{1}{3}x_2 + \cdots = x_0$$

$$\frac{1}{2}x_1 + \frac{1}{3}x_2 + \cdots = x_1$$

$$\frac{1}{3}x_2 + \cdots = x_2$$

...

$$\frac{1}{n}x_{n-1} + \frac{1}{n+1}x_n + \cdots = x_{n-1}$$

$$\frac{1}{n+1}x_n + \cdots = x_n$$

...

we obtain

$$\begin{aligned} x_1 &= \left(1 - \frac{1}{\lambda}\right) x_0 \\ x_2 &= \left(1 - \frac{1}{\lambda}\right) \left(1 - \frac{1}{2\lambda}\right) x_0 \\ (2.2) \quad x_3 &= \left(1 - \frac{1}{\lambda}\right) \left(1 - \frac{1}{2\lambda}\right) \left(1 - \frac{1}{3\lambda}\right) x_0. \end{aligned}$$

$$\dots$$

$$x_N = \prod_{n=1}^N \left(1 - \frac{1}{n\lambda}\right) x_0.$$

By Lemma 1.6, $(x_N)_1^\infty \in bs$ if and only if $\operatorname{Re}(1/\lambda) \geq 1$, that is, $|\lambda - \frac{1}{2}| \leq \frac{1}{2}$. Hence the result follows.

2.2. **THEOREM.** Let $C_1: bv_0 \rightarrow bv_0$. Then the spectrum of C_1 is

$$\sigma(C_1) = \{\lambda \in \mathbb{C} : |\lambda - \tfrac{1}{2}| \leq \tfrac{1}{2}\}.$$

2.3. **DEFINITION** (Weighted mean method). The weighted mean method is a matrix $A = (a_{nk})$ with

$$a_{nk} = p_k |P_n|, \quad P_n = \sum_{k=0}^n p_k \neq 0.$$

2.4. **LEMMA.** If $(M, p) = (N, p)$ is a regular (conservative) weighted mean method then $(M, p) = (N, p)$ is absolutely regular (conservative).

(See [1], [14] for further details.)

PROOF. Since (N, p) is a regular (conservative) mean method we have by the Kojima-Schur conditions

$$(2.3) \quad |P_n| \rightarrow \infty \quad \text{as } n \rightarrow \infty,$$

where $P_n = \sum_{\nu=0}^n p_\nu$ and

$$(2.4) \quad P_n^* = \sum_{\nu=0}^n |p_\nu| = O(P_n).$$

We need to prove that (N, p) is absolutely regular (conservative), that is, that

$$(2.5) \quad P_{k-1} \sum_{n=k}^{\infty} \left| \frac{1}{P_n} - \frac{1}{P_{n-1}} \right| \leq M.$$

Let $P_n^* = \sum_{\nu=0}^n |p_\nu|$. Then (2.4) becomes $P_n^* \leq K|P_n|$ for all $n \geq 1$ (K some constant). Thus

$$\begin{aligned} |P_{k-1}| \sum_{n=k}^{\infty} \frac{|p_n|}{|P_n| |P_{n-1}|} &\leq |P_{k-1}| \sum_{n=k}^{\infty} \frac{|P_n| K^2}{P_n^* \cdot P_{n-1}^*} \\ &\leq K^2 |P_{k-1}| \sum_{n=k}^{\infty} \left(\frac{1}{P_{n-1}^*} - \frac{1}{P_n^*} \right). \end{aligned}$$

Since $|P_n| \rightarrow \infty$ as $n \rightarrow \infty$ by (2.3), we have that $P_n^* \rightarrow \infty$ (since $P_n^* \geq |P_n|$), therefore $\sum_{n=k}^{\infty} (1/P_{n-1}^* - 1/P_n^*) = 1/P_{k-1}^*$ and so (2.5) follows, provided that $|P_{k-1}|/P_{k-1}^* \leq M$ for some M . But $M = 1$ will do and the result follows.

We now prove Theorem 2.2.

PROOF. By virtue of Theorem 2.1 and the fact that $\sigma(C_1) = \sigma(C_1^*)$ (see Theorem 1.1), it is enough to prove that $B = (C_1 - \lambda I)^{-1} \in B(bv_0)$ for all $|\lambda - \frac{1}{2}| > \frac{1}{2}$, that is, that Q is absolutely regular where $B = -I/\lambda - Q/\lambda(\lambda - 1)$ except when λ is the reciprocal of a positive integer, $B = (C - \lambda I)^{-1} = I/\lambda - Q/\lambda(\lambda - 1)$, where $Q = (q_{nk})$, $q_{nk} = A_{k-1}^{-1/\lambda}/A_{n-1}^{1-1/\lambda}$,

$$A_n^\alpha = \binom{n+\alpha}{n} = \frac{(\alpha+n) \cdots (\alpha+1)}{n!}$$

is a Hausdorff matrix (μ, μ_n) ,

$$\mu_n = \frac{1}{\lambda} \left(-1 - \frac{\frac{1}{\lambda}}{(n+1) - \frac{1}{\lambda}} \right).$$

It is also clear that Q is the Hausdorff matrix $(\mu, (1 - \frac{1}{\lambda})/((n+1) - \frac{1}{\lambda}))$. The proof of this is trivial (see Rhoades [11]).

Now Q is a regular Hausdorff transformation when $\operatorname{Re}(1/\lambda) < 1$. To see this we simply check the regularity conditions, namely:

(i) $\lim_{n \rightarrow \infty} q_{nk} = \lim_{n \rightarrow \infty} A_{n-1}^{-1/\lambda}/A_{n-1}^{1-1/\lambda} = 0$ since

$$|q_{nk}| = |A_{k-1}^{-1/\lambda}/A_{n-1}^{1-1/\lambda}| = |A_{k-1}^{-1/\lambda}| \cdot O(n^{\alpha-1})$$

and $\alpha = \operatorname{Re}(1/\lambda) < 1$, whence $q_{nk} \rightarrow 0$ as $n \rightarrow \infty$;

(ii) $\sum_{k=1}^n A_{k-1}^{-1/\lambda} = A_{n-1}^{1-1/\lambda}$, and therefore $\lim_{n \rightarrow \infty} \sum_{k=1}^n q_{nk} = 1$;

(iii) $\sum_{k=1}^n A_{k-1}^{-1/\lambda} = \sum_{k=1}^n O(k^{-\alpha}) = O(n^{1-\alpha}) \asymp O(|A_{n-1}^{1-1/\lambda}|)$ and therefore $\sum_{k=1}^n |A_{k-1}^{-1/\lambda}| = O(|A_{n-1}^{1-1/\lambda}|)$.

It is clear that $Q = (q_{nk})$ is a weighted mean method (matrix) $(N, A_{k-1}^{-1/\lambda})$ with $\sum_{k=1}^n A_{k-1}^{-1/\lambda} = A_{n-1}^{1-1/\lambda}$. Since Q is a weighted mean method and a regular Hausdorff method, theorem 2.2 follows.

Acknowledgement

I would like to thank Professor A. Kuttner and Dr. B. Thorpe, both of the University of Birmingham (U.K.), for their guidance throughout the production of this paper.

References

- [1] D. Borwein, 'On strong and absolute summability', *Proc. Glasgow Math. Assoc.* **4** (1959), 81–83.
- [2] N. Dunford and J. T. Schwartz, *Linear operators; Part I, General theory* (John Wiley and Sons, 1967).
- [3] S. Goldberg, *Unbounded Linear operators-theory and applications*, (McGraw-Hill, 1966).
- [4] G. H. Hardy, *Divergent series*, (Oxford, 1949).
- [5] A. Jakimovski and S. M. Ramanujan, 'A uniform approximation theorem and its application to moment problems', *Math. Z.* **84** (1964), 143–153.
- [6] A. Jakimovski and D. C. Russel, 'Matrix mappings between BK -spaces', *Bull. London Math. Soc.* **4** (1972), 345–353.
- [7] E. Kreyszig, *Introductory functional analysis with applications*, (John Wiley and Sons, 1980).
- [8] G. Leibowitz, 'The Cesàro operators and their generalizations: Examples in infinite dimensional linear analysis.' *Amer. Math. Monthly* **80** (1973), 654–661.
- [9] I. J. Maddox, *Elements of functional analysis*, (Cambridge Univ. Press, 1970).
- [10] J. B. Reade, 'On the spectrum of the Cesàro operator,' *Bull. London Math. Soc.* **17** (1985), 263–267.
- [11] B. E. Rhoades, 'Spectra of some Hausdorff operators,' *Acta Sci. Math. (Szeged)* **32** (1971), 91–100.
- [12] M. Stieglitz and H. Tietz, 'Matrixtransformationen von Folgenräumen. Eine Ergebnisübersicht,' *Math Z.* **154** (1977), 1–16.
- [13] A. E. Taylor and D. C. Lay, *Introduction to functional analysis* (2nd ed., John Wiley and Sons, 1980).

Kenyatta University
P.O. Box 43844
Nairobi
Kenya