

MULTIPLIERS ON WEIGHTED HARDY SPACES OVER LOCALLY COMPACT VILENKIN GROUPS, I

C. W. ONNEWEER and T. S. QUEK

(Received 31 March 1989)

Communicated by I. Raeburn

Abstract

Let G denote a locally compact Vilenkin group with dual group Γ . We give sufficient conditions for a function $\varphi \in L^\infty(\Gamma)$ to be a multiplier from the power-weighted Hardy space $H_\alpha^p(G)$ to itself or the corresponding power-weighted Lebesgue space $L_\alpha^p(G)$, $0 < p \leq 1$, $-1 < \alpha \leq 0$.

1980 *Mathematics subject classification* (Amer. Math. Soc.) (1985 Revision): primary 43 A 22; secondary 43 A 15, 43 A 70.

1. Introduction

In a number of recent papers by T. Kitada [4], [5], [6] and by the present authors [7], [8], [9] various multiplier theorems for spaces of functions or distributions defined on locally compact Vilenkin groups were proved. The spaces considered in these papers were the L^p -spaces with power weights, $1 \leq p < \infty$, the H^p -spaces, $0 < p < 1$, and the power-weighted H^1 spaces. In the present paper we consider multipliers on power-weighted Hardy spaces H_α^p , where $0 < p \leq 1$ and $-1 < \alpha \leq 0$. Our results are of two kinds: the first result, Theorem 4.5, gives a sufficient condition for a function to be a multiplier from H_α^p to the corresponding power-weighted Lebesgue space L_α^p , the second result, Theorem 4.7, deals with multipliers from H_α^p to H_α^p . As a consequence of this last result we prove a multiplier theorem for H_α^p spaces, where the multiplier satisfies a Hörmander-type condition; see Theorem 4.15.

Whereas some of the multiplier theorems in [4] have an analogue for function or distribution spaces on \mathbb{R}^n , for the multiplier theorems presented here no comparable version on \mathbb{R}^n seems to be known.

We now give a brief outline of the paper. In the next section we introduce the necessary definitions and notation. In Section 3 we prove the equivalence of the maximal function characterization of the H_α^p spaces and their characterization in terms of weighted atoms. We also give an interpolation theorem for operators on H^{p_0} spaces and L^{p_1} spaces, $0 < p_0 \leq 1 < p_1 < \infty$. Section 4 is devoted to proofs of our main results and a brief discussion of the sharpness of the second of these results. We conclude that section, and the paper, by deriving the Hörmander-type multiplier theorem for the spaces H_α^p .

2. Definitions and notation

Throughout this paper G will denote a locally compact Abelian group containing a strictly decreasing sequence of open compact subgroups $(G_n)_{n=-\infty}^\infty$ such that

- (i) $\sup\{\text{order}(G_n/G_{n+1}): n \in \mathbb{Z}\} < \infty$,
- (ii) $\bigcup_{n=-\infty}^\infty G_n = G$ and $\bigcap_{n=-\infty}^\infty G_n = \{0\}$.

Such groups are the locally compact analogue of the so-called Vilenkin groups which were first described by N. Ya. Vilenkin in 1947 [13]. Examples of such groups are given in [2, Section 4.1.2]. Additional examples are the additive group of the p -adic numbers and, more general, of a local field, see [11].

Let Γ denote the dual group of G and for each $n \in \mathbb{Z}$ let

$$\Gamma_n = \{\gamma \in \Gamma: \gamma(x) = 1 \text{ for all } x \in G_n\}.$$

We choose Haar measures μ on G and λ on Γ so that $\mu(G_0) = \lambda(\Gamma_0) = 1$. Then $\mu(G_n) = (\lambda(\Gamma_n))^{-1} := (m_n)^{-1}$ for each $n \in \mathbb{Z}$.

There exists a metric d on $G \times G$ defined by $d(x, x) = 0$ and $d(x, y) = (m_n)^{-1}$ if $x - y \in G_n \setminus G_{n+1}$. Then the topology on G determined by the metric d coincides with the original topology on G . For $x \in G$ we set $\|x\| = d(x, 0)$. For each $\alpha \in \mathbb{R}$ we define the function v_α on G by $v_\alpha(x) = \|x\|^\alpha$; the corresponding measure $v_\alpha d\mu = \|x\|^\alpha d\mu$ will also be denoted by $d\mu_\alpha$. We mention here that a simple computation shows that $\mu_\alpha(G_n) \leq C(m_n)^{-(\alpha+1)}$, provided $\alpha > -1$, and that $\mu_\alpha(x + G_n) = (m_j)^{-\alpha}(m_n)^{-1}$ if $x \in G_j \setminus G_{j+1}$ for some $j < n$. Here, like elsewhere, C will denote a constant whose value may change from one occurrence to the next. The Lebesgue spaces on G with respect to the measures $d\mu_\alpha$ will be denoted by $L_\alpha^p(G)$ or

L_α^p , and for $f \in L_\alpha^p$, $0 < p < \infty$ and $\alpha \in \mathbb{R}$ we set

$$\|f\|_{p,\alpha} = \left(\int_G |f(x)|^p d\mu_\alpha(x) \right)^{1/p}.$$

If $\alpha = 0$ we write, as usual, L^p and $\|f\|_p$ instead of L_0^p and $\|f\|_{p,0}$.

As a further generalization of the usual L^p spaces we give here the definition of the Herz spaces on G . We shall use, both here and elsewhere, the notation χ_A for the characteristic function of a set A .

DEFINITION 2.1. Let $0 < p, q \leq \infty$ and $\alpha \in \mathbb{R}$. A measurable function $f: G \rightarrow \mathbb{C}$ belongs to the Herz space $K(\alpha, p, q; G) = K(\alpha, p, q)$ if

$$\|f\|_{K(\alpha,p,q)} := \left(\sum_{l=-\infty}^{\infty} \|(m_l)^{-\alpha} f \chi_{G_l \setminus G_{l+1}}\|_p^q \right)^{1/q} < \infty,$$

with the usual modification if $q = \infty$.

It is easy to see that $K(\alpha/p, p, p) = L_\alpha^p$ for $\alpha \in \mathbb{R}$ and $0 < p < \infty$.

We can also define a metric δ on $\Gamma \times \Gamma$ compatible with the topology on Γ . In this case we have $\|\gamma\| = \delta(\gamma, \gamma_0) = m_n$ if $\gamma \in \Gamma_{n+1} \setminus \Gamma_n$, where $\gamma_0 \in \Gamma$ is defined by $\gamma_0(x) = 1$ for all $x \in G$.

The symbols $^\wedge$ and $^\vee$ will be used to denote the Fourier transform and inverse Fourier transform, respectively. An easy computation shows that

$$(\chi_{G_n})^\wedge = (\lambda(\Gamma_n))^{-1} \chi_{\Gamma_n} = (m_n)^{-1} \chi_{\Gamma_n}$$

and, hence

$$(\chi_{\Gamma_n})^\vee = (\mu(G_n))^{-1} \chi_{G_n} = m_n \chi_{G_n} := \Delta_n.$$

We now briefly review the definition of the spaces of test functions, $S(G)$, and distributions, $S'(G)$; for more details, see [11]. A function $\varphi: G \rightarrow \mathbb{C}$ belongs to $\varphi(G)$ if there exist integers k, l , depending on φ , so that $\text{supp } \varphi \subset G_k$ and φ is constant on the cosets of G_l in G . A sequence $(\varphi_n)_1^\infty$ of functions in $S(G)$ converges to $\varphi \in S(G)$ if there exist $k, l \in \mathbb{Z}$ so that every φ_n and φ has support in G_k and is constant on the cosets of G_l in G and if $\lim_{n \rightarrow \infty} \varphi_n(x) = \varphi(x)$ uniformly on G .

Next, $S'(G)$ is the space of continuous linear functionals on $S(G)$. A sequence $(f_n)_1^\infty$ in $S'(G)$ converges to $f \in S'(G)$ if for all $\varphi \in S(G)$ we have $\lim_{n \rightarrow \infty} \langle f_n, \varphi \rangle = \langle f, \varphi \rangle$.

3. Power-weighted Hardy spaces on G

In [5] Kitada gave a definition for the Hardy spaces $H_\alpha^1(G)$ with respect to the weight functions $v_\alpha(x) = \|x\|^\alpha$, where $-1 < \alpha \leq 0$. In the following

we extend Kitada's definition. If $f \in S'(G)$ we first define its regularization on $G \times \mathbb{Z}$ by $f(x, n) = f_n(x) = f * \Delta_n(x)$. Then f_n is a function on G which is constant on the cosets of G_n in G . Moreover, $\lim_{n \rightarrow \infty} f_n = f$ in $S'(G)$; see [11, Chapter IV]. For $f \in S'(G)$ we define its maximal function f^* by $f^*(x) = \sup_n |f * \Delta_n(x)|$.

DEFINITION 3.1. Let $0 < p < \infty$ and $\alpha \in \mathbb{R}$. The space $H_\alpha^p(G) = H_\alpha^p$ is the space of all $f \in S'(G)$ for which $f^* \in L_\alpha^p$. We set

$$\|f\|_{H_\alpha^p} = \|f^*\|_{p, \alpha},$$

and we denote H_0^p and $\|f\|_{H_0^p}$ by H^p and $\|f\|_{H^p}$, respectively.

We now turn to the definition of the atomic Hardy spaces with power weight

DEFINITION 3.2. Let $0 < p \leq 1$ and $\alpha > -1$. A function $a: G \rightarrow \mathbb{C}$ is a $(p, \infty)_\alpha$ atom if there exists a set $I = x + G_n$ such that

- (i) $\text{supp } a \subset I$,
- (ii) $\|a\|_\infty \leq (\mu_\alpha(I))^{-1/p}$,
- (iii) $\int_G a(x) d\mu(x) = 0$.

Clearly every $(p, \infty)_\alpha$ atom defines an element of $S'(G)$. Moreover, an argument like in [1, page 611] shows that each $(p, \infty)_\alpha$ atom a belongs to H_α^p with $\|a\|_{H^p} \leq 1$.

DEFINITION 3.3. Let $0 < p \leq 1$ and $\alpha > -1$. The space $H_\alpha^{p, \infty}(G) = H_\alpha^{p, \infty}$ is the space of all $f \in S'(G)$ for which there exists a sequence $(\lambda_i)_1^\infty \in l^p$ and a sequence of $(p, \infty)_\alpha$ atoms $(a_i)_1^\infty$ such that

$$(3.4) \quad f = \sum_{i=1}^{\infty} \lambda_i a_i \quad \text{in } S'(G).$$

We set

$$\|f\|_{H_\alpha^{p, \infty}} = \inf \left\{ \left(\sum_{i=1}^{\infty} |\lambda_i|^p \right)^{1/p} \right\},$$

where the infimum is taken over all decompositions of f of the form (3.4).

THEOREM 3.5. Let $0 < p \leq 1$ and $-1 < \alpha \leq 0$. Then $H_\alpha^p = H_\alpha^{p, \infty}$ and the "norms" on these spaces are equivalent.

The proof of Theorem 3.5 will be preceded by a lemma.

LEMMA 3.6. Let $0 < p \leq 1$ and $-1 < \alpha \leq 0$. If $f \in H_\alpha^p$ then each $f_n = f * \Delta_n$ belongs to $H_\alpha^{p, \infty}$ and

$$\|f_n\|_{H_\alpha^{p, \infty}} \leq C \|f\|_{H_\alpha^p},$$

with C independent of $n \in \mathbb{Z}$.

PROOF. Let $f \in H_\alpha^p$ and for each $k \in \mathbb{Z}$ let

$$\Omega_k = \{x \in G: f^*(x) > 2^k\}.$$

If $y \in \Omega_k$ then there exists an $N \in \mathbb{Z}$ so that $f_N(y) > 2^k$ and this implies that $y + G_N \subset \Omega_k$. If $A(y) = \{n \in \mathbb{Z}: y + G_n \subset \Omega_k\}$, then $A(y)$ is bounded from below because $f^* \in L_\alpha^p$. Thus there exists an $\alpha(y) \in \mathbb{Z}$ so that $y + G_{\alpha(y)} \subset \Omega_k$ and $y + G_n \not\subset \Omega_k$ for all $n < \alpha(y)$. We shall denote the at most countably many different sets $y + G_{\alpha(y)}$ with $y \in \Omega_k$ by $y_{k,i} + G_{\alpha(k,i)} := I_{k,i}$. Then $\Omega_k = \bigcup_i I_{k,i}$ and $I_{k,i} \cap I_{k,j} = \emptyset$ for $i \neq j$.

Next, let $\tilde{I}_{k,i} = y_{k,i} + G_{\alpha(k,i)-1}$ and let $\tilde{\Omega}_k = \bigcup_i \tilde{I}_{k,i}$. If necessary we first rename the sets $\tilde{I}_{k,i}$ so that they are mutually disjoint.

Also, observe that for each $k \in \mathbb{Z}$, $\Omega_{k+1} \subset \Omega_k$ and, since $f \in H_\alpha^p$, $\mu_\alpha(\Omega_k) < \infty$ and $\mu_\alpha(\bigcap_{-\infty}^\infty \Omega_k) = 0$, which implies that $\lim_{k \rightarrow \infty} \mu_\alpha(\Omega_k) = 0$.

Next, for each function $f_n = f * \Delta_n$ and each $k \in \mathbb{Z}$, let

$$\Omega_k^n = \{x \in G: |f_n(x)| > 2^k\}.$$

Then $\Omega_k^n \subset \Omega_k$.

For $k, n \in \mathbb{Z}$ we define the function $g_k^n: G \rightarrow \mathbb{C}$ by

$$g_k^n(x) = \begin{cases} f_n(x) & \text{if } x \notin \tilde{\Omega}_k, \\ p_{k,i}^n & \text{if } x \in \tilde{I}_{k,i}, \end{cases}$$

where

$$p_{k,i}^n = (\mu(\tilde{I}_{k,i}))^{-1} \int_{\tilde{I}_{k,i}} f_n(x) d\mu(x).$$

We first show that for a.e. $x \in G$,

$$(i) \lim_{k \rightarrow -\infty} g_k^n(x) = 0,$$

$$(ii) \lim_{k \rightarrow \infty} g_k^n(x) = f_n(x).$$

To prove (i), consider $x \in \tilde{I}_{k,i} = y + G_l$, say. If $n \leq l$ then f_n is constant on $y + G_l$ and, since $\tilde{I}_{k,i} \not\subset \Omega_k$ we see that $|f_n(x)| \leq 2^k$ on $\tilde{I}_{k,i}$ and this implies that $|p_{k,i}^n| \leq 2^k$. If $n > l$, then

$$p_{k,i}^n = (f * \Delta_n) * \Delta_l(y) = f * \Delta_l(y) = f_l(y),$$

which again implies that $|p_{k,i}^n| \leq 2^k$. Therefore, we see that $|g_k^n(x)| \leq 2^k$ for all $x \in G$ and hence (i) holds.

To prove (ii), observe that $\mu_\alpha(\bigcap_{-\infty}^\infty \Omega_k) = 0$ implies that $\mu(\bigcap_{-\infty}^\infty \Omega_k) = 0$ and hence, $\mu(\bigcap_{-\infty}^\infty \tilde{\Omega}_k) = 0$. This last equality immediately implies (ii). It

follows from (i) and (ii) that for a.e. $x \in G$,

$$f_n(x) = \sum_{k=-\infty}^{\infty} (g_{k+1}^n - g_k^n)(x),$$

that is,

$$(3.7) \quad f_n(x) = \sum_{k=-\infty}^{\infty} \sum_i (g_{k+1}^n - g_k^n)(x) \chi_{\tilde{I}_{k,i}}(x).$$

For each k, i, n let

$$b_{k,i}^n = (g_{k+1}^n - g_k^n) \chi_{\tilde{I}_{k,i}}.$$

Then $\text{supp } b_{k,i}^n \subset \tilde{I}_{k,i}$, $\|b_{k,i}^n\|_{\infty} \leq 2^{k+2}$ and a routine calculation shows that

$$(3.8) \quad \int_G b_{k,i}^n(x) d\mu(x) = 0.$$

We now prove that

$$(3.9) \quad f_n = \sum_{k=-\infty}^{\infty} \sum_i b_{k,i}^n,$$

with the series in (3.9) converging to f_n in $S'(G)$. To do so, take any $\varphi \in S(G)$ with, say, $\text{supp } \varphi \subset G_t$ for some $t \in \mathbb{Z}$. We need to prove that

$$(3.10) \quad \lim_{\substack{n_1 \rightarrow -\infty \\ n_2, n_3 \rightarrow \infty}} \int_G \sum_{k=n_1}^{n_2} \sum_{i \leq n_3} b_{k,i}^n(x) \varphi(x) d\mu(x) = \int_G f_n(x) \varphi(x) d\mu(x).$$

We first prove three auxiliary results, (3.11), (3.12) and (3.13).

(3.11) There exists an $N_1 \in -\mathbb{N}$ such that

$$A := \sum_{k=-\infty}^{N_1} \sum_i \|b_{k,i}^n \varphi\|_1 \leq 1.$$

We have

$$\begin{aligned} A &\leq \sum_{k=-\infty}^{N_1} \|g_{k+1}^n - g_k^n\|_{\infty} \|\varphi\|_1 \\ &\leq \sum_{k=-\infty}^{N_1} 2^{k+2} \|\varphi\|_1 \leq 2^{N_1+3} \|\varphi\|_1 \leq 1, \end{aligned}$$

for suitably chosen $N_1 \in -\mathbb{N}$.

(3.12) There exists an $N_2 \in \mathbb{N}$ so that for every $k > N_2$, every $i \in \mathbb{N}$ and $n \in \mathbb{Z}$,

$$\langle b_{k,i}^n, \varphi \rangle = \int_G b_{k,i}^n(x) \varphi(x) d\mu(x) = 0.$$

Since $\varphi \in \mathcal{S}(G)$, there exists an $s \in \mathbb{Z}$ such that φ is constant on the cosets of G_s in G but not on the cosets of G_{s-1} (unless $\varphi(x) \equiv 0$). Hence there exist $x_1, \dots, x_r \in G$ such that $x_i + G_s \cap x_j + G_s = \emptyset$ for $i \neq j$ and $\text{supp } \varphi = \bigcup_{j=1}^r x_j + G_s$. Also, since $\lim_{k \rightarrow \infty} \mu_\alpha(\Omega_k) = 0$, [7, Lemma 1(c)] implies that $\lim_{k \rightarrow \infty} \mu_\alpha(\tilde{\Omega}_k) = 0$. Consequently, there exists an $N_2 \in \mathbb{N}$ such that for all $k > N_2$ and all $i \in \mathbb{N}$ we have

$$\mu_\alpha(\tilde{I}_{k,i}) \leq \mu_\alpha(\tilde{\Omega}_k) \leq \min\{\mu_\alpha(x_j + G_s): 1 \leq j \leq r\}.$$

Because each $\tilde{I}_{k,i}$ is a coset of some subgroup G_l of G we see that for $k \geq N_2$ we have either $\tilde{I}_{k,i} \subset x_j + G_s$ for some j , or else $\tilde{I}_{k,i} \cap x_j + G_s = \emptyset$ for all j , $1 \leq j \leq r$. In the latter case we have $\tilde{I}_{k,i} \cap \text{supp } \varphi = \emptyset$ and hence $\langle b_{k,i}^n, \varphi \rangle = 0$ for all $n \in \mathbb{Z}$; in case $\tilde{I}_{k,i} \subset x_j + G_s$ for some j with $1 \leq j \leq r$, we again have $\langle b_{k,i}^n, \varphi \rangle = 0$ for all $n \in \mathbb{Z}$, because (3.8) holds. This proves (3.12).

(3.13) With N_1, N_2 chosen so that (3.11) and (3.12) hold, there exists an $N_3 \in \mathbb{N}$ so that

$$B := \sum_{k=N_1+1}^{N_2} \sum_{i \geq N_3} \|b_{k,i}^n \varphi\|_1 \leq 1.$$

We have

$$B \leq \sum_{k=N_1+1}^{N_2} \sum_{i \geq N_3} \int_{\tilde{I}_{k,i}} |(g_{k+1}^n - g_k^n)(x)| |\varphi(x)| v_{-\alpha}(x) d\mu_\alpha(x).$$

Since $v_{-\alpha}(x) \leq (m_l)^\alpha$ for $\alpha \leq 0$ and $x \in G_l$, we see that

$$B \leq \|\varphi\|_\infty (m_l)^\alpha \sum_{k=N_1+1}^{N_2} \sum_{i \geq N_3} 2^{k+2} \mu_\alpha(\tilde{I}_{k,i}).$$

Now we observe that for every $k \in \mathbb{Z}$ there exists an $i_k \in \mathbb{N}$ so that

$$\sum_{i \geq i_k} \mu_\alpha(\tilde{I}_{k,i}) < (2^{N_2+3} \|\varphi\|_\infty (m_l)^\alpha)^{-1}.$$

Let $N_3 = \max\{i_k: N_1 < k \leq N_2\}$. Then for this choice of N_3 we immediately obtain (3.13).

Applying (3.11), (3.12) and (3.13) it is easy to see that for every $n_1 \in -\mathbb{N}$ and $n_2, n_3 \in \mathbb{N}$,

$$\sum_{k=n_1}^{n_2} \sum_{i \geq n_3} b_{k,i}^n(x) \varphi(x)$$

is dominated pointwise on G by an integrable function. Thus, in view of (3.7), the Lebesgue Dominated Convergence Theorem implies (3.10) and, therefore, (3.9).

Finally, let

$$\lambda_{k,i} = 2^{k+2}(\mu_\alpha(\tilde{I}_{k,i}))^{1/p} \quad \text{and} \quad a_{k,i}^n = (\lambda_{k,i})^{-1} b_{k,i}^n.$$

Then each $a_{k,i}^n$ is a $(p, \infty)_\alpha$ atom and

$$f_n = \sum_{k,i} \lambda_{k,i} a_{k,i}^n \quad \text{in } S'(G).$$

Furthermore, a straightforward computation shows that

$$\sum_{k,i} |\lambda_{k,i}|^p \leq C \|f_n^*\|_{p,\alpha}^p \leq C \|f^*\|_{p,\alpha}^p = C \|f\|_{H_\alpha^p}^p.$$

This completes the proof of Lemma 3.6.

PROOF OF THEOREM 3.5. Take any $f \in H_\alpha^p$. Using the same notation as in the proof of Lemma 3.6, we see from the definition of the $(p, \infty)_\alpha$ atoms $a_{k,i}^n$ that

$$\sup_{n \in \mathbb{N}} \|a_{0,1}^n\|_\infty \leq (\mu_\alpha(\tilde{I}_{0,1}))^{-1/p}.$$

Thus the Banach-Alaoglu theorem implies the existence of a subsequence $(a_{0,1}^{n_{\nu(0,1)}})$ of $(a_{0,1}^n)$ so that this subsequence converges in the weak* topology of $L^\infty(G)$ to, say, $a_{0,1} \in L^\infty(G)$. Clearly, $a_{0,1}$ is a $(p, \infty)_\alpha$ atom with $\text{supp } a_{0,1} \subset \tilde{I}_{0,1}$. Next, since

$$\sup_{n_{\nu(0,1)}} \|a_{1,1}^{n_{\nu(0,1)}}\|_\infty \leq (\mu_\alpha(\tilde{I}_{1,1}))^{-1/p},$$

a second application of the Banach-Alaoglu theorem yields a subsequence $(a_{1,1}^{n_{\nu(1,1)}})$ of $(a_{1,1}^{n_{\nu(0,1)}})$ and a $(p, \infty)_\alpha$ atom $a_{1,1}$ with $\text{supp } a_{1,1} \subset \tilde{I}_{1,1}$ so that the subsequence converges weak* in $L^\infty(G)$ to $a_{1,1}$. Arranging the pairs of subscripts (k, i) with $k \in \mathbb{Z}$ and $i \in \mathbb{N}$ in a sequence we can repeat the process described above for each (k, i) . By the usual diagonalization method we obtain a sequence (n_ν) and a sequence of $(p, \infty)_\alpha$ atoms $a_{k,i}$ with $\text{supp } a_{k,i} \subset \tilde{I}_{k,i}$ so that for all (k, i) we have

$$(3.14) \quad \lim_{\nu \rightarrow \infty} a_{k,i}^{n_\nu} = a_{k,i} \quad \text{weak* in } L^\infty.$$

We shall prove that

$$(3.15) \quad f = \sum_{k=-\infty}^{\infty} \sum_i \lambda_{k,i} a_{k,i} \quad \text{in } S'(G).$$

To do so, take any $\varphi \in \mathcal{S}(G)$ and assume, like in Lemma 3.6, that $\text{supp } \varphi \subset G_t$. Let $\varepsilon > 0$ be given. We first derive three auxiliary inequalities, (3.16), (3.17) and (3.18).

(3.16) There exists an $M_1 \in -\mathbb{N}$ so that for all $n \in \mathbb{Z}$ we have

$$(i) \sum_{k=-\infty}^{M_1} \sum_i |\langle \lambda_{k,i} a_{k,i}^n, \varphi \rangle| < \frac{\varepsilon}{12},$$

$$(ii) \sum_{k=-\infty}^{M_1} \sum_i |\langle \lambda_{k,i} a_{k,i}, \varphi \rangle| < \frac{\varepsilon}{12}.$$

The proof of (3.16) is virtually the same as the proof of (3.11).

(3.17) There exists an $M_2 \in \mathbb{N}$ so that for all $k > M_2$, every $i \in \mathbb{N}$ and $n \in \mathbb{Z}$ we have

$$(i) \langle \lambda_{k,i} a_{k,i}^n, \varphi \rangle = 0,$$

$$(ii) \langle \lambda_{k,i} a_{k,i}, \varphi \rangle = 0.$$

This is essentially a restatement of (3.12) with $M_2 = N_2$.

(3.18) With M_1, M_2 chosen as in (3.16) and (3.17), there exists an $M_3 \in \mathbb{N}$ and an $n_{\nu_1} \in (n_{\nu})_{\nu=1}^{\infty}$ so that for all $n_{\nu} \geq n_{\nu_1}$ we have

$$\sum_{k=M_1+1}^{M_2} \sum_i |\lambda_{k,i}| |\langle a_{k,i}^{n_{\nu}} - a_{k,i}, \varphi \rangle| < \frac{\varepsilon}{6}.$$

To prove (3.18), we observe that for each $k \in \mathbb{Z}$ the sets $\tilde{I}_{k,i}$ are mutually disjoint so that at most r of the sets $\tilde{I}_{k,i}$ will contain at least one of the sets $x_j + G_s$, with $x_j + G_s$ as defined in the proof of (3.12). Let

$$\tilde{i}_k = \max\{i: x_j + G_s \subset \tilde{I}_{k,i} \text{ for some } j \text{ with } 1 \leq j \leq r\},$$

and let

$$M_3 = \max\{\tilde{i}_k: M_1 < k \leq M_2\}.$$

Clearly, if $M_1 < k \leq M_2$, $i > M_3$ and $n \in \mathbb{Z}$, then $\langle a_{k,i}^n, \varphi \rangle = \langle a_{k,i}, \varphi \rangle = 0$. Furthermore, in view of (3.14) there exists an n_{ν_1} so that for all $n_{\nu} \geq n_{\nu_1}$ we have

$$\begin{aligned} & \sum_{k=M_1+1}^{M_2} \sum_i |\lambda_{k,i}| |\langle a_{k,i}^{n_{\nu}} - a_{k,i}, \varphi \rangle| \\ &= \sum_{k=M_1+1}^{M_2} \sum_{i \leq M_3} |\lambda_{k,i}| |\langle a_{k,i}^{n_{\nu}} - a_{k,i}, \varphi \rangle| < \frac{\varepsilon}{6}, \end{aligned}$$

which proves (3.18).

Now we observe that since $\lim_{n \rightarrow \infty} f_n = f$ in $\mathcal{S}'(G)$, there exists an $n_{\nu_2} \geq n_{\nu_1}$ so that

$$(3.19) \quad |\langle f_{n_{\nu_2}} - f, \varphi \rangle| < \frac{\varepsilon}{3}.$$

In the proof of Lemma 3.6 we saw that there exist $N_1 \in -\mathbb{N}$ and $N_2, N_3 \in \mathbb{N}$, with N_1, N_2, N_3 depending on n_{ν_2} , so that if $n_1 \leq N_1$, $n_2 \geq N_2$ and $n_3 \geq N_3$ then

$$(3.20) \quad \left| \sum_{k=n_1}^{n_2} \sum_{i \leq n_3} \langle \lambda_{k,i} a_{k,i}^{n_{\nu_2}} - f_{n_{\nu_2}}, \varphi \rangle \right| < \frac{\varepsilon}{3}.$$

Consequently, if $l_1 \leq \min\{M_1, N_1\}$, $l_2 \geq N_2$ and $l_3 \geq \max\{M_3, N_3\}$ then

$$\begin{aligned} & \left| \left\langle f - \sum_{k=l_1}^{l_2} \sum_{i \leq l_3} \lambda_{k,i} a_{k,i}, \varphi \right\rangle \right| \\ &= \left| \left\langle f - \sum_{k=l_1}^{N_2} \sum_{i \leq l_3} \lambda_{k,i} a_{k,i}, \varphi \right\rangle \right| \quad (\text{by (3.12)}) \\ &\leq |\langle f - f_{n_{\nu_2}}, \varphi \rangle| + \left| \left\langle f_{n_{\nu_2}} - \sum_{k=l_1}^{N_2} \sum_{i \leq l_3} \lambda_{k,i} a_{k,i}^{n_{\nu_2}}, \varphi \right\rangle \right| \\ &\quad + \sum_{k=M_1+1}^{N_2} \sum_{i \leq l_3} |\lambda_{k,i}| |\langle a_{k,i}^{n_{\nu_2}} - a_{k,i}, \varphi \rangle| \\ &\quad + \sum_{k=l_1}^{M_1} \sum_{i \leq l_3} |\langle \lambda_{k,i} a_{k,i}^{n_{\nu_2}}, \varphi \rangle| + \sum_{k=l_1}^{M_1} \sum_{i \leq l_3} |\langle \lambda_{k,i} a_{k,i}, \varphi \rangle| \\ &\leq \varepsilon/3 + \varepsilon/3 + \varepsilon/6 + \varepsilon/12 + \varepsilon/12 = \varepsilon. \end{aligned}$$

This proves (3.15). In the proof of Lemma 3.6 we saw that

$$\sum_{k,i} |\lambda_{k,i}|^p \leq C \|f\|_{H_\alpha^p}^p.$$

Therefore, $f \in H_\alpha^{p,\infty}$ and

$$\|f\|_{H_\alpha^{p,\infty}} \leq C \|f\|_{H_\alpha^p}.$$

To prove the converse, take any $f \in H_\alpha^{p,\infty}$. Then $f = \sum_k \lambda_k a_k$ in $\mathcal{S}'(G)$, where $(\lambda_k) \in l^p$ and each a_k is a $(p, \infty)_\alpha$ atom so that $\|a_k^*\|_{p,\alpha} \leq 1$. Consequently,

$$|f^*(x)|^p \leq \sum_k |\lambda_k|^p |a_k^*(x)|^p$$

and this implies that $\|f^*\|_{p,\alpha}^p \leq \sum_k |\lambda_k|^p$, that is, $f \in H_\alpha^p$ and

$$\|f\|_{H_\alpha^p} \leq \|f\|_{H_\alpha^{p,\infty}}.$$

This completes the proof of Theorem 3.5.

We mention here the following corollary whose simple proof will be omitted.

COROLLARY 3.21. *For each q with $1 \leq q < \infty$ we have $L^q \cap H_\alpha^p$ is dense in H_α^p .*

The last theorem of this section is an interpolation theorem for operators on H^p and L^p spaces. The theorem is a version for locally compact Vilenkin groups of [3, Theorems III.6.4 and 6.5] or [1, Theorem D], where also the precise definitions of some of the concepts used here can be found.

THEOREM 3.22. *Let $0 < p_0 \leq 1 < p_1 < \infty$. Suppose T is a sublinear operator of weak type (H^{p_0}, p_0) on H^{p_0} and of weak type (p_1, p_1) on L^{p_1} . Then T is bounded from H^p to L^p for $p_0 < p \leq 1$ and T is bounded from L^p to L^p for $1 < p < p_1$.*

PROOF (Outline). Let $f \in L^p$ with $1 < p < p_1$ and choose q so that $1 < q < p$. For $t > 0$ let

$$E_t = \{x: M_q(|f|)(x) = (|f|^q)^*(x) > t^q\}.$$

As in the proof of Lemma 3.6 we can express E_t as a disjoint union of maximal cosets of certain subgroups G_n of G , say $E_t = \bigcup_j I_j$.

Define $g_t: G \rightarrow \mathbb{C}$ by

$$g_t(x) = \begin{cases} f(x) & \text{if } x \notin E_t, \\ (\mu(I_j))^{-1} \int_{I_j} f(x) d\mu(x) & \text{if } x \in I_j, \end{cases}$$

and define $b_t: G \rightarrow \mathbb{C}$ by $b_t(x) = f(x) - g_t(x)$. Then

$$b_t(x) = \sum_j (f - g_t)(x) \chi_{I_j}(x) = \sum_j b_j(x).$$

We have

$$((\mu(I_j))^{-1} \int_{I_j} |b_j(x)|^q d\mu(x))^{1/q} \leq Ct,$$

and if we set

$$a_j(x) = (Ct(\mu(I_j))^{1/p_0})^{-1} b_j(x),$$

then each a_j is a (p_0, q) atom and

$$b_t(x) = \sum_j Ct(\mu(I_j))^{1/p} a_j(x).$$

Thus $b_t \in H^{p_0, q}$ and $\|b_t\|_{H^{p_0, q}} \leq Ct(\mu(E_t))^{1/p}$. Moreover, $|g_t(x)| \leq Ct$ for $x \in E_t$, and for $x \notin E_t$ we have $|f(x)| \leq M(|f|)(x) \leq \{M_q(|f|)(x)\}^{1/q} \leq t$.

Consequently,

$$\begin{aligned} \int_G |g_t(x)|^{p_1} d\mu(x) &= \int_{G \setminus E_t} |g_t(x)|^{p_1} d\mu(x) + \int_{E_t} |g_t(x)|^{p_1} d\mu(x) \\ &\leq \int_{|f| \leq t} |f(x)|^{p_1} d\mu(x) + (Ct)^{p_1} \mu(E_t) \\ &\leq Ct^{p_1-p} \|f\|_p^p, \end{aligned}$$

that is, $g_t \in L^{p_1}$ for every $t > 0$. The rest of the proof is virtually the same as the proof of [3, Theorems III.6.4 and 6.5] and will be omitted.

4. Multipliers on $H_\alpha^p(G)$

As mentioned in the introduction, in this section we shall present our multiplier theorems for the spaces H_α^p . Throughout this section, if $\varphi \in L^\infty(\Gamma)$ and if $k \in \mathbb{Z}$ we let $\varphi_k = \varphi \chi_{\Gamma_k}$ and $\varphi^k = \varphi_{k+1} - \varphi_k$. We begin with a definition which extends a definition given by Kitada in [5].

DEFINITION 4.1. Let $0 < p \leq 1$ and $-1 < \alpha \leq 0$. Let X denote H_α^p and let Y denote H_α^p or L_α^p . A function $\varphi \in L^\infty(\Gamma)$ is a multiplier from X to Y ($\varphi \in \mathcal{M}(X, Y)$ or $\varphi \in \mathcal{M}(X)$ in case $X = Y$) if there exists a constant $C > 0$ so that for all $f \in X \cap L^2$ we have $(\varphi \hat{f})^\vee \in Y$ and $\|(\varphi \hat{f})^\vee\|_Y \leq C \|f\|_X$.

REMARK 4.2. In order to prove that $\varphi \in \mathcal{M}(X, Y)$ it is sufficient to prove that there exists a $C > 0$ so that for every $(p, \infty)_\alpha$ atom a and for every $k \in \mathbb{Z}$ we have $\|(\varphi_k)^\vee * a\|_Y = \|(\varphi_k \hat{a})^\vee\|_Y \leq C$. To see this, take any $(p, \infty)_\alpha$ atom a and let $\hat{a}_k = \hat{a} \chi_{\Gamma_k}$. Then $\lim_{k \rightarrow \infty} \hat{a}_k = \hat{a}$ in $L^2(\Gamma)$. Consequently,

$$\lim_{k \rightarrow \infty} \varphi_k \hat{a} = \lim_{k \rightarrow \infty} \varphi \hat{a}_k = \varphi \hat{a} \quad \text{in } L^2(\Gamma)$$

and hence,

$$(4.3) \quad \lim_{k \rightarrow \infty} (\varphi_k \hat{a})^\vee = (\varphi \hat{a})^\vee \quad \text{in } L^2(G).$$

Now we distinguish two cases.

(i) Let $Y = L_\alpha^p$. Then (4.3) implies the existence of a subsequence (k_i) so that

$$\lim_{i \rightarrow \infty} (\varphi_{k_i} \hat{a})^\vee(x) = (\varphi \hat{a})^\vee(x) \quad \text{for a.e. } x \in G.$$

Thus, Fatou's Lemma implies that

$$\|(\varphi \hat{a})^\vee\|_{p, \alpha} \leq \liminf \|(\varphi_{k_i} \hat{a})^\vee\|_{p, \alpha} \leq C.$$

From this inequality we easily derive that $\varphi \in \mathcal{M}(X, Y)$.

(ii) Let $Y = X = H_\alpha^p$. Then (4.3) implies that

$$\lim_{k \rightarrow \infty} (\varphi_k \hat{a})^\vee = (\varphi \hat{a})^\vee \quad \text{in } S'(G).$$

Now a simple argument shows that for every $x \in G$,

$$(((\varphi \hat{a})^\vee)^*(x))^p \leq \liminf (((\varphi_k \hat{a})^\vee)^*(x))^p$$

and an application of Fatou's Lemma shows that

$$\|(\varphi \hat{a})^\vee\|_{H_\alpha^p} \leq \liminf \|(\varphi_k \hat{a})^\vee\|_{H_\alpha^p} \leq C$$

and this inequality immediately implies that $\varphi \in \mathcal{M}(X)$.

We now turn to the discussion of our multiplier theorems for the spaces H_α^p . Our first result deals with multipliers from the spaces H_α^p to the corresponding spaces L_α^p . We start with a lemma in which we consider the case $\alpha = 0$.

LEMMA 4.4. *Let $\varphi \in L^\infty(\Gamma)$ and let $0 < p \leq 1$. If*

$$\sup_k (m_k)^{1/p-1} \|(\varphi_k)^\vee\|_{K(1/p-1/r, r, p)} < \infty$$

for some r with $p < r < \infty$ then

(i) $\varphi \in \mathcal{M}(H^s, L^s)$ for $p \leq s \leq 1$

and

(ii) $\varphi \in \mathcal{M}(L^s)$ for $1 < s < \infty$.

PROOF. Let a be a (p, ∞) atom with $\text{supp } a \subset I = x_0 + G_n$ for some $x_0 \in G$ and $n \in \mathbb{Z}$. For every $k \in \mathbb{Z}$ we have

$$\begin{aligned} \|(\varphi_k \hat{a})^\vee\|_p^p &= \|(\varphi_k)^\vee * a\|_p^p \\ &= \|((\varphi_k)^\vee * a)\chi_I\|_p^p + \|((\varphi_k)^\vee * a)\chi_{G \setminus I}\|_p^p := A + B. \end{aligned}$$

Applying Hölder's inequality we see that

$$\begin{aligned} A &\leq \left(\int_I |(\varphi_k)^\vee * a(x)|^2 d\mu(x) \right)^{p/2} \cdot (\mu(I))^{1-p/2} \\ &\leq C \|(\varphi_k)^\vee * a\|_2^p \cdot (m_n)^{-(1-p/2)} \\ &\leq C \|\varphi_k\|_\infty^p \|a\|_2^p \cdot (m_n)^{-(1-p/2)} \\ &\leq C \|\varphi\|_\infty^p, \end{aligned}$$

because a is a (p, ∞) atom.

For B we have

$$\begin{aligned} B &= \int_{G \setminus I} \left| \int_G (\varphi_k)^\vee(t) a(x-t) d\mu(t) \right|^p d\mu(x) \\ &\leq \|a\|_\infty^p \int_{G \setminus I} \left(\int_{x-I} |(\varphi_k)^\vee(t)| d\mu(t) \right)^p d\mu(x). \end{aligned}$$

Since $\varphi_k(\gamma) = 0$ for $\gamma \in \Gamma \setminus \Gamma_k$, $(\varphi_k)^\vee$ is constant on the cosets of G_k . Thus, if $(x_i + G_k)_{i=0}^\infty$ represent the different cosets of G_k in G , then

$$(\varphi_k)^\vee(t) = \sum_{i=0}^\infty (\varphi_k)^\vee(x_i) \chi_{x_i + G_k}(t),$$

so

$$\begin{aligned} \left(\int_{x-I} |(\varphi_k)^\vee(t)| d\mu(t) \right)^p &= \left(\sum_{\{i: x_i \in x-I\}} |(\varphi_k)^\vee(x_i)| (m_k)^{-1} \right)^p \\ &\leq \sum_{\{i: x_i \in x-I\}} |(\varphi_k)^\vee(x_i)|^p (m_k)^{-p} \\ &= (m_k)^{-(p-1)} \int_{x-I} |(\varphi_k)^\vee(t)|^p d\mu(t). \end{aligned}$$

Therefore,

$$\begin{aligned} B &\leq \|a\|_\infty^p \cdot (m_k)^{1-p} \int_{G \setminus I} \int_I |(\varphi_k)^\vee(x-t)|^p d\mu(t) d\mu(x) \\ &= \|a\|_\infty^p \cdot (m_k)^{1-p} \int_{G_n} \int_{G \setminus G_n} |(\varphi_k)^\vee(y-u)|^p d\mu(y) d\mu(u). \end{aligned}$$

Next we observe that for each $u \in G_n$,

$$\begin{aligned} \int_{G \setminus G_n} |(\varphi_k)^\vee(y-u)|^p d\mu(y) &= \int_{G \setminus G_n} |(\varphi_k)^\vee(y)|^p d\mu(y) \\ &= \sum_{j=-\infty}^{n-1} \int_{G_j \setminus G_{j+1}} |(\varphi_k)^\vee(y)|^p d\mu(y) \\ &\leq \sum_{j=-\infty}^{n-1} \left(\int_{G_j \setminus G_{j+1}} |(\varphi_k)^\vee(y)|^r d\mu(y) \right)^{p/r} \cdot (\mu(G_j \setminus G_{j+1}))^{1-p/r} \\ &\leq C \sum_{j=-\infty}^{n-1} ((m_j)^{-(1/p-1/r)} \|(\varphi_k)^\vee \chi_{G_j \setminus G_{j+1}}\|_r)^p \\ &\leq C \|(\varphi_k)^\vee\|_{K(1/p-1/r, r, p)}^p. \end{aligned}$$

Since $\|a\|_\infty \leq (\mu(I))^{-1/p} \leq C(m_n)^{1/p}$, we see that

$$B \leq C m_n (m_k)^{1-p} (m_n)^{-1} (m_k)^{p-1} = C.$$

Consequently, $\varphi \in \mathcal{M}(H^p, L^p)$. Since $\varphi \in L^\infty(\Gamma)$, we have $\varphi \in \mathcal{M}(L^2)$. Thus, an application of Theorem 3.22 and a duality argument complete the proof of the lemma.

THEOREM 4.5. *Let $\varphi \in L^\infty(\Gamma)$ and let $0 < p \leq 1$. If*

$$\sup_k (m_k)^{1/p-1} \|(\varphi_k)^\vee\|_{K(1/p-1/r, r, p)} < \infty$$

for some r with $p < r < \infty$, then $\varphi \in \mathcal{M}(H_\alpha^p, L_\alpha^p)$ for $-1 + p/r < \alpha \leq 0$.

PROOF. Let a be a $(p, \infty)_\alpha$ atom. We shall distinguish two cases, depending on $\text{supp } a$. First assume $\text{supp } a \subset I = x_0 + G_n$ with $x_0 \notin G_n$. Then $x_0 \in G_j \setminus G_{j+1}$ for some $j < n$ and $\mu_\alpha(I) = (m_j)^{-\alpha} (m_n)^{-1}$, so that $\|a\|_\infty \leq ((m_j)^\alpha m_n)^{1/p}$. For each $k \in \mathbb{Z}$ we have

$$\begin{aligned} \|(\varphi_k \hat{a})^\vee\|_{p, \alpha}^p &= \|(\varphi_k)^\vee * a\|_{p, \alpha}^p \\ &= \|((\varphi_k)^\vee * a)\chi_I\|_{p, \alpha}^p + \|((\varphi_k)^\vee * a)\chi_{G \setminus I}\|_{p, \alpha}^p := A + B, \end{aligned}$$

where

$$\begin{aligned} A &= \int_I |(\varphi_k)^\vee * a(x)|^p v_\alpha(x) d\mu(x) \\ &\leq (m_j)^{-\alpha} \left(\int_I |(\varphi_k)^\vee * a(x)|^2 d\mu(x) \right)^{p/2} \cdot (\mu(I))^{1-p/2} \\ &\leq (m_j)^{-\alpha} (m_n)^{-(1-p/2)} \|(\varphi_k)^\vee * a\|_2^p \\ &\leq (m_j)^{-\alpha} (m_n)^{-(1-p/2)} \|\varphi\|_\infty^p \|a\|_2^p \leq \|\varphi\|_\infty^p. \end{aligned}$$

To estimate B we observe that, as in the proof of Lemma 4.4,

$$\begin{aligned} B &= \int_{G \setminus I} \left| \int_G (\varphi_k)^\vee(x-t) a(t) d\mu(t) \right|^p d\mu_\alpha(x) \\ &\leq \|a\|_\infty^p \int_{G \setminus I} \left(\int_{x-I} |(\varphi_k)^\vee(t)| d\mu(t) \right)^p d\mu_\alpha(x) \\ &\leq \|a\|_\infty^p \cdot (m_k)^{1-p} \int_{G \setminus I} \int_I |(\varphi_k)^\vee(x-t)|^p d\mu(t) d\mu_\alpha(x) \\ &\leq \|a\|_\infty^p \cdot (m_k)^{1-p} \int_{G_n} \int_{G \setminus G_n} |(\varphi_k)^\vee(y-u)|^p v_\alpha(x_0+y) d\mu(y) d\mu(u). \end{aligned}$$

We now estimate the inner integral, first writing it as a sum of three integrals

$$\begin{aligned} \int_{G \setminus G_n} \cdots d\mu(y) &= \int_{G \setminus G_j} \cdots + \int_{G_j \setminus G_{j+1}} \cdots + \int_{G_{j+1} \setminus G_n} \cdots d\mu(y) \\ &:= B_1 + B_2 + B_3. \end{aligned}$$

For $x_0 \in G_j \setminus G_{j+1}$ and $y \notin G_j$ we have $x_0 + y \notin G_j$, so that $v_\alpha(x_0 + y) \leq (m_j)^{-\alpha}$. Therefore, if $u \in G_n$ we obtain, as in the proof of Lemma 4.4,

$$\begin{aligned} B_1 &\leq (m_j)^{-\alpha} \int_{G \setminus G_n} |(\varphi_k)^\vee(y - u)|^p d\mu(y) \\ &\leq C(m_j)^{-\alpha} \|(\varphi_k)^\vee\|_{K(1/p-1/r, r, p)}^p \leq C(m_j)^{-\alpha} (m_k)^{p-1}. \end{aligned}$$

For $x_0 \in G_j \setminus G_{j+1}$ and $y \in G_{j+1} \setminus G_n$ we have $x_0 + y \in G_j \setminus G_{j+1}$ and hence, $v_\alpha(x_0 + y) = (m_j)^{-\alpha}$. Therefore, if $u \in G_n$ then

$$B_3 \leq (m_j)^{-\alpha} \int_{G \setminus G_n} |(\varphi_k)^\vee(y - u)|^p d\mu(y) \leq C(m_j)^{-\alpha} (m_k)^{p-1}.$$

Finally, to find the appropriate estimate for B_2 , observe that for $u \in G_n$,

$$\begin{aligned} B_2 &\leq \left(\int_{G_j \setminus G_{j+1}} |(\varphi_k)^\vee(y - u)|^r d\mu(y) \right)^{p/r} \\ &\quad \cdot \left(\int_{G_j \setminus G_{j+1}} (v_\alpha(x_0 + y))^{r/(r-p)} d\mu(y) \right)^{(r-p)/r} \\ &\leq C \|(\varphi_k)^\vee\|_{\chi_{G_j \setminus G_{j+1}}}^p \cdot (m_j)^{-(\alpha+1-p/r)} \\ &\leq C(m_j)^{-\alpha} \|(\varphi_k)^\vee\|_{K(1/p-1/r, r, p)}^p \\ &\leq C(m_j)^{-\alpha} (m_k)^{p-1}. \end{aligned}$$

Therefore,

$$B \leq C \|a\|_\infty^p \cdot (m_k)^{1-p} (m_j)^{-\alpha} (m_k)^{p-1} (m_n)^{-1} \leq C,$$

because a is a $(p, \infty)_\alpha$ atom. Thus we see that $\|(\varphi_k \hat{a})^\vee\|_{p, \alpha}^p \leq C$.

In case $\text{supp } a \subset G_n$ we have $\|a\|_\infty \leq (\mu_\alpha(G_n))^{-1/p} \leq C(m_n)^{(\alpha+1)/p}$, and for each $k \in \mathbb{Z}$,

$$\begin{aligned} \|(\varphi_k \hat{a})^\vee\|_{p, \alpha}^p &= \|(\varphi_k)^\vee * a\|_{p, \alpha}^p \\ &= \|((\varphi_k)^\vee * a)\chi_{G_n}\|_{p, \alpha}^p + \|((\varphi_k)^\vee * a)\chi_{G \setminus G_n}\|_{p, \alpha}^p \\ &:= A + B. \end{aligned}$$

Choose $s > 1$ so that $-1 + p/s < \alpha$. Then, according to Lemma 4.4, $\varphi_k \in \mathcal{M}(L^s)$ and we see that

$$\begin{aligned} A &\leq \left(\int_{G_n} |(\varphi_k)^\vee * a(x)|^s d\mu(x) \right)^{p/s} \cdot \left(\int_{G_n} (v_\alpha(x))^{s/(s-p)} d\mu(x) \right)^{(s-p)/s} \\ &\leq C \|(\varphi_k)^\vee * a\|_s^p \cdot (m_n)^{-(\alpha+1-p/s)} \\ &\leq C \|a\|_s^p \cdot (m_n)^{-(\alpha+1-p/s)} \leq C. \end{aligned}$$

Moreover, as in the first part of the proof, we have

$$\begin{aligned} B &\leq \|a\|_\infty^p \cdot (m_k)^{1-p} \int_{G_n} \int_{G \setminus G_n} |(\varphi_k)^\vee(y-u)|^p v_\alpha(y) d\mu(y) d\mu(u) \\ &\leq C \|a\|_\infty^p (m_k)^{1-p} (m_n)^{-\alpha} (m_k)^{p-1} (m_n)^{-1} \leq C. \end{aligned}$$

Thus, we see again that $\|((\varphi_k \hat{a})^\vee)\|_{p,\alpha}^p \leq C$. According to Remark 4.2 we may conclude that $\varphi \in \mathcal{M}(H_\alpha^p, L_\alpha^p)$.

The next theorem deals with multipliers from H_α^p to H_α^p . We begin with a lemma which extends [9, Theorem 2].

LEMMA 4.6. *Let $\varphi \in L^\infty(\Gamma)$ and $0 < p \leq 1$. If*

$$\sup_k (m_k)^{1/p-1} \sum_{j=k}^{\infty} \|(\varphi^j)^\vee\|_{K(1/p-1/r, r, 1)} < \infty$$

for some r with $1 \leq r < \infty$ then $\varphi \in \mathcal{M}(H^s)$ for $1 \leq s < \infty$.

PROOF. We first prove that $\varphi \in \mathcal{M}(H^1)$ by showing that there exists a $C > 0$ so that for all $(1, \infty)$ atoms a we have $\|(\varphi \hat{a})^\vee\|_{H^1} \leq C$. We may assume that $\text{supp } a \subset G_n$ for some $n \in \mathbb{Z}$. Let $f = (\varphi \hat{a})^\vee$ and let $f^* = \sup_l |f * \Delta_l|$. Kitada showed in [4, Theorem 2] that

$$\int_{G_n} f^*(x) d\mu(x) \leq C$$

and

$$\int_{G \setminus G_n} f^*(x) d\mu(x) \leq \sum_{j=n}^{\infty} \sum_{k=-\infty}^{n-1} \|(\varphi^j)^\vee \chi_{G_k \setminus G_{k+1}}\|_1.$$

Applying Hölder's inequality we see that for $k < n$

$$\begin{aligned} \|(\varphi^j)^\vee \chi_{G_k \setminus G_{k+1}}\|_1 &\leq \|(\varphi^j)^\vee \chi_{G_k \setminus G_{k+1}}\|_r \cdot (m_k)^{-1/r'} \\ &= (m_k)^{1/p-1} (m_k)^{-(1/p-1/r)} \|(\varphi^j)^\vee \chi_{G_k \setminus G_{k+1}}\|_r \\ &\leq (m_n)^{1/p-1} (m_k)^{-(1/p-1/r)} \|(\varphi^j)^\vee \chi_{G_k \setminus G_{k+1}}\|_r. \end{aligned}$$

Hence, we have

$$\begin{aligned} \int_{G \setminus G_n} f^*(x) d\mu(x) &\leq (m_n)^{1/p-1} \sum_{j=n}^{\infty} \sum_{k=-\infty}^{n-1} (m_k)^{-(1/p-1/r)} \|(\varphi^j)^\vee \chi_{G_k \setminus G_{k+1}}\|_r \\ &\leq (m_n)^{1/p-1} \sum_{j=n}^{\infty} \|(\varphi^j)^\vee\|_{K(1/p-1/r, r, 1)} \leq C. \end{aligned}$$

Therefore,

$$\|f\|_{H^1} = \int_{G_n} f^*(x) d\mu(x) + \int_{G \setminus G_n} f^*(x) d\mu(x) \leq C,$$

that is, $\varphi \in \mathcal{M}(H^1)$.

We now show that $\varphi \in \mathcal{M}(H^s)$ for $1 < s < \infty$. Since $\varphi \in \mathcal{M}(H^1)$ there exists a $C > 0$ so that for all $f \in H^1 \cap L^2$ we have

$$\|Tf\|_1 := \|(\varphi \hat{f})^\vee\|_{L^1} \leq \|(\varphi \hat{f})^\vee\|_{H^1} \leq C\|f\|_{H^1},$$

that is, $\varphi \in \mathcal{M}(H^1, L^1)$. Since $H^1 \cap L^2$ is a dense subset of H^1 , the operator T can be extended to H^1 so that $\|Tf\|_{H^1} \leq C\|f\|_{H^1}$ for all $f \in H^1$. This implies immediately that T is of weak type $(H^1, 1)$ on H^1 . Since $\varphi \in \mathcal{M}(L^2)$, T is of type $(2, 2)$ on L^2 . Thus, it follows from Theorem 3.22 and a standard duality argument that T is of type (s, s) , that is, $\varphi \in \mathcal{M}(L^s) = \mathcal{M}(H^s)$ for each $1 < s < \infty$.

THEOREM 4.7. *Let $\varphi \in L^\infty(\Gamma)$ and $0 < p \leq 1$. If*

$$\sup_k (m_k)^{1-p} \sum_{j=k}^{\infty} \|(\varphi^j)^\vee\|_{K(1/p-1/r, r, p)}^p < \infty$$

for some r with $1 \leq r < \infty$ then $\varphi \in \mathcal{M}(H_\alpha^p)$ for $-1 + p/r < \alpha \leq 0$.

PROOF. Since $0 < p \leq 1$ we have

$$\begin{aligned} (m_k)^{1-p} \left(\sum_{j=k}^{\infty} \|(\varphi^j)^\vee\|_{K(1/p-1/r, r, 1)} \right)^p &\leq (m_k)^{1-p} \sum_{j=k}^{\infty} \|(\varphi^j)^\vee\|_{K(1/p-1/r, r, 1)}^p \\ &\leq C(m_k)^{1-p} \sum_{j=k}^{\infty} \|(\varphi^j)^\vee\|_{K(1/p-1/r, r, p)}^p < \infty. \end{aligned}$$

It follows from Lemma 4.6 that $\varphi \in \mathcal{M}(H^s)$ for $1 \leq s < \infty$.

To see that $\varphi \in \mathcal{M}(H_\alpha^p)$ for $-1 + p/r < \alpha \leq 0$, let a be a $(p, \infty)_\alpha$ atom with $\text{supp } a \subset I = x_0 + G_n$. Take any $k \in \mathbb{Z}$ and let $f = (\varphi_k \hat{a})^\vee = (\varphi_k)^\vee * a$

and let $f^* = \sup_I |f * \Delta_I|$. We have

$$\begin{aligned} & \int_G (f^*(x))^p d\mu_\alpha(x) \\ &= \int_I (f^*(x))^p d\mu_\alpha(x) + \int_{G \setminus I} (f^*(x))^p d\mu_\alpha(x) := A + B. \end{aligned}$$

To estimate A we distinguish two cases.

(i) If $x_0 \in G_n$ then for every $r \in [1, \infty)$ and each α with $-1 + p/r < \alpha \leq 0$ we have

$$\begin{aligned} A &\leq \left(\int_{G_n} (f^*(x))^r d\mu(x) \right)^{p/r} \cdot \left(\int_{G_n} (v_\alpha(x))^{r/(r-p)} d\mu(x) \right)^{(r-p)/r} \\ &\leq C \|a\|_{H^r}^p \cdot (m_n)^{-(\alpha+1-p/r)} \leq C, \end{aligned}$$

where the second inequality is obtained by observing that $\varphi \in \mathcal{M}(H^r)$.

(ii) If $x_0 \notin G_n$ then $x_0 \in G_l \setminus G_{l+1}$ for some $l < n$ and $I \subset G_l \setminus G_{l+1}$. With r and α as in (i) we have

$$\begin{aligned} A &\leq \left(\int_I (f^*(x))^r d\mu(x) \right)^{p/r} \cdot \left(\int_I (v_\alpha(x))^{r/(r-p)} d\mu(x) \right)^{(r-p)/r} \\ &\leq C \|a\|_{H^r}^p \cdot (m_l)^{-\alpha} (m_n)^{-(1-p/r)} \leq C. \end{aligned}$$

To find the appropriate estimate for B we closely follow Kitada's proof of [5, Theorem 2]. If we set $\psi(\gamma) = \overline{\gamma(x_0)}\varphi(\gamma)$, $\psi^j = \psi \chi_{\Gamma_{l+1} \setminus \Gamma_l}$ and $b(x) = a(x + x_0)$, then Kitada showed that

$$f^*(x) \leq \sum_{j=n}^{\infty} |(\psi^j)^\vee * b(x)|.$$

Therefore,

$$\begin{aligned} B &= \int_{G \setminus I} (f^*(x))^p d\mu_\alpha(x) \\ &\leq \sum_{j=n}^{\infty} \sum_{i=-\infty}^{n-1} \int_{J_i} |(\psi^j)^\vee * b(x)|^p d\mu_\alpha(x), \end{aligned}$$

where $J_i = I_i \setminus I_{i+1}$ and $I_i = x_0 + G_i$. For each of the integrals in this sum Kitada showed that

$$\begin{aligned} B_{ij} &:= \int_{J_i} |(\psi^j)^\vee * b(x)|^p d\mu_\alpha(x) \\ &= \int_{J_i} |(\psi^j)^\vee \chi_{J_i} * b(x)|^p d\mu_\alpha(x). \end{aligned}$$

Consequently, applying [7, Lemma 1(b)] to obtain the third inequality, we see that

$$\begin{aligned}
 B_{ij} &\leq \left(\int_{J_i} |(\psi^j)^\vee \chi_{J_i} * b(x)|^r d\mu(x) \right)^{p/r} \\
 &\quad \cdot \left(\int_{J_i} (v_\alpha(x))^{r/(r-p)} d\mu(x) \right)^{(r-p)/r} \\
 &\leq \|b\|_1^p \|(\psi^j)^\vee \chi_{J_i}\|_r^p \cdot (\mu_{\alpha r/(r-p)}(I_i))^{(r-p)/r} \\
 &\leq \|a\|_1^p \|(\varphi^j)^\vee \chi_{G_i \setminus G_{i+1}}\|_r^p \cdot C((m_i)^{-1} \inf\{v_{\alpha r/(r-p)}(y): y \in I_i \setminus \{0\}\})^{(r-p)/r} \\
 &\leq C \|a\|_1^p \|(\varphi^j)^\vee \chi_{G_i \setminus G_{i+1}}\|_r^p \\
 &\quad \cdot (m_i)^{(p-r)/r} (\inf\{v_{\alpha r/(r-p)}(y): y \in I_i \setminus \{0\}\})^{(r-p)/r}
 \end{aligned}$$

(a) If $I_n = I = G_n$ we have

$$B_{ij} \leq C(m_n)^{\alpha+1-p} \|(\varphi^j)^\vee \chi_{G_i \setminus G_{i+1}}\|_r^p \cdot (m_i)^{(p-r)/r} (m_n)^{-\alpha}.$$

(b) If $I_n \subset G_l \setminus G_{l+1}$ for some $l < n$ we have

$$B_{ij} \leq C(m_l)^\alpha (m_n)^{1-p} \|(\varphi^j)^\vee \chi_{G_i \setminus G_{i+1}}\|_r^p \cdot (m_i)^{(p-r)/r} (m_l)^{-\alpha}.$$

Thus, in both cases we see that

$$\begin{aligned}
 B &\leq C(m_n)^{1-p} \sum_{j=n}^{\infty} \sum_{i=-\infty}^{n-1} ((m_i)^{1/r-1/p} \|(\varphi^j)^\vee \chi_{G_i \setminus G_{i+1}}\|_r)^p \\
 &\leq C(m_n)^{1-p} \sum_{j=n}^{\infty} \|(\varphi^j)^\vee\|_{K(1/p-1/r, r, p)}^p \leq C.
 \end{aligned}$$

Thus, $\|f^*\|_{p, \alpha} = \|f\|_{H_\alpha^p} \leq C$ and this implies that $\varphi \in \mathcal{M}(H_\alpha^p)$.

For $0 < p < 1$ we have the following corollary.

COROLLARY 4.8. *Let $\varphi \in L^\infty(\Gamma)$ and $0 < p < 1$. If*

$$\sup_k (m_k)^{1/p-1} \|(\varphi^k)^\vee\|_{K(1/p-1/r, r, p)} < \infty$$

for some r with $1 \leq r < \infty$ then $\varphi \in \mathcal{M}(H_\alpha^p)$ for $-1 + p/r < \alpha \leq 0$.

PROOF. For $0 < p < 1$ we have

$$\sum_{j=k}^{\infty} \|(\psi^j)^\vee\|_{K(1/p-1/r, r, p)}^p \leq C \sum_{j=k}^{\infty} (m_j)^{p-1} \leq C (m_k)^{p-1}.$$

The result follows immediately from Theorem 4.7.

We now show that Corollary 4.8 is sharp in a certain sense. The example we use to prove the sharpness result is a variation of the example used in [9] to prove that certain results of Kitada for H^p multipliers, $0 < p < 1$, were best possible.

THEOREM 4.9. *Let $0 < p < 1$ and $1 \leq r < \infty$. There exists a $\varphi \in L^\infty(\Gamma)$ so that*

- (i) $\sup_k (m_k)^{1/p-1} \|(\varphi^k)^\vee\|_{K(1/p-1/r, r, q)} < \infty$ for every $q > p$;
- (ii) $\varphi \in \mathcal{M}(H_\alpha^q)$ for all q with $p < q < 1$ and α with $-1 + q/r < \alpha \leq 0$;
- (iii) $\varphi \notin \mathcal{M}(H_\alpha^p)$ for any α with $-1 < \alpha \leq 0$.

PROOF. Choose $\gamma_1 \in \Gamma_1 \setminus \Gamma_0$ and define $f: G \rightarrow \mathbb{C}$ by

$$f(x) = \sum_{k=-\infty}^{-1} \left(\frac{m_k}{|k|} \right)^{1/p} \gamma_1(x) \chi_{G_k \setminus G_{k+1}}(x).$$

Then $f \in L^1(G)$ and for every $r \geq p$ we have

$$\begin{aligned} \|f\|_{K(1/p-1/r, r, q)}^q &= \sum_{k=-\infty}^{\infty} (m_k)^{-(1/p-1/r)q} \|f \chi_{G_k \setminus G_{k+1}}\|_r^q \\ &\cong \sum_{k=-\infty}^{-1} \left(\frac{1}{|k|} \right)^{q/p} < \infty \Leftrightarrow q > p. \end{aligned}$$

Moreover, if $q > p$ then

$$\begin{aligned} \|f\|_{K(1/q-1/r, r, q)}^q &= \sum_{k=-\infty}^{\infty} (m_k)^{-(1/q-1/r)q} \|f \chi_{G_k \setminus G_{k+1}}\|_r^q \\ &= \sum_{k=-\infty}^{-1} (m_k)^{-1+q/r} \left(\frac{m_k}{|k|} \right)^{q/p} (\mu(G_k \setminus G_{k+1}))^{q/r} \\ &\leq C \sum_{k=-\infty}^{-1} \left(\frac{1}{|k|} \right)^{q/p} (m_k)^{-1+q/p} < \infty. \end{aligned}$$

Also,

$$\hat{f}(\gamma) = \sum_{k=-\infty}^{-1} \left(\frac{m_k}{|k|} \right)^{1/p} (\hat{\chi}_{G_k} - \hat{\chi}_{G_{k+1}})(\gamma - \gamma_1),$$

with $\hat{\chi}_{G_k} = m_k \chi_{\Gamma_k}$. Thus $\text{supp } \hat{f} \subset \gamma_1 + \Gamma_0 \subset \Gamma_1 \setminus \Gamma_0$. Let $\varphi = \hat{f}$. Then $\varphi \in L^\infty(\Gamma)$. Moreover, $\varphi^k = 0$ for $k \neq 1$ and $\varphi^k = \varphi$ for $k = 1$, so that $(\varphi^1)^\vee = f$ and $(\varphi^k)^\vee = 0$ for $k \neq 1$. Therefore φ satisfies (i) and,

according to Corollary 4.8, φ satisfies (ii). To see that φ satisfies (iii), choose for every $i < 0$ an $x_i \in G_i \setminus G_{i+1}$ and define, for $-1 < \alpha \leq 0$, functions $g_i: G \rightarrow \mathbb{C}$ by

$$g_i(x) = (m_i)^{\alpha/p} (m_1 \chi_{x_i + G_1} - m_0 \chi_{x_i + G_0})(x).$$

Then g_i is a multiple of a $(p, \infty)_\alpha$ atom and $\|g_i\|_{H_\alpha^p} \leq m_1$. Moreover,

$$\hat{g}_i(\gamma) = (m_i)^{\alpha/p} \overline{\gamma(x_i)} (\chi_{\Gamma_1} - \chi_{\Gamma_0})(\gamma),$$

so $\text{supp } \hat{g}_i \subset \Gamma_1 \setminus \Gamma_0$.

Furthermore, if we define $h_i: G \rightarrow \mathbb{C}$ by

$$h_i(x) = (m_i)^{\alpha/p} \sum_{k=-\infty}^{-1} \left(\frac{m_k}{|k|} \right)^{1/p} \gamma_1(x - x_i) \chi_{G_k \setminus G_{k+1}}(x - x_i)$$

then $h_i \in L^1(G)$, and a straightforward computation shows that $\hat{h}_i = \varphi \hat{g}_i$, that is, $h_i = (\varphi \hat{g}_i)^\vee$. Furthermore, we have

$$\|h_i\|_{p, \alpha}^p = \int_G |h_i(x)|^p d\mu_\alpha(x) \geq C \sum_{k=i}^{-1} |k|^{-1}$$

so $\lim_{i \rightarrow -\infty} \|h_i\|_{p, \alpha} = \infty$. Since each $h_i \in L^1(G)$ we have

$$\|h_i\|_{H_\alpha^p} = \|h_i^*\|_{p, \alpha} \geq \|h_i\|_{p, \alpha},$$

so

$$\lim_{i \rightarrow -\infty} \|(\varphi \hat{g}_i)^\vee\|_{H_\alpha^p} = \lim_{i \rightarrow -\infty} \|h_i\|_{H_\alpha^p} = \lim_{i \rightarrow -\infty} \|h_i^*\|_{p, \alpha} = \infty$$

and this implies that $\varphi \notin \mathcal{M}(H_\alpha^p)$.

In his most recent paper on multipliers on $H^p(G)$ spaces [6], Kitada proved a multiplier result for Hardy spaces on locally compact Vilenkin groups in which his assumptions are the natural analogue for G of the usual Hörmander condition for multipliers for function spaces on \mathbb{R}^n . Before stating Kitada's main result we first repeat a definition given in [6].

DEFINITION 4.10. Let $\varphi \in L^\infty(\Gamma)$. For $\lambda > 0$ and $j \in \mathbb{Z}$ let $D^\lambda \varphi^j$ be defined by

$$D^\lambda \varphi^j = (|x|^\lambda (\varphi^j)^\vee(x))^\wedge.$$

We say that $\varphi \in \mathcal{M}(s, \lambda)$, where $1 \leq s \leq \infty$, if

$$B(\varphi, s, \lambda) := \|\varphi\|_\infty + \sup_j (m_j)^{\lambda-1/s} \|D^\lambda \varphi^j\|_s < \infty.$$

In [6, Theorem 2] Kitada proved the following, which is the analogue for G of [12, Theorem (4.11)].

THEOREM K. *Let $0 < p \leq 1$ and $1 \leq s < \infty$. If $\varphi \in M(s, \lambda)$ for $\lambda > 1/p - 1/\max(2, s')$ then $\varphi \in M(H^p)$.*

We conclude this paper by extending Theorem K to power-weighted Hardy spaces. Our proof depends on Corollary 4.8 and is somewhat different from Kitada's proof of Theorem K. We first establish a simple lemma.

LEMMA 4.11. *Let $\varphi \in L^\infty(\Gamma)$, let $0 < p \leq 1$ and $1 \leq r < \infty$. If*

$$(4.12) \quad \sup_k (m_k)^{1/p-1+\varepsilon} \|(\varphi^k)^\vee\|_{K(1/p-1/r+\varepsilon, r, \infty)} < \infty \quad \text{for some } \varepsilon > 0,$$

then

$$(4.13) \quad \sup_k (m_k)^{1/p-1} \|(\varphi^k)^\vee\|_{K(1/p-1/r, r, p)} < \infty.$$

PROOF. We have

$$\begin{aligned} \|(\varphi^k)^\vee\|_{K(1/p-1/r, r, p)}^p &= \sum_{i=-\infty}^k (m_i)^{-1+p/r} \|(\varphi^k)^\vee \chi_{G_i \setminus G_{i+1}}\|_r^p \\ &\quad + \sum_{i=k+1}^{\infty} (m_i)^{-1+p/r} \|(\varphi^k)^\vee \chi_{G_i \setminus G_{i+1}}\|_r^p \\ &:= A + B. \end{aligned}$$

Assumption (4.12) implies that

$$\begin{aligned} A &\leq C \sum_{i=-\infty}^k (m_i)^{-1+p/r} (m_i)^{1-p/r+\varepsilon p} (m_k)^{-1+p-\varepsilon p} \\ &\leq C(m_k)^{-1+p-\varepsilon p} \sum_{i=-\infty}^k (m_i)^{\varepsilon p} \leq C(m_k)^{-1+p}, \end{aligned}$$

since $\varepsilon p > 0$.

To estimate B first observe that

$$\|(\varphi^k)^\vee\|_\infty \leq \|\varphi^k\|_1 \leq \|\varphi^k\|_\infty \cdot \lambda(\Gamma_{k+1} \setminus \Gamma_k) \leq C\|\varphi\|_\infty \cdot m_k.$$

Therefore,

$$\begin{aligned} B &\leq C \sum_{i=k+1}^{\infty} (m_i)^{-1+p/r} (m_k)^p (m_i)^{-p/r} \\ &= C(m_k)^p \sum_{i=k+1}^{\infty} (m_i)^{-1} \leq C(m_k)^{p-1}. \end{aligned}$$

From the inequalities for A and B we immediately obtain (4.13).

COROLLARY 4.14. Let $\varphi \in L^\infty(\Gamma)$, let $0 < p \leq 1$ and $1 \leq r < \infty$. If φ satisfies (4.12) then $\varphi \in \mathcal{M}(H_\alpha^p)$ for $-1 + p/r < \alpha \leq 0$.

PROOF. For $p = 1$ this is [5, Theorem 2]. For $0 < p < 1$ we apply Lemma 4.11 and Corollary 4.8.

THEOREM 4.15. Let $\varphi \in L^\infty(\Gamma)$, let $0 < p \leq 1$, $1 < r \leq \infty$ and $\lambda > 1/p - 1/\max(2, r')$. If $\varphi \in M(r, \lambda)$ then $\varphi \in \mathcal{M}(H_\alpha^p)$ for $-1 + p/\min(2, r') < \alpha \leq 0$.

PROOF. We first assume that $1 < r \leq 2$ so that $\max(2, r') = r'$. Since $\lambda > 1/p - 1/r'$, there exists an $\varepsilon > 0$ so that $\lambda = 1/p - 1/r' + \varepsilon$. Now we consider

$$\begin{aligned} \|(\varphi^j)^\vee\|_{K(1/p-1/r'+\varepsilon, r', \infty)} &= \|(\varphi^j)^\vee\|_{K(\lambda, r', \infty)} \leq \|(\varphi^j)^\vee\|_{K(\lambda, r', r')} \\ &= \|(\varphi^j)^\vee\|_{r', \lambda r'} = \| |x|^\lambda (\varphi^j)^\vee \|_{r'} = \| (D^\lambda \varphi^j)^\vee \|_{r'} \leq \| (D^\lambda \varphi^j) \|_{r'}. \end{aligned}$$

Thus, if $\varphi \in M(r, \lambda)$ then φ satisfies inequality (4.12), and Corollary 4.14 implies that $\varphi \in \mathcal{M}(H_\alpha^p)$ for $-1 + p/r' < \alpha \leq 0$.

If $2 < r \leq \infty$ then $\max(2, r') = 2$. In this case there exists an $\varepsilon > 0$ such that $\lambda = 1/p - 1/2 + \varepsilon$ and we have

$$\|(\varphi^j)^\vee\|_{K(1/p-1/2+\varepsilon, 2, \infty)} = \|(\varphi^j)^\vee\|_{K(\lambda, 2, \infty)} \leq \|(\varphi^j)^\vee\|_{K(\lambda, 2, 2)} \leq \|D^\lambda \varphi^j\|_2.$$

An application of [6, Proposition 2] to obtain the third inequality, shows that

$$\begin{aligned} \sup_j (m_j)^{1/p-1+\varepsilon} \|(\varphi^j)^\vee\|_{K(1/p-1/2+\varepsilon, 2, \infty)} \\ \leq \sup_j (m_j)^{\lambda-1/2} \|D^\lambda \varphi^j\|_2 \leq B(\varphi, 2, \lambda) \\ \leq CB(\varphi, r, \lambda) < \infty \end{aligned}$$

and the conclusion of the theorem follows again from Corollary 4.14. This completes the proof of Theorem 4.15.

REMARK. Professor Kitada informed the authors that he obtained independently essentially the same result as our Theorem 4.15.

References

- [1] R. Coifman and G. Weiss, 'Extensions of Hardy spaces and their use in analysis', *Bull. Amer. Math. Soc.* **83** (1977), 569–645.
- [2] R. E. Edwards and G. I. Gaudry, *Littlewood-Paley and multiplier theory*, (Springer-Verlag, Berlin, 1977).

- [3] J. Garcia-Cuerva and J. L. Rubio de Francia, *Weighted norm inequalities and related topics*, (North-Holland Mathematics Studies 116, North-Holland, Amsterdam, 1985).
- [4] T. Kitada, ' H^p -multiplier theorems on certain totally disconnected groups', *Sci. Rep. Hirosaki Univ.* **34** (1987), 1–7.
- [5] T. Kitada, 'Multipliers on weighted Hardy spaces over certain totally disconnected groups', *Internat. J. Math. Math. Sci.* **11** (1988), 665–675.
- [6] T. Kitada, ' H^p multiplier theorem on certain totally disconnected groups, II', preprint.
- [7] C. W. Onneweer, 'Multipliers on weighted L^p -spaces over certain totally disconnected groups', *Trans. Amer. Math. Soc.* **288** (1985), 347–357.
- [8] C. W. Onneweer and T. S. Quek, 'Multipliers on weighted L^p -spaces over locally compact Vilenkin groups', *Proc. Amer. Math. Soc.* **105** (1989), 622–631.
- [9] C. W. Onneweer and T. S. Quek, ' H^p multiplier results on locally compact Vilenkin groups', *Quart. J. Math. Oxford Ser. (2)* **40** (1989), 313–323.
- [10] M. H. Taibleson, 'Harmonic analysis on n -dimensional vector spaces over local fields, III. Multipliers', *Math. Ann.* **187** (1970), 259–271.
- [11] M. H. Taibleson, *Fourier analysis on local fields*, (Mathematical Notes 15, Princeton University Press, 1975).
- [12] M. H. Taibleson and G. Weiss, *The molecular characterization of certain Hardy spaces*, (Astérisque 77, pp. 68–149, 1980).
- [13] N. Ya. Vilenkin, *On a class of complete orthonormal systems*, (Amer. Math. Soc. Transl. (2) 28, pp. 1–35, 1963).

Department of Mathematics and
Statistics
University of New Mexico
Albuquerque, New Mexico 87131
U.S.A.

Department of Mathematics
National University of Singapore
Singapore 0511
Republic of Singapore