

EIGENELEMENTS OF PERTURBED OPERATORS

B. V. LIMAYE and M. T. NAIR

(Received 23 January 1989)

Communicated by S. Yamamuro

Abstract

Let λ_0 be a semisimple eigenvalue of an operator T_0 . Let Γ_0 be a circle with centre λ_0 containing no other spectral value of T_0 . Some lower bounds are obtained for the convergence radius of the power series for the spectral projection $P(t)$ (and for trace $T(t)P(t)$) associated with a linear perturbation family $T(t) = T_0 + tV_0$ and the circle Γ_0 . They are useful when T_0 is a member of a sequence (T_n) which approximates an operator T in a collectively compact manner. These bounds result from a modification of Kato's method of majorizing series, based on an idea of Redont. If λ_0 is simple, it is shown that the same lower bounds are valid for the convergence radius of a power series yielding an eigenvector of $T(t)$.

1980 *Mathematics subject classification* (Amer. Math. Soc.) (1985 Revision): 47 A 55, 47 A 70, 47 A 10, 41 A 35.

1. Introduction

Let T_0 and V_0 be closed (linear) operators on a domain D_0 , which is a dense subspace of a Banach space X over the field \mathbb{C} of complex numbers. For $t \in \mathbb{C}$, consider the operator

$$T(t) = T_0 + tV_0.$$

Then $T(0) = T_0$ is called the *unperturbed operator*, tV_0 the *perturbation* and $T(t)$ the *perturbed operator*. This situation typically arises in quantum mechanics as follows. Let T_0 be the Hamiltonian of a quantum mechanical system and V_0 be a potential energy operator. Then $T(1) = T_0 + V_0$ is the Hamiltonian of the perturbed system. Also, in operator approximation

theory, one considers a sequence of operators (T_n) which approximates a given operator T . One can let $T_0 = T_n$ for some fixed integer n and $V_0 = T - T_0$, so as to have $T = T(1)$.

Let $\sigma(T_0)$ denote the spectrum of T_0 and λ_0 be an isolated point of it. Let Γ_0 be a simple closed curve which separates λ_0 from the rest of $\sigma(T_0)$. For $z \notin \sigma(T_0)$, let $R_0(z) = (T_0 - zI)^{-1}$. Consider the following open disk in \mathbb{C} :

$$\partial(\Gamma_0) = \left\{ t \in \mathbb{C} : |t| < \min_{z \in \Gamma_0} (1/r_\sigma(V_0 R_0(z))) \right\},$$

where r_σ denotes the spectral radius. For all $t \in \partial(\Gamma_0)$, it follows that $T(t)$ is a closed operator on the domain D_0 , and Γ_0 does not intersect the spectrum $\sigma(T(t))$ of $T(t)$. For $z \notin \sigma(T(t))$, let $R(t, z) = (T(t) - zI)^{-1}$. Then the spectral projection

$$P(t) = -\frac{1}{2\pi i} \int_{\Gamma_0} R(t, z) dz,$$

associated with $T(t)$ and Γ_0 , is an analytic function on $\partial(\Gamma_0)$. If the algebraic multiplicity of λ_0 is $m < \infty$, that is, if $P(0)$ has rank m , then for all $t \in \partial(\Gamma_0)$, $P(t)$ has rank m and Γ_0 contains precisely m eigenvalues of $T(t)$: $\lambda_1(t), \dots, \lambda_m(t)$, repeated according to their algebraic multiplicities. Also, the function

$$\hat{\lambda}(t) \equiv \frac{1}{m} \text{trace } T(t)P(t) = (\lambda_1(t) + \dots + \lambda_m(t))/m$$

is analytic for t in $\partial(\Gamma_0)$.

It is of interest to know for which values of the parameter t , the power series for $P(t)$ and $\hat{\lambda}(t)$ converge. The method of majorizing series developed by Kato [2, pages 89–91] allows one to find lower bounds for the convergence radii of these series. A modification of Kato's method yields improved results for the case where λ_0 is a semisimple eigenvalue of T_0 (Lemma 2.1 and Remark 2.2). This procedure also gives an upper bound for $|\lambda_i(t) - \lambda_0|$, $i = 1, \dots, m$, and a disk of isolation for the set $\{\lambda_1(t), \dots, \lambda_m(t)\}$ from the rest of the spectrum of $T(t)$ (Theorem 2.3). The basic idea behind these results is given in Redont's unpublished work [7, pages 79–80] (compare also [6, Proposition 2.1] and [4, Theorem 10.5]. The results are applicable when a compact operator T is approximated in a collectively compact manner (Remark 2.5).

In case λ_0 is a simple eigenvalue of T_0 , let ϕ_0 (respectively ϕ_0^*) be an eigenvector of T_0 (respectively T_0^*) corresponding to λ_0 (respectively $\overline{\lambda_0}$) such that $\langle \phi_0, \phi_0^* \rangle = 1$, and consider the meromorphic function

$$\phi(t) = P(t)\phi_0 / \langle P(t)\phi_0, \phi_0^* \rangle.$$

If $t_0 \in \partial(\Gamma_0)$ is not a pole of the function ϕ , then $\phi(t_0)$ is an eigenvector of $T(t_0)$.

We describe a subregion of $\partial(\Gamma_0)$, which is devoid of any pole of the function ϕ (Proposition 3.2). This again yields a lower bound for the convergence radius of the power series for $\phi(t)$ about 0 (Theorem 3.3 and Remark 3.4).

2. Eigenvalues of $T(t)$ near λ_0

We assume that λ_0 is a semisimple eigenvalue of T_0 of algebraic multiplicity m , that is, the range of the spectral projection $P(0) = P_0$ coincides with the eigenspace of T_0 corresponding to λ_0 and is of dimension m . Let S_0 denote the reduced resolvent associated with T_0 and λ_0 :

$$(1) \quad S_0 = \lim_{z \rightarrow \lambda_0} R_0(z)(I - P_0).$$

Then $S_{0|(I-P_0)D_0}$ is the inverse of $(T_0 - \lambda_0 I)_{|(I-P_0)D_0}$. Since $\sigma(T_{0|(I-P_0)D_0}) = \sigma(T_0) \setminus \{\lambda_0\}$, it follows that

$$(2) \quad \frac{1}{\text{dist}(\lambda_0, \sigma(T_0) \setminus \{\lambda_0\})} = r_\sigma(S_0) \leq \|S_0\|.$$

Let $0 < \varepsilon < 1$. Consider the circle Γ_ε with centre λ_0 and radius $\varepsilon/\|S_0\|$. The convergence radii of the series for $P(t)$ and $\hat{\lambda}(t)$ are greater than or equal to the radius of $\partial(\Gamma_\varepsilon)$. To find a lower bound for the radius of $\partial(\Gamma_\varepsilon)$, we introduce the following notation:

$$(3) \quad \begin{aligned} s &= \|S_0\|, \quad p = \|V_0 P_0\|, \quad q = \|V_0 S_0\|, \\ r^2 &= \sup\{\|V_0 S_0^{k+1} V_0 S_0\|/\|S_0\|^k, k = 0, 1, 2, \dots\}. \end{aligned}$$

LEMMA 2.1. *Let $0 < \varepsilon < 1$. The radius of $\partial(\Gamma_\varepsilon)$ is at least*

$$\left[\frac{p^2 s^2}{\varepsilon^2} + \frac{2psq}{\varepsilon(1-\varepsilon)} + \frac{r^2}{(1-\varepsilon)^2} \right]^{-1/2}.$$

The convergence radii of the power series for $P(t)$ and $\hat{\lambda}(t)$ about 0 are at least equal to

$$(4) \quad m_0 = \max_{0 < \varepsilon < 1} \left[\frac{p^2 s^2}{\varepsilon^2} + \frac{2psq}{\varepsilon(1-\varepsilon)} + \frac{r^2}{(1-\varepsilon)^2} \right]^{-1/2}.$$

PROOF. Since λ_0 is semisimple, we have $T_0 P_0 = \lambda_0 P_0$. Hence λ_0 is a pole of order 1 of $R_0(z)$. The Laurent expansion of $R_0(z)$ about λ_0 is given by

$$R_0(z) = \sum_{k=0}^{\infty} S_0^{k+1} (z - \lambda_0)^k - \frac{P_0}{z - \lambda_0}$$

for $0 < |z - \lambda_0| < \text{dist}(\lambda_0, \sigma(T_0) \setminus \{\lambda_0\})$ (compare [2, (5.18)]). Writing $R_0(z) = R_0(z)P_0 + R_0(z)(I - P_0)$, we see that

$$[V_0 R_0(z)]^2 = A(z) + B_1(z) + B_2(z) + C(z),$$

where

$$\begin{aligned} A(z) &= [V_0 R_0(z) P_0]^2 = (V_0 P_0)^2 / (z - \lambda_0)^2, \\ B_1(z) &= V_0 R_0(z) P_0 V_0 R_0(z) (I - P_0) \\ &= - \sum_{k=0}^{\infty} V_0 P_0 V_0 S_0^{k+2} (z - \lambda_0)^k - (V_0 P_0 V_0 S_0) / (z - \lambda_0), \\ B_2(z) &= V_0 R_0(z) (I - P_0) V_0 R_0(z) P_0 \\ &= - \sum_{k=0}^{\infty} V_0 S_0^{k+2} V_0 P_0 (z - \lambda_0)^k - (V_0 S_0 V_0 P_0) / (z - \lambda_0) \end{aligned}$$

and

$$C(z) = [V_0 R_0(z) (I - P_0)]^2 = \sum_{k=0}^{\infty} \sum_{j=0}^{\infty} V_0 S_0^{k+1} V_0 S_0^{j+1} (z - \lambda_0)^{k+j}.$$

For $z \in \Gamma_\varepsilon$, we have $|z - \lambda_0| = \varepsilon/s$, so that

$$\|A(z)\| = \frac{p^2 s^2}{\varepsilon^2}, \quad \|B_1(z)\|, \|B_2(z)\| \leq pqs \left(\frac{1}{1-\varepsilon} + \frac{1}{\varepsilon} \right) = \frac{psq}{\varepsilon(1-\varepsilon)},$$

and

$$\begin{aligned} (5) \quad \|C(z)\| &\leq \left(\sum_{k=0}^{\infty} \|V_0 S_0^{k+1} V_0 S_0\| |z - \lambda_0|^k \right) \left(\sum_{j=0}^{\infty} s^j |z - \lambda_0|^j \right) \\ &\leq r^2 / (1 - |z - \lambda_0|s)^2 = r^2 / (1 - \varepsilon)^2, \end{aligned}$$

as $\|V_0 S_0^{k+1} V_0 S_0\|/s^k \leq r^2$ for $k = 0, 1, 2, \dots$. Thus, it follows that for $z \in \Gamma_\varepsilon$

$$\|[V_0 R_0(z)]^2\| \leq \frac{p^2 s^2}{\varepsilon^2} + \frac{2psq}{\varepsilon(1-\varepsilon)} + \frac{r^2}{(1-\varepsilon)^2}.$$

Also, by the spectral radius formula,

$$r_\sigma(V_0 R_0(z)) = \inf_{n=1,2,\dots} \|[V_0 R_0(z)]^n\|^{1/n} \leq \|[V_0 R_0(z)]^2\|^{1/2}.$$

As the radius of $\partial(\Gamma_\varepsilon)$ is $\min_{z \in \Gamma_\varepsilon} 1/r_\sigma(V_0 R_0(z))$, we see that it must at least be

$$\left[\frac{p^2 s^2}{\varepsilon^2} + \frac{2psq}{\varepsilon(1-\varepsilon)} + \frac{r^2}{(1-\varepsilon)^2} \right]^{-1/2}.$$

Since for every ε in $(0, 1)$, the functions $P(t)$ and $\hat{\lambda}(t)$ are analytic on $\partial(\Gamma_\varepsilon)$, it follows that their power series converge if $|t|$ is less than m_0 , given by (4).

REMARK 2.2. Letting $\varepsilon = \sqrt{ps}/(\sqrt{ps} + \sqrt{q})$, we see that

$$(6) \quad (\sqrt{ps} + \sqrt{q})^{-1} [ps + 2\sqrt{psq} + r^2/q]^{-1/2} \leq m_0.$$

Since $r^2 \leq \|V_0 S_0\|^2 = q^2$, it follows that (6) improves upon the lower bound $(\sqrt{ps} + \sqrt{q})^{-2}$ given by Kato [2, (3.21)]. This bound was obtained by majorizing the series

$$V_0 R_0(z) = \frac{-V_0 P_0}{z - \lambda_0} + \sum_{k=0}^{\infty} V_0 S_0^{k+1} (z - \lambda_0)^k$$

term by term by the power series for the function

$$\Phi_1(z - \lambda_0) = \frac{p}{z - \lambda_0} + \frac{q}{1 - s(z - \lambda_0)}.$$

(See [2, (3.14)].) On the other hand, the lower bound m_0 given by (4) is obtained by majorizing the series for $[V_0 R_0(z)]^2$ by the power series for the function

$$\Phi_2(z - \lambda_0) = \frac{p}{(z - \lambda_0)^2} + \frac{2pq}{(z - \lambda_0)[1 - s(z - \lambda_0)]} + \frac{r^2}{[1 - s(z - \lambda_0)]^2}.$$

THEOREM 2.3. The convergence radii of the power series for $P(t)$ and $\hat{\lambda}(t)$ are at least equal to $1/4u$, where

$$(7) \quad u = \max\{ps, \sqrt{psq}, r\}.$$

If $|t| < 1/4u$, then the operator $T(t) = T_0 + tV_0$ has m eigenvalues $\lambda_1(t), \dots, \lambda_m(t)$, counted according to their algebraic multiplicities, and they satisfy

$$(8) \quad |\lambda_i(t) - \lambda_0| \leq \frac{1 - \sqrt{1 - 4u|t|}}{2s}, \quad 1 \leq i \leq m.$$

Further, $\lambda_1(t), \dots, \lambda_m(t)$ are the only spectral values of $T(t)$ lying in the open disk

$$(9) \quad \left\{ z \in \mathbb{C} : |z - \lambda_0| < \frac{1 + \sqrt{1 - 4u|t|}}{2s} \right\}.$$

PROOF. Let $0 < \varepsilon < 1$. Then by Lemma 2.1, the radius of the disk $\partial(\Gamma_\varepsilon)$ is at least

$$\left[\frac{u^2}{\varepsilon^2} + \frac{2u^2}{\varepsilon(1 - \varepsilon)} + \frac{u^2}{(1 - \varepsilon)^2} \right]^{-1/2} = \frac{\varepsilon(1 - \varepsilon)}{u}.$$

It can be easily seen that $|t| < \varepsilon(1 - \varepsilon)/u$ if and only if $u_1(t) < \varepsilon < u_2(t)$, where

$$(10) \quad u_1(t) = \frac{1 - \sqrt{1 - 4u|t|}}{2} \quad \text{and} \quad u_2(t) = \frac{1 + \sqrt{1 - 4u|t|}}{2}.$$

Hence for every ε satisfying $u_1(t) < \varepsilon < u_2(t)$, we see that $t \in \partial(\Gamma_\varepsilon)$. Thus, Γ_ε contains m eigenvalues $\lambda_1(t), \dots, \lambda_m(t)$ of $T(t)$ (counted according to their algebraic multiplicities) and does not contain any other spectral value of $T(t)$. In particular,

$$|\lambda_i(t) - \lambda_0| < \varepsilon/s, \quad i = 1, \dots, m.$$

Letting $\varepsilon \rightarrow u_1(t)$, we have

$$|\lambda_i(t) - \lambda_0| \leq \frac{1 - \sqrt{1 - 4u|t|}}{2s}, \quad i = 1, \dots, m,$$

while letting $\varepsilon \rightarrow u_2(t)$, we find that the open disk

$$\left\{ z \in \mathbb{C} : |z - \lambda_0| < \frac{1 + \sqrt{1 - 4u|t|}}{2s} \right\}$$

does not contain any other spectral value of $T(t)$.

REMARK 2.4. The functions $u_1(t)$ and $u_2(t)$ introduced in (10) satisfy $0 \leq u_1(t) \leq \frac{1}{2} \leq u_2(t) \leq 1$ for $|t| < 1/4u$, and they depend only on $|t|$. As $|t| \downarrow 0$, $u_1(t)$ monotonically decreases to 0, while $u_2(t)$ monotonically increases to 1, and as $|t| \uparrow 1/4u$, $u_1(t)$ monotonically increases to $\frac{1}{2}$, while $u_2(t)$ monotonically decreases to $\frac{1}{2}$. Thus, a smaller absolute value of the parameter t (corresponding to a smaller perturbation tV_0) yields a better estimate for $|\lambda_i(t) - \lambda_0|$, $i = 1, \dots, m$, and at the same time a larger disk of isolation of $\{\lambda_1(t), \dots, \lambda_m(t)\}$ from the rest of $\sigma(T(t))$. For all t with $|t| < 1/4u$, we have

$$\left\{ z \in \mathbb{C} : |z - \lambda_0| < \frac{1}{2s} \right\} \cap \sigma(T(t)) = \{\lambda_1(t), \dots, \lambda_m(t)\}.$$

These phenomena are illustrated in Figure 1.

REMARK 2.5. The lower bounds

$$(\sqrt{ps} + \sqrt{q})^{-1} \left[ps + 2\sqrt{psq} + \frac{r^2}{q} \right]^{-1/2} \quad \text{and} \quad \frac{1}{4 \max\{ps, \sqrt{psq}, r\}}$$

for the convergence radii of the series for $P(t)$ and $\hat{\lambda}(t)$ obtained in Remark 2.2 and Theorem 2.3 are significant in the operator approximation theory. We note that both these bounds tend to infinity if p and r approach 0,




Figure 1

and if s and q remain bounded. Let a compact operator T on X be approximated by a sequence of bounded operators (T_n) in a collectively compact manner (that is, $T_n x \rightarrow T x$ for every $x \in X$ and the set $\bigcup_{n=1}^{\infty} \{(T - T_n)x : x \in X, \|x\| \leq 1\}$ is totally bounded). Let λ be a simple eigenvalue of T such that $|\lambda| > \text{dist}(\lambda, \sigma(T) \setminus \{\lambda\})$. It is shown in [3, Theorem 2.1] and [5, Theorem 3.4] that for all large n , T_n has a simple eigenvalue λ_n such that $\lambda_n \rightarrow \lambda$, and if we let $T_0 = T_n$, $V_0 = T - T_n$, $\lambda_0 = \lambda_n$, $s_n = \|S_n\|$, $p_n = \|V_n P_n\|$, $q_n = \|V_n S_n\|$ and

$$r_n^2 = \sup\{\|V_n S_n^{k+1} V_n S_n\| / \|S_n\|^k : k = 0, 1, 2, \dots\},$$

then $p_n \rightarrow 0$ and $r_n \rightarrow 0$, while (s_n) and (q_n) remain bounded as $n \rightarrow \infty$. The above lower bounds then guarantee that for a suitably large fixed n , the point 1 lies in the convergence disks of $P(t)$ and $\hat{\lambda}(t)$, and hence the spectral projection $P(1)$ and the eigenvalue $\lambda = \lambda(1)$ of $T(1) = T$ can be approximated by the partial sums of the power series for $P(t)$ and $\hat{\lambda}(t)$, respectively.

We note that since (q_n) may not tend to zero as $n \rightarrow \infty$, the lower bound $(\sqrt{p_n s_n} + \sqrt{q_n})^{-2}$ for the convergence radii of $P(t)$ and for $\lambda(t)$ obtained by Kato is not useful for ascertaining the convergence at $t = 1$. In case (T_n) converges to T in the norm, then (q_n) does tend to zero as $n \rightarrow \infty$, and Kato's lower bound becomes applicable. We mention that there are many important cases where (T_n) approximates T in the collectively compact

manner, but not in the norm. For example, for a nonzero compact Fredholm integral operator T on $X = C([a, b])$, let (T_n) be a sequence of Nyström or Fredholm approximations of T , or if π_n is an interpolatory projection on X with $\pi_n x \rightarrow x$ for all $x \in X$, let $T_n = T\pi_n$ or $\pi_n T\pi_n$ ([1, pages 18–19] and [4, pages 292–294]).

In case T_0 is a normal operator on a Hilbert space X , then every eigenvalue of T_0 of finite algebraic multiplicity is semisimple. For this case, we refer to [4, Theorem 10.6] and [2, Theorem 3.9] for an especially elegant analogue of Theorem 2.3.

3. Convergence radius of an eigenvector series

Let λ_0 be a simple eigenvalue of T_0 , that is, λ_0 is a semisimple eigenvalue and the associated eigenspace is one dimensional. Then for $t \in \partial(\Gamma_0)$, $P(t)$ is of rank 1, and the operator $T(t) = T_0 + tV_0$ has only one simple eigenvalue $\lambda(t)$ lying inside Γ_0 , so $\hat{\lambda}(t) = \lambda(t)$ is analytic for $t \in \partial(\Gamma_0)$. Let ϕ_0 be an eigenvector of T_0 corresponding to λ_0 . Then the adjoint operator T_0^* is closed and densely defined in the adjoint space $X^* = \{x^*: X \rightarrow \mathbb{C}; x^* \text{ is conjugate linear and continuous}\}$. There is a unique eigenvector ϕ_0^* of T_0^* corresponding to $\bar{\lambda}_0$ such that $\langle \phi_0, \phi_0^* \rangle = 1$ and we have

$$P_0 x = \langle x, \phi_0^* \rangle \phi_0, \quad x \in X.$$

As mentioned in the introduction, the function

$$(11) \quad \phi(t) = P(t)\phi_0 / \langle P(t)\phi_0, \phi_0^* \rangle$$

is meromorphic in $\partial(\Gamma_0)$ and $\phi(0) = \phi_0$. If $t_0 \in \partial(\Gamma_0)$ is not a pole of the function ϕ , then $\phi(t_0)$ is an eigenvector of $T(t_0)$ corresponding to $\lambda(t_0)$. We have not been able to decide whether the function ϕ ever has a pole in $\partial(\Gamma_0)$. Nevertheless, we shall obtain a subregion of $\partial(\Gamma_0)$ which contains 0 and which is free of any pole of the function ϕ .

LEMMA 3.1. *Let $t_0 \in \partial(\Gamma_0)$. Then there is a unique nonnegative integer k_0 such that*

$$(12) \quad \psi(t_0) = \lim_{t \rightarrow t_0} P(t)\phi_0 / (t - t_0)^{k_0}$$

is an eigenvector of $T(t_0)$ corresponding to $\lambda(t_0)$. Further, t_0 is a pole of the function ϕ if and only if $\langle \psi(t_0), \phi_0^ \rangle = 0$, and in this case, $\psi(t_0)$ is an eigenvector also of the operator $(I - P_0)T(t_0)|_{(I - P_0)D_0}$.*

PROOF. Let $t_0 \in \partial(\Gamma_0)$. Since $P(0)\phi_0 = \phi_0 \neq 0$, the X -valued analytic function $t \mapsto P(t)\phi_0$ is not identically zero on $\partial(\Gamma_0)$. Hence either it has an isolated zero of finite order at t_0 or it is nonzero at t_0 . In any case, there is a unique nonnegative integer k_0 and an analytic function ψ in a neighbourhood U of t_0 such that $P(t)\phi_0 = (t - t_0)^{k_0}\psi(t)$ and $\psi(t_0) \neq 0$. For $t \in U$, we have

$$\langle P(t)\phi_0, \phi_0^* \rangle = (t - t_0)^{k_0} \langle \psi(t), \phi_0^* \rangle.$$

Hence the function

$$\phi(t) = \frac{\psi(t)}{\langle \psi(t), \phi_0^* \rangle}, \quad t \neq t_0,$$

has a pole at t_0 if and only if $\langle \psi(t_0), \phi_0^* \rangle = 0$. Clearly, $\lim_{t \rightarrow t_0} \psi(t)$ exists and equals $\psi(t_0)$. Now,

$$\psi(t) = P(t)\phi_0 / (t - t_0)^{k_0}, \quad t \in U, \quad t \neq t_0,$$

is an eigenvector of $T(t)$ corresponding to $\lambda(t)$. The continuity of the functions $t \mapsto P(t)$ and $t \mapsto \psi(t)$ at $t = t_0$ shows that $\psi(t_0)$ is an eigenvector of $T(t_0)$ corresponding to $\lambda(t_0)$.

Assume now that t_0 is a pole of the function ϕ . Then

$$P_0\psi(t_0) = \langle \psi(t_0), \phi_0^* \rangle \phi_0 = 0,$$

that is, $\psi(t_0) \in (I - P_0)D_0$. Since

$$(I - P_0)T(t_0)\psi(t_0) = \lambda(t_0)(I - P_0)\psi(t_0) = \lambda(t_0)\psi(t_0),$$

it is apparent that $\psi(t_0)$ is an eigenvector of $(I - P_0)T(t_0)|_{(I - P_0)D_0}$ corresponding to $\lambda(t_0)$.

PROPOSITION 3.2. *The meromorphic function ϕ given by (11) has no pole in the open set*

$$G = \{t \in \partial(\Gamma_0) : s|\lambda(t) - \lambda_0| + r|t| < 1\}.$$

For every $t \in G$, $\phi(t)$ is an eigenvector of $T(t)$ corresponding to $\lambda(t)$.

PROOF. Let $t \in G$. We show that $\lambda(t)$ is not in the spectrum of the operator $(I - P_0)(T_0 + tV_0)|_{(I - P_0)D_0} = (I - P_0)T(t)|_{(I - P_0)D_0}$. Lemma 3.1 then implies the desired result.

Let us denote $(I - P_0)D_0$ by Z and $\lambda(t)$ by z for brevity. Since $t \in G$, we have $|z - \lambda_0| < 1/s$. By (2), λ_0 is the only spectral value of T_0 in the open disk with centre λ_0 and radius $1/s$, and $\sigma(T_0|_Z) = \sigma(T_0) \setminus \{\lambda_0\}$. Hence

it follows that $z \notin \sigma(T_{0|Z})$. In order to conclude $z \notin \sigma(I - P_0)(T_0 + tV_0)|_Z$, it is enough to show that $r_\sigma(E(z)) < 1$, where

$$E(z) = [T_{0|Z} - (I - P_0)(T_0 + tV_0)|_Z][T_{0|Z} - zI_Z]^{-1}.$$

Now,

$$E(\lambda_0) = -t(I - P_0)V_0S_{0|Z},$$

and if $z \neq \lambda_0$, then

$$E(z) = -t(I - P_0)V_0R_0(z)|_Z.$$

In the first case,

$$\begin{aligned} r_\sigma(E(\lambda_0)) &\leq |t|r_\sigma((I - P_0)V_0S_0) = |t|r_\sigma(V_0S_0(I - P_0)) \\ &= |t|r_\sigma(V_0S_0) \leq |t| \|(V_0S_0)^2\|^{1/2} \leq |t|r, \end{aligned}$$

which is less than 1 since $t \in G$. In the second case, that is, when $z \neq \lambda_0$,

$$\begin{aligned} r_\sigma(E(z)) &\leq |t|r_\sigma((I - P_0)V_0R_0(z)) = |t|r_\sigma(V_0R_0(z)(I - P_0)) \\ &= |t| \|C(z)\|^{1/2} \leq |t|r/(1 - s|z - \lambda_0|), \end{aligned}$$

by (5). Since $z = \lambda(t)$ and $t \in G$, it again follows that $r_\sigma(E(z)) < 1$.

THEOREM 3.3. *Let $0 < \varepsilon < 1$ and $t \in \mathbb{C}$ be such that*

$$|t| < \left[\frac{p^2s^2}{\varepsilon^2} + \frac{2psq}{\varepsilon(1-\varepsilon)} + \frac{r^2}{(1-\varepsilon)^2} \right]^{-1/2}.$$

Then $\phi(t) = P(t)\phi_0 / \langle P(t)\phi_0, \phi_0^ \rangle$ is an eigenvector of $T(t)$ corresponding to $\lambda(t)$ such that $\langle \phi(t), \phi_0^* \rangle = 1$. The radius of convergence of the power series for $\phi(t)$ about 0 is at least equal to m_0 given by (4).*

PROOF. By Lemma 2.1, t belongs to $\partial(\Gamma_\varepsilon)$, so that $|\lambda(t) - \lambda_0| < \varepsilon/s$. Hence

$$s|\lambda(t) - \lambda_0| + r|t| < \varepsilon + \frac{r}{\left[\frac{p^2s^2}{\varepsilon^2} + \frac{2psq}{\varepsilon(1-\varepsilon)} + \frac{r^2}{(1-\varepsilon)^2} \right]^{1/2}} < 1.$$

By Proposition 3.2, we see that $\phi(t)$ is an eigenvector of $T(t)$ corresponding to $\lambda(t)$. It is obvious that $\langle \phi(t), \phi_0^* \rangle = 1$. Thus, the function ϕ is analytic on

$$\left\{ t \in \mathbb{C} : |t| < \left[\frac{p^2s^2}{\varepsilon^2} + \frac{2psq}{\varepsilon(1-\varepsilon)} + \frac{r^2}{(1-\varepsilon)^2} \right]^{-1/2} \right\}$$

for every ε , $0 < \varepsilon < 1$. The desired result now follows from the very definition of m_0 .

REMARK 3.4. As the quantities

$$(\sqrt{ps} + \sqrt{q})^{-1} \left[ps + 2\sqrt{psq} + \frac{r^2}{q} \right]^{-1/2} \quad \text{and} \quad \frac{1}{4 \max\{ps, \sqrt{psq}, r\}}$$

are at most equal to m_0 , they also give lower bounds for the convergence radius of the power series for $\phi(t)$, just as we had noted this for $P(t)$ and $\lambda(t)$. The lower bound $(\sqrt{ps} + \sqrt{q})^{-2}$ for $\phi(t)$ given in [2, Problem 3.7] (obtained by finding a majorizing series for $\phi(t)$) is thus improved to $(\sqrt{ps} + \sqrt{q})^{-1} [ps + 2\sqrt{psq} + r^2/q]^{1/2}$. (Note $r \leq q$.)

The lower bound $1/(4 \max\{ps, \sqrt{psq}, r\})$ for the convergence radii of the series for $\lambda(t)$ and for $\phi(t)$ is obtained in [5] by considering estimates for the iteratively defined n th terms of the power series for $\lambda(t)$ and $\phi(t)$, while in [7, pages 82 and 85], a similar lower bound is obtained by considering the power series for $P(t)$.

References

- [1] P. M. Anselone, *Collectively compact operator approximation theory*, (Prentice-Hall, Englewood Cliffs, N. J., 1971).
- [2] T. Kato, *Perturbation theory for linear operators*, 2nd ed. (Springer-Verlag, Berlin, Heidelberg, New York and Tokyo, 1976).
- [3] R. P. Kulkarni and B. V. Limaye, 'On the error estimates for the Rayleigh-Schrödinger series and the Kato-Rellich perturbation series', *J. Austral. Math. Soc. Ser. A* **46** (1989), 456–468.
- [4] B. V. Limaye, *Spectral perturbation and approximation with numerical experiments*, Proc. Centre Math. Anal., Vol. 13, (Australian National Univ., 1986).
- [5] B. V. Limaye and M. T. Nair, 'On the accuracy of Rayleigh-Schrödinger approximation,' *J. Math. Anal. Appl.* **139** (1989), 413–431.
- [6] M. T. Nair, 'A note on the Rayleigh-Schrödinger series', *J. Math. Phys. Sci.* **23** (1989), 185–193.
- [7] P. Redont, *Application de la théorie de la perturbation des opérateurs linéaires à l'obtention de bornes d'erreur sur les éléments propres et à leur calcul*, (Thèse Doct.-Ing., Univ. de Grenoble, 1979).

Indian Institute of Technology
Powai, Bombay 400076
India

University of Goa
Santa Cruz, Goa 403005
India