

## ERGODICITY AND DIFFERENCES OF FUNCTIONS ON SEMIGROUPS

BOLIS BASIT and A. J. PRYDE

(Received 18 July 1997)

Communicated by E. N. Dancer

### Abstract

Iseki [11] defined a general notion of ergodicity suitable for functions  $\varphi : J \rightarrow X$  where  $J$  is an arbitrary abelian semigroup and  $X$  is a Banach space. In this paper we develop the theory of such functions, showing in particular that it fits the general framework established by Eberlein [9] for ergodicity of semigroups of operators acting on  $X$ . Moreover, let  $\mathcal{A}$  be a translation invariant closed subspace of the space of all bounded functions from  $J$  to  $X$ . We prove that if  $\mathcal{A}$  contains the constant functions and  $\varphi$  is an ergodic function whose differences lie in  $\mathcal{A}$  then  $\varphi \in \mathcal{A}$ . This result has applications to spaces of sequences facilitating new proofs of theorems of Gelfand and Katznelson-Tzafriri [12]. We also obtain a decomposition for the space of ergodic vectors of a representation  $T : J \rightarrow L(X)$  generalizing results known for the case  $J = \mathbb{Z}^+$ . Finally, when  $J$  is a subsemigroup of a locally compact abelian group  $G$ , we compare the Iseki integrals with the better known Cesàro integrals.

1991 *Mathematics subject classification* (Amer. Math. Soc.): primary 43A60; secondary 47A10, 47D03, 28B05.

*Keywords and phrases*: Ergodic, semigroup, differences, Beurling spectrum, invariant means, system of invariant integrals.

### 1. Introduction

In a successful attempt to unify and extend the growing collection of ergodic theorems, Eberlein [9] introduced systems of almost invariant integrals for semigroups of continuous linear transformations on locally convex spaces. A semigroup possessing such a system he called ergodic, and for such semigroups he proved a very general mean ergodic theorem ([9, Theorem 3.1]). Since that time many more ergodic theorems have appeared and many have been revealed as special cases of Eberlein's classical theorem. See for example [17].

In a different direction, Iseki [11] introduced the notion of ergodicity of functions  $\varphi : J \rightarrow X$  where  $J$  is a semigroup and  $X$  is a locally convex space. With it he was able to show that every such function which is almost periodic in the sense of Maak is necessarily ergodic.

Ruess and Summers [18] considered asymptotically almost periodic functions  $\varphi : \mathbb{R}^+ \rightarrow X$ . They showed that if the indefinite integral  $\Phi$  of  $\varphi$  is weakly almost periodic in the sense of Eberlein, then  $\Phi$  is asymptotically almost periodic. Subsequently Basit [3] observed that weak almost periodicity could be replaced by the more general property of ergodicity, that is the Cesàro integrals of  $\Phi$  converge uniformly to a constant. Moreover, he replaced asymptotically almost periodic functions by large classes of functions. Ruess and Phóng [16] independently obtained some of these results.

Basit also observed that the integral problem discussed above is closely related to the *difference problem*: if  $\varphi \in C_b(J, X)$  and  $\Delta_t \varphi \in \mathcal{A} \subseteq C_b(J, X)$  for all  $t \in J$ , find conditions that ensure  $\varphi \in \mathcal{A}$ . Basit investigated this problem for the cases  $J = \mathbb{R}^+$  or  $\mathbb{R}$  and gave applications to the solutions of certain integro-differential difference equations [3] and to the abstract Cauchy problem [4]. Once again ergodicity of  $\varphi$  played an important role.

In the present paper we develop the theory of (Iseki) ergodic functions  $\varphi : J \rightarrow X$  where  $J$  is an arbitrary semigroup and  $X$  is a Banach space. For the sake of simplicity and clarity, we restrict ourselves to the case of abelian  $J$ . In particular, we show how this theory fits into the framework established by Eberlein. Our main result concerns the difference problem and its relationship with ergodicity. This is in Section 2.

In Section 3 we apply our results to spaces of sequences. Among other things we obtain new proofs of theorems of Gelfand and Katznelson-Tzafriri on power bounded elements of Banach algebras. Section 4 deals with representations of semigroups on Banach spaces. We obtain a decomposition for the subspace of ergodic vectors generalizing known results for the case  $J = \mathbb{Z}^+$ .

Finally, in section 5 we exhibit a large class of semigroups  $J$  for which one can take limits of Cesàro integrals of functions  $\varphi$  in  $C_{ub}(J, X)$ . We show that these limits, when they exist, are identical to the Iseki means. Similarly, when  $G$  is a locally compact abelian group, we show that the means studied by Argabright [2] and Datry and Muraz [7] for  $\varphi \in C_b(G, X)$  are identical to the Iseki means. We conclude by giving a simple condition on the Beurling spectrum of a function  $\varphi \in C_{ub}(G, X)$  that ensures  $\varphi$  is ergodic.

## 2. Ergodicity

Throughout this paper,  $J$  will denote an abelian semigroup and  $X$  a Banach space over  $\mathbb{R}$  or  $\mathbb{C}$ . By  $B(J, X)$  we denote the space of bounded functions  $\varphi : J \rightarrow X$ , endowed with the norm  $\|\varphi\|_\infty = \sup_{t \in J} \|\varphi(t)\|$ . For such a function,  $\varphi_s$  and  $\Delta_s \varphi$  will denote the translate and difference by  $s$  of  $\varphi$ , defined by  $\varphi_s(t) = \varphi(t + s)$  and  $\Delta_s \varphi = \varphi_s - \varphi$  for  $s, t \in J$ . The closed subspaces of  $B(J, X)$  consisting of continuous and uniformly continuous functions respectively are denoted  $C_b(J, X)$  and  $C_{ub}(J, X)$ . We will use the same symbol, say  $x$ , for an element of  $X$  and for the function in  $B(J, X)$  taking the constant value  $x$ .

Following Iseki [11, I] we say that a function  $\varphi : J \rightarrow X$  is *ergodic* if  $\varphi \in B(J, X)$  and there exists  $M_\varphi \in X$  such that for each  $\varepsilon > 0$  there are elements  $t_1, \dots, t_n \in J$  with  $\|(1/n) \sum_{i=1}^n (\varphi_{t_i} - M_\varphi)\|_\infty < \varepsilon$ . The element  $M_\varphi$ , clearly unique, is called the (Iseki) *mean* of  $\varphi$  and the class of all such ergodic functions is denoted  $E(J, X)$ . We define  $M : E(J, X) \rightarrow X$  by  $M(\varphi) = M_\varphi$ .

**PROPOSITION 2.1.** *The space  $E(J, X)$  is a translation invariant closed subspace of  $B(J, X)$  containing all the constant functions. Moreover,  $M : E(J, X) \rightarrow X$  is a bounded linear map.*

**PROOF.** Let  $\varphi, \psi \in E(J, X)$ . By the definition of ergodicity, for each  $\varepsilon > 0$  there exist elements  $s_1, \dots, s_m, t_1, \dots, t_n \in J$  such that  $\|(1/m) \sum_{i=1}^m (\varphi_{s_i} - M_\varphi)\|_\infty < \varepsilon$  and  $\|(1/n) \sum_{j=1}^n (\psi_{t_j} - M_\psi)\|_\infty < \varepsilon$ . Since  $\|\varphi_t\|_\infty \leq \|\varphi\|_\infty$  for all  $t \in J$ , we obtain  $\|(1/nm) \sum_{i=1}^m \sum_{j=1}^n (\varphi_{s_i+t_j} + \psi_{s_i+t_j} - M_\varphi - M_\psi)\|_\infty < 2\varepsilon$ . Hence  $\varphi + \psi \in E(J, X)$  and  $M(\varphi + \psi) = M(\varphi) + M(\psi)$ . The rest of the proposition is proved similarly.

The following result shows that there are many ergodic functions. Further examples will be provided later.

**PROPOSITION 2.2.** *If  $\varphi \in B(J, X)$  and  $s \in J$  then  $\Delta_s \varphi \in E(J, X)$  and  $M(\Delta_s \varphi) = 0$ .*

**PROOF.** Given  $\varepsilon > 0$ , choose  $n \in \mathbb{N}$  such that  $\|(1/n)\varphi\|_\infty < \varepsilon/2$ . Since  $(\Delta_s \varphi)_t = \Delta_{s+t} \varphi - \Delta_t \varphi$ , we have  $\|(1/n) \sum_{j=1}^n (\Delta_s \varphi)_{t_j}\|_\infty < \varepsilon$ . This proves the proposition.

The following alternative characterization of ergodic functions will be useful. For this we set  $\mathcal{F}(J) = \{F \subseteq J : |F| < \infty\}$  where  $|F|$  is the cardinality of  $F$ . Then  $\mathcal{F}(J)$  becomes a directed set if we define  $F_1 \leq F_2$  whenever there exists  $F \in \mathcal{F}(J)$  such that  $F_2 = F_1 + F$ .

**PROPOSITION 2.3.** *Let  $\varphi \in B(J, X)$ . Then  $\varphi \in E(J, X)$  if and only if there exists  $y \in X$  such that  $\lim_{F \in \mathcal{F}(J)} ((1/|F|) \sum_{t \in F} \varphi_t) = y$ . In this case,  $y = M_\varphi$ .*

PROOF. Let  $\varphi \in E(J, X)$ . For each  $\varepsilon > 0$  there is a set  $F_\varepsilon \in \mathcal{F}(J)$  such that  $\|(1/|F_\varepsilon|) \sum_{t \in F_\varepsilon} (\varphi_t - M_\varphi)\|_\infty < \varepsilon$ . If  $F \in \mathcal{F}(J)$  satisfies  $F \geq F_\varepsilon$ , that is  $F = F_\varepsilon + H$  for some  $H \in \mathcal{F}(J)$ , then

$$\left\| \frac{1}{|F|} \sum_{u \in F} (\varphi_u - M_\varphi) \right\|_\infty = \left\| \frac{1}{|F_\varepsilon|} \cdot \frac{1}{|H|} \sum_{t \in F_\varepsilon} \sum_{s \in H} (\varphi_{t+s} - M_\varphi) \right\|_\infty < \varepsilon,$$

showing that  $\lim_{F \in \mathcal{F}(J)} (1/|F|) \sum_{t \in F} \varphi_t = M_\varphi$ . The converse is clear.

Our next task is to set Iseki ergodicity in the framework of Eberlein. For this, let  $\mathcal{S}$  be a sub-semigroup under composition of the Banach algebra  $L(E)$  of all bounded operators  $A : E \rightarrow E$  where  $E$  is a Banach space. The orbit of  $x \in E$  under  $\mathcal{S}$  is  $\text{orb}_{\mathcal{S}}(x) = \{Sx : S \in \mathcal{S}\}$ . A net  $(A_\alpha)_{\alpha \in \Lambda}$  in  $L(E)$  is called a *system of invariant integrals* for  $\mathcal{S}$  if

$$(2.1) \quad A_\alpha x \in \overline{\text{co}} \text{orb}_{\mathcal{S}}(x) \text{ for all } x \in E \text{ and } \alpha \in \Lambda,$$

$$(2.2) \quad \sup_{\alpha \in \Lambda} \|A_\alpha\| < \infty,$$

$$(2.3) \quad \lim_{\alpha \in \Lambda} \|(A_\alpha S - A_\alpha)x\| = \lim_{\alpha \in \Lambda} \|(SA_\alpha - A_\alpha)x\| = 0 \text{ for all } x \in E \text{ and } S \in \mathcal{S}.$$

If (2.1), (2.2) hold but (2.3) only holds at  $x_0 \in E$  then we say  $(A_\alpha)$  is a *system of invariant integrals for  $\mathcal{S}$  at  $x_0$* .

For  $\varphi \in B(J, X)$ ,  $F \in \mathcal{F}(J)$  and  $s \in J$ , define  $R_F \varphi = (1/|F|) \sum_{t \in F} \varphi_t$ , interpreted as 0 if  $F = \emptyset$ , and  $R_s = R_{\{s\}}$ . Hence  $R_F, R_s \in L(E)$  where  $E = B(J, X)$ .

PROPOSITION 2.4. *The net  $(R_F)_{F \in \mathcal{F}(J)}$  is a system of invariant integrals for the translation semigroup  $\mathcal{R} = \{R_s : s \in J\}$ .*

PROOF. For  $\varphi \in B(J, X)$ ,  $(R_F R_s - R_F)\varphi = R_F(\Delta_s \varphi)$ . By Proposition 2.2,  $M(\Delta_s \varphi) = 0$  and so by Proposition 2.3,  $\lim_{F \in \mathcal{F}(J)} (R_F R_s - R_F)\varphi = 0$ . Hence (2.3) follows, and (2.1), (2.2) are obvious.

By Eberlein's mean ergodic theorem [9, Theorem 3.1] we have immediately

COROLLARY 2.5. *For  $\varphi \in B(J, X)$  the following are equivalent*

- (1)  $\varphi \in E(J, X)$  and  $M(\varphi) = y$ ,
- (2) the net  $(R_F \varphi)_{F \in \mathcal{F}(J)}$  converges to  $y$ ,
- (3) some subnet of  $(R_F \varphi)_{F \in \mathcal{F}(J)}$  converges weakly to  $y$ ,
- (4)  $y \in \overline{\text{co}} \text{orb}_{\mathcal{R}}(\varphi)$  with  $y$  a constant function.

Recall that the space  $W(J, X)$  of Eberlein weakly almost periodic functions consists of the bounded functions  $\varphi : J \rightarrow X$  for which  $\text{orb}_{\mathcal{A}}(\varphi)$  is weakly relatively compact. From Corollary 2.5 we obtain

**COROLLARY 2.6.**  $W(J, X)$  is a closed linear subspace of  $E(J, X)$ .

Note that  $M : E(J, X) \rightarrow X$  is a (translation) invariant mean in the sense of [6, p.79] for scalar  $X$  and [21] for general  $X$ . The latter proved the existence of an invariant mean on  $W(J, X)$  for certain non-abelian semigroups  $J$  [21, Theorem 8.7]. However, the invariant means in these references are not given explicitly.

To conclude this section we prove our main result for ergodic functions. With the additional assumption that  $\mathcal{A}$  contains the constant functions, this theorem provides a solution of the difference problem.

**THEOREM 2.7.** Let  $\mathcal{A}$  be a translation invariant closed subspace of  $B(J, X)$ . If  $\varphi \in E(J, X)$  and  $\Delta_t \varphi \in \mathcal{A}$  for all  $t \in J$ , then  $\varphi - M(\varphi) \in \mathcal{A}$ .

**PROOF.** For each non-empty  $F \in \mathcal{F}(J)$  we have  $\varphi - R_F \varphi = -(1/|F|) \sum_{t \in F} \Delta_t \varphi \in \mathcal{A}$ . The theorem follows from Corollary 2.5 by taking the limit over  $F$  in  $\mathcal{F}(J)$ .

### 3. Sequence spaces

In this section we give some applications of our results to spaces of sequences. Here we take  $J = \mathbb{Z}, \mathbb{Z}^+$  or  $\mathbb{Z}^-$  and use the condition

$$(3.1) \quad \mathcal{A} \text{ is a closed subspace of } B(J, X) \text{ such that } \psi_t|_J \in \mathcal{A} \text{ whenever } \psi \in B(\mathbb{Z}, X), \\ t \in \mathbb{Z} \text{ and } \psi|_J \in \mathcal{A}.$$

Examples of such subspaces  $\mathcal{A}$  include  $E(J, X)$ , the space  $C_0(J, X)$  of functions convergent to 0 at infinity, the space  $AP(\mathbb{Z}, X)$  of almost periodic functions and the space  $WAP(J, X)$  of Eberlein weakly almost periodic functions.

Following [3, Definition 4.1.2] we define the *spectrum with respect to*  $\mathcal{A}$  of a function  $\varphi \in B(\mathbb{Z}, X)$  by  $\text{sp}_{\mathcal{A}}(\varphi) = \{\gamma \in \widehat{\mathbb{Z}} : \hat{f}(\gamma) = 0 \text{ for all } f \in I_{\mathcal{A}}(\varphi)\}$  where  $\widehat{\mathbb{Z}}$  is the (unitary) character group of  $\mathbb{Z}$ ,  $\hat{f} : \widehat{\mathbb{Z}} \rightarrow \mathbb{C}$  is the Fourier transform of  $f$ , and  $I_{\mathcal{A}}(\varphi) = \{f \in L^1(\mathbb{Z}) : (\varphi * f)|_J \in \mathcal{A}\}$ .

The following proposition is well-known for the case  $\mathcal{A} = \{0\}$  and  $J = \mathbb{Z}$ , in which case  $\text{sp}_{\mathcal{A}}(\varphi) = \text{sp}(\varphi)$ , the *Beurling spectrum* of  $\varphi$ .

**PROPOSITION 3.1.** Let  $\varphi, \psi \in B(\mathbb{Z}, X)$ ,  $f \in L^1(\mathbb{Z})$ ,  $\gamma \in \widehat{\mathbb{Z}}$  and  $\mathcal{A}$  satisfy condition (3.1).

- (i)  $\text{sp}_{\mathcal{A}}(\varphi) = \text{sp}_{\mathcal{A}}(\varphi_t)$  for all  $t \in \mathbb{Z}$ .

- (ii)  $\text{sp}_{\mathcal{A}}(\varphi * f) \subseteq \text{sp}_{\mathcal{A}}(\varphi) \cap \text{supp}(f)$ .
- (iii)  $\text{sp}_{\mathcal{A}}(\varphi + \psi) \subseteq \text{sp}_{\mathcal{A}}(\varphi) \cup \text{sp}_{\mathcal{A}}(\psi)$ .
- (iv)  $\text{sp}_{\mathcal{A}}(\gamma\varphi) = \gamma + \text{sp}_{\mathcal{A}}(\varphi)$ .
- (v)  $\text{sp}_{\mathcal{A}}(\varphi) = \emptyset$  if and only if  $\varphi|_J \in \mathcal{A}$ .

PROOF. The arguments are the same as for the Beurling spectrum. See for example [8, part II, p.988] or [5]. We present a proof for (v). If  $\varphi|_J \in \mathcal{A}$  then by (3.1),  $\varphi_t|_J \in \mathcal{A}$  for all  $t \in \mathbb{Z}$ . Hence for  $f \in L^1(\mathbb{Z})$ ,  $(\varphi * f)|_J = \sum_{n \in \mathbb{Z}} f(n)\varphi_{-n}|_J \in \mathcal{A}$ . So  $I_{\mathcal{A}}(\varphi) = L^1(\mathbb{Z})$  and  $\text{sp}_{\mathcal{A}}(\varphi) = \emptyset$ . Conversely, if  $\text{sp}_{\mathcal{A}}(\varphi) = \emptyset$  then  $I_{\mathcal{A}}(\varphi) = L^1(\mathbb{Z})$ . Choose  $f_n \in L^1(\mathbb{Z})$  such that  $\varphi * f_n \rightarrow \varphi$  in  $B(\mathbb{Z}, X)$ . Since  $f_n \in I_{\mathcal{A}}(\varphi)$ ,  $(\varphi * f_n)|_J \in \mathcal{A}$  and since  $\mathcal{A}$  is closed,  $\varphi|_J \in \mathcal{A}$ .

In the sequel we denote the elements of  $\widehat{\mathbb{Z}}$  by  $\gamma_\lambda$  or  $\lambda$ , where  $\lambda \in \mathbb{T}$  the circle group and  $\gamma_\lambda(n) = \lambda^n$  for  $n \in \mathbb{Z}$ . Hence  $\gamma_1$  or 1 is the unit in  $\widehat{\mathbb{Z}}$ .

PROPOSITION 3.2. *Suppose  $\mathcal{A}$  satisfies (3.1),  $\varphi \in B(J, X)$ ,  $\varphi|_J \in E(J, X)$  and  $\text{sp}_{\mathcal{A}}(\varphi) \subseteq \{1\}$ . Then  $\varphi|_J - M(\varphi|_J) \in \mathcal{A}$ .*

PROOF. By Wiener's tauberian theorem [15, 7.2.5] the condition  $\text{sp}_{\mathcal{A}}(\varphi) \subseteq \{1\}$  is equivalent to  $I_{\mathcal{A}}(\varphi) \supseteq \{f \in L^1(\mathbb{Z}) : \hat{f}(1) = 0\}$ . For  $t \in \mathbb{Z}$ ,  $g \in L^1(\mathbb{Z})$  and  $\lambda \in \mathbb{T}$  we have  $(\Delta_t g)\hat{\gamma}(\lambda) = (\gamma_\lambda(t) - 1)\hat{g}(\lambda)$ . Hence  $\Delta_t g \in I_{\mathcal{A}}(\varphi)$ . In other words,  $(\Delta_t \varphi * g)|_J = (\varphi * \Delta_t g)|_J \in \mathcal{A}$ . Setting  $g = \chi_{\{0\}}$ , the characteristic function of  $\{0\}$  in  $\mathbb{Z}$  we have  $\Delta_t \varphi = \Delta_t \varphi * g$  and so  $\Delta_t \varphi|_J \in \mathcal{A}$ . By Theorem 2.7,  $\varphi|_J - M(\varphi|_J) \in \mathcal{A}$ .

As a consequence we have the following application of spectra to the difference problem.

THEOREM 3.3. *Suppose  $\mathcal{A}$  satisfies (3.1) and  $\varphi \in B(\mathbb{Z}, X)$ . Then  $\text{sp}_{\mathcal{A}}(\varphi) \subseteq \{1\}$  if and only if  $\Delta_t \varphi|_J \in \mathcal{A}$  for all  $t \in J$ .*

PROOF. Let  $\Delta_t \varphi|_J \in \mathcal{A}$  for all  $t \in J$ . If  $g \in L^1(\mathbb{Z})$  then by (3.1),  $(\varphi * \Delta_t g)|_J = \sum_{n \in \mathbb{Z}} g(n)(\Delta_t \varphi)_{-n}|_J \in \mathcal{A}$ . So  $I_{\mathcal{A}}(\varphi) \supseteq \{\Delta_t g : t \in J, g \in L^1(\mathbb{Z})\}$ . But  $(\Delta_t g)\hat{\gamma}(\lambda) = (\gamma_\lambda(t) - 1)\hat{g}(\lambda)$  is zero for all  $t \in J$  and  $g \in L^1(\mathbb{Z})$  only when  $\lambda = 1$ . So  $\text{sp}_{\mathcal{A}}(\varphi) \subseteq \{1\}$ . Conversely, let  $\text{sp}_{\mathcal{A}}(\varphi) \subseteq \{1\}$ . By Proposition 2.2,  $\Delta_t \varphi|_J \in E(J, X)$  and  $M(\Delta_t \varphi|_J) = 0$  for each  $t \in J$ . By Proposition 3.2,  $\Delta_t \varphi|_J \in \mathcal{A}$ .

In order to apply Theorem 3.3, we first prove the following result. In it,  $\sigma(x)$  denotes the Banach algebra spectrum of  $x$ .

THEOREM 3.4. *Let  $X$  be a unital Banach algebra. Suppose  $\mathcal{A} \subseteq B(J, X)$  satisfies (3.1) and in addition  $y\mathcal{A} \subseteq \mathcal{A}$  for all  $y \in X$ . Let  $\varphi : \mathbb{Z} \rightarrow X$  be a bounded solution of the recurrence equation  $\varphi(n+1) = x\varphi(n) + \psi(n)$  for some  $x \in X$  and  $\psi \in C_b(\mathbb{Z}, X)$ . If  $\psi|_J \in \mathcal{A}$  then  $\text{sp}_{\mathcal{A}}(\varphi) \subseteq \sigma(x) \cap \mathbb{T}$ .*

PROOF. Let  $\lambda_0 \in \mathbb{T} \setminus \sigma(x)$ . Choose  $\delta > 0$  such that  $B_\delta(\lambda_0) = \{\lambda \in \mathbb{C} : |\lambda - \lambda_0| < \delta\} \subseteq \mathbb{C} \setminus \sigma(x)$ . Take  $f \in L^1(\mathbb{Z})$  with  $\hat{f}(\lambda_0) = 1$  and  $\text{supp}(\hat{f}) \subseteq B_{\delta/4}(\lambda_0)$ . Let  $\xi = \varphi * f$ . It suffices to prove  $\xi|_J \in \mathcal{A}$ , for then  $f \in I_{\mathcal{A}}(\varphi)$  and  $\lambda_0 \notin \text{sp}_{\mathcal{A}}(\varphi)$ .

To do this, let  $g \in L^1(\mathbb{Z})$  be such that  $\hat{g}(\lambda) = 1$  for  $\lambda \in B_{\delta/2}(\lambda_0) \cap \mathbb{T}$ ,  $\text{supp}(\hat{g}) \subseteq B_\delta(\lambda_0)$  and  $\hat{g} \in C^1(\mathbb{T})$ . Define  $h : \mathbb{T} \rightarrow X$  by  $\hat{h}(\lambda) = \hat{g}(\lambda)(\lambda e - x)^{-1}$ , interpreted as 0 outside  $B_\delta(\lambda_0)$ , where  $e$  is the unit in  $X$ . Then  $\hat{h} \in C^1(\mathbb{T}, X)$  so  $\hat{h}(\lambda) = \sum_{n=-\infty}^{\infty} h(n)\lambda^{-n}$  for some  $h \in L^1(\mathbb{Z}, X)$  with  $h(n)x = xh(n)$  for all  $n \in \mathbb{Z}$ . Moreover, if  $\eta_\lambda(n) = \gamma_\lambda(n+1)e - \gamma_\lambda(n)x$ , where  $\gamma_\lambda(n) = \lambda^n$  and  $\lambda \in B_{\delta/2}(\lambda_0) \cap \mathbb{T}$ , then  $h * \eta_\lambda = \gamma_\lambda$ . Indeed,

$$\begin{aligned} h * \eta_\lambda(n) &= \sum_j h(j)(\lambda^{n+1-j}e - \lambda^{n-j}x) = \lambda^n(\lambda e - x) \sum_j h(j)\lambda^{-j} \\ &= \lambda^n(\lambda e - x)\hat{g}(\lambda)(\lambda e - x)^{-1} = \lambda^n. \end{aligned}$$

Now  $\xi = \varphi * f \in B(\mathbb{Z}, X)$  and  $\text{sp}(\xi) \subseteq \text{supp}(\hat{f}) \subseteq B_{\delta/4}(\lambda_0)$ , so there is a sequence of trigonometric polynomials  $\pi_m \in B(\mathbb{Z}, X)$  converging pointwise to  $\xi$  and with  $\text{sp}(\pi_m) \subseteq B_{\delta/2}(\lambda_0)$ . Let  $\eta_m(n) = \pi_m(n+1)e - x\pi_m(n)$ . Then  $h * \eta_m = \pi_m$ .

From the recurrence equation,  $\eta_m(n) \rightarrow \xi(n+1) - x\xi(n) = \psi * f(n)$  for each  $n \in \mathbb{Z}$ . Hence  $\xi = h * \psi * f$ . Since  $\xi = \sum_{n \in \mathbb{Z}} h(n)(\psi * f)_{-n}$  and  $y_{\mathcal{A}} \subseteq \mathcal{A}$  for each  $y \in X$ , it follows from (3.1) that  $\xi|_J \in \mathcal{A}$  as required.

As a consequence we easily obtain the following two results. The first was proved by Gelfand (see [12]) and the second by Katznelson and Tzafriri [12]. Recall that an element  $x$  of a unital Banach algebra  $X$  is called *power bounded* if  $\{x^n : n \in \mathbb{Z}^+\}$  is bounded and *doubly power bounded* if  $\{x^n : n \in \mathbb{Z}\}$  is bounded.

**COROLLARY 3.5.** *Let  $x$  be a doubly power bounded element of a unital Banach algebra  $X$ . If  $\sigma(x) = \{1\}$  then  $x = e$ .*

PROOF. We may apply Theorem 3.4 with  $\mathcal{A} = \{0\}$ ,  $J = \mathbb{Z}$ ,  $\psi = 0$  and  $\varphi(n) = x^n$ . So  $\text{sp}(\varphi) \subseteq \sigma(x) \cap \mathbb{T} = \{1\}$ . By Theorem 3.3,  $\Delta_t \varphi = 0$  for all  $t \in \mathbb{Z}$  and hence  $x = e$ .

**COROLLARY 3.6.** *Let  $x$  be a power bounded element of a unital Banach algebra  $X$ . If  $\sigma(x) \cap \mathbb{T} \subseteq \{1\}$  then  $\|x^{n+1} - x^n\| \rightarrow 0$  as  $n \rightarrow \infty$ .*

PROOF. Apply Theorem 3.4 with  $\mathcal{A} = C_0(J, X)$ ,  $J = \mathbb{Z}^+$  and  $\varphi, \psi$  as follows. For  $n \geq 0$  set  $\varphi(n) = x^n$ ,  $\psi(n) = 0$  and for  $n < 0$  set  $\varphi(n) = e$ ,  $\psi(n) = e - x$ . So  $\text{sp}_{\mathcal{A}}(\varphi) \subseteq \{1\}$  and by Theorem 3.3,  $\Delta_t \varphi|_J \in \mathcal{A}$  for all  $t \in J$ . This gives the corollary.

In a subsequent paper we will use ergodicity and the difference problem to obtain generalizations of these last two results.

#### 4. Ergodic vectors of representations

Throughout this section  $J$  will denote an abelian semigroup and  $T : J \rightarrow L(X)$  a representation. That is,  $T$  is a semigroup homomorphism mapping  $J$  into the semigroup under composition  $L(X)$ . The dual representation  $T^* : J \rightarrow L(X^*)$  is defined by  $\langle x, T^*(t)\varphi \rangle = \langle T(t)x, \varphi \rangle$  for  $x \in X$ ,  $t \in J$  and  $\varphi \in X^*$ .

The space of fixed points of  $T$  is  $N = N(T) = \bigcap_{t \in J} \ker(T(t) - I)$  and its complementary space is  $R = R(T) = \text{span}\{T(s)x - x : x \in X, s \in J\}$ . The closure of  $R$  is denoted  $\bar{R} = \bar{R}(T)$ . The set of ergodic vectors of  $T$  is  $X_{\text{erg}} = X_{\text{erg}}(T) = \{x \in X : T(\cdot)x \in E(J, X)\}$ .

Next let  $T(J)$  be the range of  $T$  in  $L(X)$  and for  $F \in \mathcal{F}(J)$  define  $T_F \in L(X)$  by  $T_F x = (1/|F|) \sum_{t \in F} T(t)x$ , again interpreted as 0 if  $F = \emptyset$ . Finally, the orbit under  $T$  of an element  $x \in X$  is  $\text{orb}_T(x) = \text{orb}_{T(J)}(x)$ .

**PROPOSITION 4.1.** *If  $T : J \rightarrow L(X)$  is a representation and  $\text{orb}_T(x)$  is bounded for some  $x \in X$ , then the set  $(T_F)_{F \in \mathcal{F}(J)}$  is a system of invariant integrals for the semigroup  $T(J)$  at  $x$ .*

**PROOF.** Let  $s \in J$ . The function  $T(\cdot)x : J \rightarrow X$  is bounded and hence by Proposition 2.2,  $\Delta_s T(\cdot)x \in E(J, X)$  and  $M(\Delta_s T(\cdot)x) = 0$ . By Corollary 2.5,  $\lim_F R_F \Delta_s T(\cdot)x = 0$  and in particular  $\lim_F \|R_F \Delta_s T(t)x\| = 0$  for each  $t \in J$ . But  $R_F \Delta_s T(t)x = (R_{F+1} T(s) - R_{F+1})x$  and so  $\lim_F \|(R_F T(s) - R_F)x\| = 0$ . Condition (2.3) follows for this  $x$ . Since (2.1) and (2.2) are clear the proposition is proved.

**COROLLARY 4.2.** *If  $T : J \rightarrow L(X)$  is a representation and  $\text{orb}_T(x)$  is bounded for some  $x \in X$  then the following are equivalent*

- (i)  $x \in X_{\text{erg}}(T)$  and  $M(T(\cdot)x) = y$ ,
- (ii)  $(T_F x)_{F \in \mathcal{F}(J)}$  converges to  $y$ ,
- (iii) some subnet of  $(T_F x)_{F \in \mathcal{F}(J)}$  converges weakly to  $y$ ,
- (iv)  $y \in N(T) \cap \overline{\text{co}} \text{orb}_T(x)$ .

**PROOF.** By Eberlein's mean ergodic theorem (see Theorem 3.1 in [9] and the remark following it) we conclude that (ii), (iii) and (iv) are equivalent. Let  $\kappa = \sup\{\|z\| : z \in \text{orb}_T(x)\}$ . Then for each  $t \in J$  and  $F \in \mathcal{F}(J)$  we have  $\|T_{F+1}x - y\| = \|R_F T(t)x - y\| \leq \|R_F T(\cdot)x - y\|_\infty \leq \kappa \|T_F x - y\|$ . Hence  $(T_F x) \rightarrow y$  in  $X$  if and only if  $(R_F T(\cdot)x) \rightarrow y$  in  $B(J, X)$ . By Corollary 2.5, (ii) is equivalent to (i).

**PROPOSITION 4.3.** *If  $T : J \rightarrow L(X)$  is a bounded representation, then  $X_{\text{erg}}$  is a closed linear subspace of  $X$ . Moreover,  $X_{\text{erg}} = N \oplus \bar{R}$ .*



PROOF. Since  $E(J, X)$  is a linear space, so too is  $X_{\text{erg}}$ . The closedness of  $X_{\text{erg}}$  follows from the boundedness of  $T$  and the closedness of  $E(J, X)$  in  $B(J, X)$ . If  $x \in N$  then  $T(t)x = x$  for all  $t \in J$ . Hence  $T(\cdot)x \in E(J, X)$  and  $M(T(\cdot)x) = x$ , showing  $N \subseteq X_{\text{erg}}$ . If  $z \in R$  then there exist  $t_1, \dots, t_n \in J$  and  $x_1, \dots, x_n \in X$  such that  $z = \sum_{j=1}^n (T(t_j)x_j - x_j)$ . Hence  $T(\cdot)z = \sum_{j=1}^n \Delta_{t_j} T(\cdot)x_j$ . By Proposition 2.2,  $T(\cdot)z \in E(J, X)$  and  $M(T(\cdot)z) = 0$ . By Proposition 2.1, the same is true for  $z \in \bar{R}$ . Hence  $\bar{R} \subseteq X_{\text{erg}}$  and moreover,  $N \cap \bar{R} = \{0\}$ .

Finally we show  $X_{\text{erg}} \subseteq N + \bar{R}$ . If  $y \in X_{\text{erg}}$  then by Corollary 4.2,  $M(T(\cdot)y) \in N$ . Setting  $z = y - M(T(\cdot)y)$  we show  $z \in \bar{R}$ . Indeed, for each  $\varepsilon > 0$  there exist  $t_1, \dots, t_n \in J$  such that  $\|(1/n) \sum_{j=1}^n [T(t)T(t_j)y - M(T(\cdot)y)]\| < \varepsilon$  for all  $t \in J$ . Now  $z_\varepsilon = (1/n) \sum_{j=1}^n [z - T(t + t_j)z] \in R$  and  $\|z - z_\varepsilon\| < \varepsilon$ , so  $z \in \bar{R}$ . Hence  $y \in N + \bar{R}$  and the proposition is proved.

The following two results provide examples of ergodic vectors.

COROLLARY 4.4. *Let  $T : J \rightarrow L(X)$  be a representation and  $x \in X$ . If  $\text{orb}_T(x)$  is weakly relatively compact then  $x \in X_{\text{erg}}(T)$ .*

PROOF. Since  $\text{orb}_T(x)$  is weakly relatively compact, it is bounded and by Proposition 4.1,  $(T_F)$  is a system of invariant integrals for  $T(J)$  at  $x$ . Moreover,  $\text{co orb}_T(x)$  is weakly relatively compact so  $(T_F x)$  has a weak limit point  $y$ . By Corollary 4.2,  $x \in X_{\text{erg}}(T)$ .

PROPOSITION 4.5. *Let  $T : J \rightarrow L(X)$  be a bounded representation. If  $X$  is reflexive, or more generally if  $N + R$  is dense in  $X$ , then  $X_{\text{erg}} = X$ .*

PROOF. Since  $N + R \subseteq X_{\text{erg}} \subseteq X$  we conclude that  $X_{\text{erg}} = X$  whenever  $N + R$  is dense in  $X$ . It remains to prove that  $N + R$  is dense in  $X$  if  $X$  is reflexive. For  $S \subseteq X$  let  $S^\perp = \{\varphi \in X^* : \langle x, \varphi \rangle = 0 \text{ for all } x \in S\}$ . It is easy to check that  $R^\perp = N(T^*)$ . Hence for reflexive  $X$ ,  $R(T^*)^\perp = N(T^{**}) = N$ . Further,  $N^\perp = R(T^*)^{\perp\perp} = \bar{R}(T^*)$ . Hence  $(N + R)^\perp = N^\perp \cap R^\perp = \bar{R}(T^*) \cap N(T^*) = \{0\}$ , showing that  $N + R$  is dense in  $X$ .

As an application we present the following

PROPOSITION 4.6. *Given  $A \in L(X)$  define  $T : \mathbb{Z}^+ \rightarrow L(X)$  by  $T(n) = A^n$ . If  $x \in X_{\text{erg}}(T)$  and  $A^{n+1}x - A^n x \rightarrow 0$  as  $n \rightarrow \infty$  then  $A^n x \rightarrow y$  for some  $y \in X$  with  $Ay = y$ .*

PROOF. We apply Theorem 2.7 with  $\mathcal{A} = C_0(J, X)$ ,  $J = \mathbb{Z}^+$  and  $\varphi(n) = A^n x$ . Since  $\Delta_t \varphi \in \mathcal{A}$  for all  $t \in J$  and  $\varphi \in E(J, X)$  we conclude that  $\varphi - M_\varphi \in \mathcal{A}$ . So  $A^n x \rightarrow y$  where  $y = M_\varphi$ .

REMARK 4.7. If  $A \in L(X)$  and  $T : \mathbb{Z}^+ \rightarrow L(X)$  is given by  $T(n) = A^n$  then  $N(T) = \ker(A - I)$  and  $R(T) = \text{range}(A - I)$ . If  $A$  is power bounded then  $T$  is a bounded representation and if the Cesàro sums  $A_n x = (1/n) \sum_{j=1}^n A^j x$  converge weakly for some  $x \in X$  then  $T(\cdot)x$  is ergodic. If in addition  $X$  is reflexive then by Propositions 4.1 and 4.5,  $X = N \oplus \overline{R}$ . This special case may be found in [20, p.214]. Also see [10].

## 5. Cesàro and other means

Throughout this section we will assume that  $J$  is a measurable sub-semigroup of a locally compact abelian group  $G$  carrying a fixed Haar measure  $\mu$ . Let  $\mathcal{K}(G)$  denote the set of compact neighbourhoods of 0 in  $G$  and set  $\mathcal{K}(J) = \{V \cap J : V \in \mathcal{K}(G) \text{ and } \mu(V \cap J) \neq 0\}$ . We shall call a net  $(K_\alpha)_{\alpha \in \Lambda}$  in  $\mathcal{K}(J)$ , a *Følner net* if

$$(5.1) \quad \lim_{\alpha \in \Lambda} \frac{\mu(K_\alpha \Delta (K_\alpha + s))}{\mu(K_\alpha)} = 0 \quad \text{for all } s \in J,$$

where  $\Delta$  denotes symmetric difference.

Condition (5.1) was introduced by Følner (see [6, p.80]). As an example, let  $G = \mathbb{R}^2$  and  $J = \{(x_1, x_2) \in \mathbb{R}^2 : |x_2| \leq m(x_1 - a)\}$  where  $a \geq 0$  and  $m > 0$ . If  $K_r = \{x \in J : |x| \leq r\}$  then  $K_r \in \mathcal{K}(J)$ ,  $\mu(K) \sim r^2$  and  $\mu(K_r \Delta (K_r + s)) \sim r$  for fixed  $s \in J$ . Hence  $(K_r)_{r>a}$  is a Følner net.

We define the *Cesàro integrals* of functions  $\varphi \in C_b(J, X)$  by  $C_K \varphi(t) = (1/\mu(K)) \int_K \varphi(t+s) d\mu(s)$  for  $K \in \mathcal{K}(J)$ ,  $t \in J$ .

PROPOSITION 5.1. *If (5.1) holds then  $(C_{K_\alpha})_{\alpha \in \Lambda}$  is a system of invariant integrals for the translation semigroup  $\mathcal{R}$  acting on  $C_{ub}(J, X)$ .*

PROOF. Let  $K \in \mathcal{K}(J)$  and  $\varphi \in C_{ub}(J, X)$ . Given  $\varepsilon > 0$  choose  $V \in \mathcal{K}(G)$  such that  $\|\varphi_s - \varphi_t\|_\infty < \varepsilon$  for all  $t \in J$  and all  $s \in (t+V) \cap J$ . Since  $\|C_K \varphi(s) - C_K \varphi(t)\| \leq \|\varphi_s - \varphi_t\|_\infty$  we conclude that  $C_K \varphi \in C_{ub}(J, X)$ . Moreover,  $C_K \in L(C_{ub}(J, X))$ . Next, by the compactness of  $K$  we can choose  $t_1, \dots, t_m \in K$  such that  $K \subseteq \bigcup_{j=1}^m (t_j + V)$ . Set  $\pi_1 = (t_1 + V) \cap K$  and for  $2 \leq j \leq m$ ,  $\pi_j = (t_j + V) \cap K \setminus \bigcup_{i=1}^{j-1} \pi_i$ . Then  $K = \bigcup_{j=1}^m \pi_j$  and the  $\pi_j$  are disjoint measurable sets. Since

$$\left\| C_K \varphi - \sum_{j=1}^m \frac{\mu(\pi_j)}{\mu(K)} \varphi_{t_j} \right\| < \varepsilon$$

we conclude that  $C_K \varphi \in \overline{\text{co}}_{\mathcal{R}}(\varphi)$ , thereby proving (2.1).

For (2.3), let  $s \in J$ . Then

$$\begin{aligned}\|(C_K R_s - C_K)\varphi\|_\infty &= \sup_{t \in J} \left\| \frac{1}{\mu(K)} \int_K [\varphi(t+s+u) - \varphi(t+u)] d\mu(u) \right\| \\ &= \sup_{t \in J} \left\| \frac{1}{\mu(K)} \int_{K \Delta (K+s)} \varphi(t+u) d\mu(u) \right\| \\ &\leq \|\varphi\|_\infty \frac{\mu(K \Delta (K+s))}{\mu(K)}\end{aligned}$$

and (2.3) follows from (5.1). Since (2.2) is clear, the proposition is proved.

**COROLLARY 5.2.** *If  $\varphi \in C_{ub}(J, X)$  and (5.1) holds, then the following are equivalent*

- (i)  $\varphi \in E(J, X)$  and  $M(\varphi) = y$ ,
- (ii) the net  $(C_{K_\alpha}\varphi)_{\alpha \in \Lambda}$  converges to  $y$ ,
- (iii) some subnet of  $(C_{K_\alpha}\varphi)_{\alpha \in \Lambda}$  converges weakly to  $y$ .

**PROOF.** By Corollary 2.5 and Eberlein's mean ergodic theorem again, each of these conditions is equivalent to  $y \in \overline{\text{co}} \text{orb}_{\mathcal{A}}(\varphi)$  with  $y$  a constant function.

We come to our final system of invariant integrals. Let  $\mathcal{P} = \{f \in L^1(G) : f \geq 0 \text{ and } \hat{f}(0) = 1\}$ . Reiter [14, p.113] has proved the existence of a net  $(f_\alpha)_{\alpha \in \Lambda}$  in  $\mathcal{P}$  satisfying  $\lim_{\alpha \in \Lambda} \|R_s f_\alpha - f_\alpha\|_1 = 0$  for all  $s \in G$ . For  $\varphi \in C_{ub}(G, X)$  we can define  $A_\alpha \varphi \in C_{ub}(G, X)$  by  $A_\alpha \varphi = \varphi * f_\alpha$ . So  $\|A_\alpha \varphi\|_\infty \leq \|\varphi\|_\infty$  and  $A_\alpha \in L(C_{ub}(G, X))$ .

**PROPOSITION 5.3.** *The net  $(A_\alpha)_{\alpha \in \Lambda}$  is a system of invariant integrals for the translation semigroup  $\mathcal{R} = (R_s)_{s \in G}$  acting on  $C_{ub}(G, X)$ .*

**PROOF.** Given  $V \in \mathcal{X}(G)$  and  $\varphi \in C_{ub}(G, X)$  let  $f_V = (1/\mu(V))\chi_{-V}$  where  $\chi_{-V}$  is the characteristic function of  $-V$ . Then  $f_V \in \mathcal{P}$  and since  $\varphi * f_V = (1/\mu(V)) \int_V \varphi_s d\mu(s) = C_V \varphi$ , it follows from Proposition 5.1 that  $\varphi * f_V \in \overline{\text{co}} \text{orb}_{\mathcal{A}}(\varphi)$ . It is easy to check that  $\mathcal{P} \subseteq \overline{\text{co}}\{f_V : V \in \mathcal{X}(G)\}$ . Hence,  $\varphi * \mathcal{P} \subseteq \overline{\text{co}} \text{orb}_{\mathcal{A}}(\varphi)$ , proving (2.1). Since  $\|A_\alpha\| \leq 1$ , (2.2) holds. Finally, for  $s \in G$  we have

$$\|(A_\alpha R_s - A_\alpha)\varphi\|_\infty = \|(R_s \varphi - \varphi) * f_\alpha\|_\infty = \|\varphi * (R_s f_\alpha - f_\alpha)\|_\infty \leq \|\varphi\|_\infty \|R_s f_\alpha - f_\alpha\|_1.$$

From the definition of  $(f_\alpha)$ , (2.3) follows and the proposition is proved.

As for Corollary 5.2 we obtain

**COROLLARY 5.4.** *For  $\varphi \in C_{ub}(G, X)$  the following are equivalent*

- (i)  $\varphi \in E(G, X)$  and  $M(\varphi) = y$ ,
- (ii) the net  $(A_\alpha \varphi)_{\alpha \in \Lambda}$  converges to  $y$ ,
- (iii) some subnet of  $(A_\alpha \varphi)_{\alpha \in \Lambda}$  converges weakly to  $y$ .

Argabright [2] used the Reiter nets  $(f_\alpha)$  to prove an ergodic limit for scalar-valued Eberlein weakly almost periodic functions on  $G$ . Datry and Muraz [7] also used them to introduce ergodicity in Banach  $L^1(G)$ -modules.

We conclude with two more examples, firstly of some ergodic functions and secondly of a non-ergodic one. Recall that for a function  $\varphi \in C_b(G, X)$  the set  $I(\varphi) = \{f \in L^1(G) : \varphi * f = 0\}$  is a closed ideal of  $L^1(G)$ . Let  $\widehat{G}$  denote the character group of  $G$ ,  $0$  the unit of  $\widehat{G}$ , and  $\hat{f} : G \rightarrow \mathbb{C}$  the Fourier transform of  $f$ . The *Beurling spectrum* of  $\varphi$  is  $\text{sp}(\varphi) = \{\gamma \in \widehat{G} : \hat{f}(\gamma) = 0 \text{ for all } f \in I(\varphi)\}$ .

**THEOREM 5.5.** *If  $\varphi \in C_{ub}(G, X)$  and  $0 \notin \text{sp}(\varphi)$  then  $\varphi \in E(G, X)$ .*

**PROOF.** Take  $V \in \mathcal{K}(\widehat{G})$  with  $V \cap \text{sp}(\varphi) = \emptyset$  and  $f \in L^1(G)$  with  $\hat{f}(0) = 1$  and  $\text{supp}(\hat{f}) \subseteq V$ . Then  $\text{sp}(\varphi * f) = \emptyset$  so  $\varphi * f = 0$ . Moreover,  $f$  is continuous. Now, given  $\varepsilon > 0$ , choose a compact set  $K$  in  $G$  such that  $\int_{G \setminus K} |f(t)| d\mu(t) < \varepsilon/(1 + 2\|\varphi\|_\infty)$ . For  $s \in G$  define  $g(s) = (\varphi - \varphi_{-s})f(s)$ . Hence  $\int_G g(s) d\mu(s) = \varphi - \varphi * f = \varphi$ . Moreover, by Proposition 2.2,  $g(s) \in E(G, X)$  and since  $\varphi$  is uniformly continuous,  $g : G \rightarrow E(G, X)$  is continuous. Since  $K$  is compact,  $g|_K$  is separably-valued and hence Bochner integrable. Therefore  $\int_K g(s) d\mu(s) \in E(G, X)$ . But  $\|\varphi - \int_K g(s) d\mu(s)\| \leq \|\int_{G \setminus K} g(s) d\mu(s)\| < \varepsilon$  and so  $\varphi \in E(G, X)$  as claimed.

**EXAMPLE 5.6.** Define  $\varphi : \mathbb{R} \rightarrow c_0$  by  $\varphi(t) = (\sin(t/n))_{n=1}^\infty$ . One easily checks that  $\varphi \in C_{ub}(\mathbb{R}, c_0)$ . Now the range of  $\varphi$  is not relatively compact in  $c_0$ . For, if it were, then given  $0 < \varepsilon < 1/4$  there would exist  $t_1, \dots, t_m \in \mathbb{R}$  such that  $\inf_j \|\varphi(t) - \varphi(t_j)\| < \varepsilon$  for all  $t \in \mathbb{R}$ . In particular we would have  $|\sin(t/n)| < 2\varepsilon$  for all  $n > N(\varepsilon)$  and all  $t \in \mathbb{R}$ , which is false. It follows that  $\varphi$  is not almost periodic. On the other hand  $\varphi'$  is almost periodic (see [1, p. 53]) and so  $\varphi \notin E(\mathbb{R}, c_0)$ . For otherwise, by Levitan [13] or Basit [3, Theorem 3.1.1] it would follow that  $\varphi$  is almost periodic. From Theorem 5.5 we conclude that  $0 \in \text{sp}(\varphi)$ .

## References

- [1] L. Amerio and G. Prouse, *Almost periodic functions and functional equations* (Van Nostrand, New York, 1971).
- [2] L. N. Argabright, 'On the mean value of weakly almost periodic functions', *Proc. Amer. Math. Soc.* **36** (1972), 315–316.
- [3] B. Basit, 'Some problems concerning different types of vector-valued almost periodic type functions', *Dissertationes Math. (Rozprawy Mat.)* **338** (1995).

- [4] ———, 'Harmonic analysis and asymptotic behavior of solutions to the abstract Cauchy problem', *Semigroup Forum* **54** (1997), 58–74.
- [5] B. Basit and A. J. Pryde, 'Polynomials and functions with finite spectra on locally compact abelian groups', *Bull. Austral. Math. Soc.* **51** (1994), 33–42.
- [6] J. F. Berglund, H. D. Junghenn and P. Milnes, *Analysis on Semigroups: Function spaces, compactifications, representations* (Wiley-Interscience, New York, 1989).
- [7] C. Datry and G. Muraz, 'Analyse harmonique dans les modules de Banach II: presque-périodicité et ergodicité', *Bull. Science Math. (2)* **120** (1996), 493–536.
- [8] N. Dunford and J. T. Schwartz, *Linear operators*, Parts I, II (Interscience, New York, 1958, 1963).
- [9] W. F. Eberlein, 'Abstract ergodic theorems and weak almost periodic functions', *Trans. Amer. Math. Soc.* **69** (1949), 217–240.
- [10] J. A. Goldstein, 'Application of operator semigroups to Fourier analysis', *Semigroup Forum* **52** (1996), 37–47.
- [11] K. Iseki, 'Vector valued functions on semigroups, I-III', *Proc. Japan Acad. Ser. A Math. Sci.* (1955), 16–19, 152–155 and 699–702.
- [12] Y. Katznelson and L. Tzafriri, 'On power bounded operators', *J. Funct. Anal.* **68** (1986), 313–328.
- [13] B. M. Levitan, 'Integration of almost periodic functions with values in Banach spaces', *Math. USSR-Izv.* **30** (1966), 1101–1110 (in Russian).
- [14] H. Reiter, *Classical Fourier analysis on locally compact groups* (Oxford University Press, 1968).
- [15] W. Rudin, *Harmonic analysis on groups* (Interscience, New York, 1963).
- [16] W. M. Ruess and V. Q. Phóng, 'Asymptotically almost periodic solutions of evolution equations in Banach spaces', *J. Differential Equations* **122** (1995), 282–301.
- [17] W. M. Ruess and W. H. Summers, 'Weak almost periodicity and the strong ergodic limit theorem for contraction semigroups', *Israel J. Math.* **64** (1988), 139–157.
- [18] ———, 'Integration of asymptotically almost periodic functions and weak almost periodicity', *Dissertationes Math. (Rozprawy Mat.)* **279** (1989).
- [19] ———, 'Ergodicity theorems for semigroups of operators', *Proc. Amer. Math. Soc.* **114** (1992), 423–432.
- [20] K. Yosida, *Functional Analysis* (Springer, Berlin, 1966).
- [21] C. Zhang, 'Vector-valued means and their applications in some vector-valued function spaces', *Dissertationes Math. (Rozprawy Mat.)* **334** (1994).

Department of Mathematics

Monash University

Clayton, VIC 3168

Australia

e-mail: [bbasit\(ajpryde\)@vaxc.cc.monash.edu.au](mailto:bbasit(ajpryde)@vaxc.cc.monash.edu.au)