

Molecular Machinery Mediating Vesicle Budding, Docking and Fusion

Thomas H. Söllner* and James E. Rothman

Cellular Biochemistry and Biophysics Program, Memorial Sloan-Kettering Cancer Center, 1275 York Avenue, New York, NY 10021, USA

Key words: vesicle budding and fusion/ARF/coat proteins/cargo receptor/NSF/SNAP/SNARE

ABSTRACT. A general machinery buds and fuses transport vesicles which connect intracellular compartments with each other and allow communication with the extracellular environment. Cytoplasmic coat proteins deform membranes to bud vesicles and interact directly or indirectly with cargo molecules. Compartment-specific SNAREs on vesicles and target membranes dock vesicles and provide a scaffolding for the general fusion machinery to initiate lipid bilayer fusion.

Each compartment in eukaryotic cells is characterized by a set of specific proteins and a distinct lipid composition, which together define its unique functional properties. The compartments along the major transport routes termed secretion, endocytosis and transcytosis exchange components with others by means of transport vesicles (43). Each route is composed of a series of sequential vesicular transport steps. At each step transport vesicles form at a donor compartment and fuse with a specific acceptor compartment. To maintain the identity of the individual organelles, cargo has to be selected during vesicle budding and has to be specifically delivered to its correct destination. Recent data have provided insight into the underlying mechanisms of these processes, thereby expanding our understanding of how cells maintain and control their temporal and spatial organization and how they communicate with their environment. The core machinery mediating these processes will be outlined in the following sections.

Vesicle budding

Shuttling of cargo between organelles requires its packaging into distinct transport containers defined by their distinct size and, to a certain degree, specialized protein and lipid composition. In every instance, the generation of defined transport units is a prerequisite to control and balance import in and export from a distinct organelle. To ensure the generation of such transport units, coat proteins derived from the cytoplasm assemble at budding sites on the donor membrane. This

process is triggered by nucleotide exchange on a class of small soluble GTP binding proteins, their membrane recruitment and the subsequent binding of coat proteins. The assembly of coat proteins deforms the membrane resulting in the generation of coated buds which finally pinch off to release coated vesicles. Following their uncoating the vesicles dock to and fuse with the target membrane.

In case of the Golgi, the vesicular protein coat is termed COPI (coat protein) and assembles from the cytoplasmic coatomer complexes (coat protomer) consisting of seven subunits (53, 64). The binding of coatomer to the donor membrane and the subsequent assembly process occur after ADP-ribosylation factor (ARF) in its GTP-bound form interacts with the membrane (52). A GDP/GTP exchange factor and an ARF receptor mediating this processes have been postulated, but still await purification. Recent data also suggest the involvement of polyphosphoinositides and phosphatidic acid in the budding step (10, 13, 18). These lipids seem to facilitate nucleotide exchange on ARF and might enhance the interaction of coatomer with the membrane. The vesicle fission step has been shown to require an additional lipophilic component, fatty acyl — coenzyme A (41). Neither the target nor the exact function of the fatty acyl requirement are known. After the vesicle has pinched off, ARF hydrolyzes its bound GTP (60), the coat disassembles and dissociates from the vesicle membrane, an uncoated vesicle is generated which goes on to dock and fuse with the target membrane. GTP hydrolysis seems to be regulated by ARF-GAP (GTPase activating protein) a cytosolic protein which is recruited to Golgi membranes and activated by a still unknown mechanism (16). Again, polyphosphoinositides have been implicated to function at this step (35).

Vesicle formation may require proper cargo selec-

* To whom correspondence should be addressed.
Telephone: 212-639-8598, FAX: 212-717-3604

Abbreviations. used in this paper: NSF, N-ethylmaleimide-sensitive fusion protein; SNAP, soluble NSF attachment protein; SNARE, SNAP receptor; ER, endoplasmic reticulum; ARF, ADP-ribosylation factor; NEM, N-ethylmaleimide; GAP, GTPase activating protein.

tion, or, in some cases, cargo could be transported by default at its bulk concentration in a process termed bulk flow (67). Cargo selection is accomplished by a protein's (or lipid's) enrichment in or exclusion from transport vesicles which requires that cargo molecules carry transport signals or that resident components contain retention signals (45). For example, the amino acid motifs KKXX and XXRR at the carboxy- and amino-termini, respectively, of integral membrane proteins act as transport signals for retrieval of resident proteins which have escaped from the endoplasmic reticulum (ER) (29, 38, 51). KKXX or related peptides have been shown to interact with COPI coats, illustrating how a signal contained in a cargo molecule can be coupled to the transport machinery and thus ensure cargo packaging (14, 32). However, since COPI-coated vesicles also carry cargo in the anterograde direction, coatomer should not interact with the KKXX motif at the ER membrane. Possible mechanisms could be that COPI coats exist in different conformational states, binding cargo either for anterograde or retrograde transport, or that posttranslational modifications or interactions with additional components alter the binding properties of coatomer for KKXX (48).

Luminal proteins, which cannot possibly interact with cytosolic coat proteins in a direct way, likely contain other types of transport signals, which might be recognized by cargo receptors, integral membrane proteins which function as adapters to mediate the interaction with the vesicle coats. A family of integral membrane proteins, major constituents of both COPI and COPII coated vesicles, has been recently identified, whose members fulfill the criteria for such cargo receptors. P24 proteins have a variable amino-terminal domain, which might interact with cargo, and a conserved carboxy-terminal domain, binding coat proteins (57) (unpublished observation). Deletion of individual family members does not affect cell viability but does alter the transport efficiency of some but not all cargo molecules (50).

Vesicle docking and fusion

Transport from the donor compartment to the acceptor compartment is in some instances facilitated by vesicle-associated motors interacting with the cytoskeleton (62, 63), but this is not a prerequisite for transport and hence cannot be critical for targeting transport vesicles selectively to their destination. Thus, signals on the vesicles must exist that are recognized by a receptor on the intended target membrane. According to the SNARE hypothesis, this process is mediated — at least in part — by the unique pairing of SNAP receptors localized on vesicles (*v*-SNAREs) with their cognate *t*-SNAREs on target membranes (56). Most SNAREs are type II membrane proteins, with short luminal or extra-cellular re-

gions and comparably large cytosolic domains. The cytoplasmic domains are predicted to form coiled coils and to mediate the unique interactions between the distinct members of the *v/t*-SNARE families (11). The system is best characterized in the yeast *Saccharomyces cerevisiae*. *T*-SNAREs (which should provide a functional definition of compartment borders) have been localized to the endoplasmic reticulum (*Ufe1*) (34), the *cis* Golgi network (*Sed5*) (3), the vacuole (*Pep12*) (30), and the plasma membrane (*Sso1,2*) (1). Physiological evidence for the role of SNARE proteins is provided by the fact that inactivation of *t*-SNAREs causes the accumulation of transport vesicles, which are no longer able to dock to the correct compartment. Massive overexpression of *t*-SNAREs has been shown to perturb cellular morphology, presumably by altering the *t*-SNARE containing compartment (25). A *t*-SNAREs localized to a compartment in which vesicular transport routes from several directions converge might serve several purposes in providing a docking site for vesicles originating from different organelles. For example, coimmunoprecipitation experiments with an antibody directed against *Sed5*, a *t*-SNARE localized to the early Golgi, revealed the presence of several *Sed5* interacting *v*-SNAREs, some of them mediating the docking of ER-derived vesicles, others presumably allowing the binding of vesicles derived from downstream Golgi compartments and returning to the early Golgi (2, 54). *V*-SNAREs implicated in Golgi to plasma membrane transport were, however, absent in the SNARE complexes containing *Sed5* (54). They form a distinct complex with the plasma membrane *t*-SNAREs, thereby illustrating the high specificity of *v*-SNARE-*t*-SNARE interactions (15, 47). Physiological evidence for the role of *v*-SNAREs in yeast is provided by the fact that some of them were originally identified in secretion deficient mutants and that inactivation of these proteins results in accumulation of transport vesicles *in vivo* (17, 37, 39).

While, studies in yeast can establish the basic mechanisms of vesicular targeting and fusion, studies in higher eukaryotes reveal an additional level of complexity. This is due to the complex organization of multicellular organisms with specialized cell types, organized in different tissues, each fulfilling specific tasks, and the establishment of elaborate intercellular communication systems. For example, in the rat four different *t*-SNAREs localized to the plasma membrane (syntaxins 1–4) have been identified, some of them showing tissue specificity (7). Another example is the neuronal synapse, which is specialized in neurotransmitter release and used as a model system to study regulated secretion. Regulated secretion involves additional proteins that temporarily lock the fusion machinery in place until an appropriate signal releases the constraint (19, 22, 23, 58). Recent data indicate that the synaptic vesicle protein, synapto-




Fig. 1. The molecular machinery mediating vesicle budding and fusion. A) Transport vesicle formation. i) Vesicle budding is initiated when GDP on ARF is exchanged against GTP. The resulting ARF [GTP] binds to the membrane and recruits coat proteins. The assembly of coat proteins on the membrane incrementally deforms the corresponding area of the lipid bilayer, resulting in a bud. ii) The bud pinches off in a reaction requiring fatty acyl coenzyme A. iii) GTP hydrolyses results in dissociating of ARF yielding presumably a still coated vesicle (vi), whose coat then disassembles and dissociates from the membrane (v). B) Targeting of vesicles. Uncoated vesicles interact via their address signal, termed v-SNARE with their cognate receptor on the target membrane, termed t-SNARE. C) Initiation of vesicle fusion. SNAPs bind to assembled SNARE complexes at the attachment site of the vesicles to the target membrane thereby allowing the subsequent binding of NSF. ATP hydrolysis by NSF disrupts the SNARE complex and initiates membrane fusion. It is not know whether additional components are required for the lipid bilayer fusion.

tagmin plays a dual role as a specialized v-SNARE (49) on the one hand and as a calcium sensor, on the other, and thereby links the signal for neurotransmitter release to the vesicle fusion machinery. The presence of two v-SNAREs on synaptic vesicles (VAMP and synaptotagmin) and a cognate t-SNARE pair on the presynaptic plasma membrane (syntaxin and SNAP-25) provides a multivalent vesicle binding site, explaining why neither the proteolytic cleavage of VAMP by tetanus toxin and certain serotypes of botulinum toxins nor the deletion of syntaxin prevents the accumulation of synaptic vesicles in the active zone (4, 6, 9, 28, 42, 59, 61).

Additional proteins regulating SNARE activity have been identified, and are likely to ensure that SNAREs are only active in their proper intracellular localization. Members of the Sec 1 protein family have been shown to interact with t-SNAREs and control their exposure to the cognate v-SNARE (26, 46). Rab proteins, small GTP-binding proteins are required for the assembly of SNARE complexes (54). Both rab proteins and Sec 1 family members are cytoplasmic proteins which are recruited to the membrane and exhibit a certain organelle specificity, indicating that these proteins likely add an additional layer of specificity to vesicular transport, maybe by a proofreading mechanism. More specialized interactions of distinct SNAREs with other components have been demonstrated. For example VAMP has been shown to interact with the synaptic vesicle protein synaptophysin in a way that is mutually exclusive with the binding to its t-SNAREs (21). The t-SNARE syntaxin 1 has been found to be associated with N-type calcium channels — yet another instance of close coupling between the vesicle docking and fusion machinery and that carrying the signal for exocytosis (6, 33, 69).

Assembled SNARE complexes provide a scaffolding for binding of general fusion components such as the N-ethylmaleimide-sensitive fusion protein (NSF) and the soluble NSF attachment proteins (SNAPs) (8, 12, 36, 66). The general fusion proteins are cytoplasmic proteins which function at many sites (5, 20, 24). Inactivation of NSF by NEM causes the accumulation of docked uncoated vesicles (40), indicating that NSF is required for consumption of vesicles docked via SNAREs. Additional evidence for the physiological role of NSF and for its general function in a variety of vesicular transport processes was provided by NSF mutants. The NSF mutant in *Drosophila* comatose causes paralysis of the flies, clearly establishing NSF's function in neurotransmission (44). Similarly, the NSF mutant in yeast, sec 18, causes an accumulation of transport vesicles and assembled SNARE complexes (31, 54, 68). SNAPs mediate the interaction of NSF with membranes, as has been demonstrated using Golgi membranes (12, 65). Three to six SNAP proteins bind to an assembled v-SNARE-t-SNARE complex (27, 55). In the

neuronal synapse, the brain specific form of SNAP (β -SNAP), but not the ubiquitous SNAP (α -SNAP) interacts with the specialized v-SNARE synaptotagmin (49). ATP hydrolysis by NSF provides the energy to disrupt the complex, likely by causing conformational changes in SNAPs and one or several of the SNARE proteins (55). The actual lipid bilayer fusion process is not understood; it cannot be excluded that additional components are necessary, but it seems likely that after ATP hydrolysis by NSF one or several of the above mentioned components are in an activated state to initiate fusion.

In summary, the rational scheme underlying transport between intra-cellular compartments seems to be established. But although the identification and isolation of the core machinery mediating these processes has proceeded during the past years, we are far from having a complete understanding of mechanistic details. On the one hand, studying the machinery in its cellular background will help us to confirm its physiological functions, and on the other reconstitution of the purified components in vitro will provide the biophysical details about the reaction mechanisms. The completion of the yeast genome project will greatly facilitate the isolation of additional components and should give us pointers as to the minimal set of proteins required for vesicular transport.

Acknowledgments. We thank Mark Craighead and Gero Miesenböck for critically reading the manuscript and for their helpful discussions.

REFERENCES

1. AALTO, M.K., RONNE, H., and KERANEN, S. 1993. Yeast syntaxins Sso1p and Sso2p belong to a family of related membrane proteins that function in vesicular transport. *Embo. J.*, **12**: 4095–4104.
2. BANFIELD, D.K., LEWIS, M.J., and PELHAM, H.R. 1995. A SNARE-like protein required for traffic through the Golgi complex. *Nature*, **375**: 806–809.
3. BANFIELD, D.K., LEWIS, M.J., RABOUILLE, C., WARREN, G., and PELHAM, H.R. 1994. Localization of Sed5, a putative vesicle targeting molecule, to the cis-Golgi network involves both its transmembrane and cytoplasmic domains. *J. Cell Biol.*, **127**: 357–371.
4. BAUMERT, M., MAYCOX, P.R., NAVONE, F., DE CAMILLI, P., and JAHN, R. 1989. Synaptobrevin: an integral membrane protein of 18,000 daltons present in small synaptic vesicles of rat brain. *Embo. J.*, **8**: 379–384.
5. BECKERS, C.J., BLOCK, M.R., GLICK, B.S., ROTHMAN, J.E., and BALCH, W.E. 1989. Vesicular transport between the endoplasmic reticulum and the Golgi stack requires the NEM-sensitive fusion protein. *Nature*, **339**: 397–398.
6. BENNETT, M.K., CALAKOS, N., and SCHELLER, R.H. 1992. Syntaxin: a synaptic protein implicated in docking of synaptic vesicles at presynaptic active zones. *Science*, **257**: 255–259.
7. BENNETT, M.K., GARCIA-ARRARAS, J.E., ELFERINK, L.A.,

- PETERSON, K., FLEMING, A.M., HAZUKA, C.D., and SCHELLER, R.H. 1993. The syntaxin family of vesicular transport receptors. *Cell*, **74**: 863–873.
8. BLOCK, M.R., GLICK, B.S., WILCOX, C.A., WIELAND, F.T., and ROTHMAN, J.E. 1988. Purification of an N-ethylmaleimide-sensitive protein catalyzing vesicular transport. *Proc. Natl. Acad. Sci. USA*, **85**: 7852–7856.
 9. BROADIE, K., PROKOP, A., BELLEN, H.J., O'KANE, C.J., SCHULZE, K.L., and SWEENEY, S.T. 1995. Syntaxin and synaptobrevin function downstream of vesicle docking in *Drosophila*. *Neuron*, **15**: 663–673.
 10. BROWN, H.A., GUTOWSKI, S., MOOMAW, C.R., SLAUGHTER, C., and STERNWEIS, P.C. 1993. ADP-ribosylation factor, a small GTP-dependent regulatory protein, stimulates phospholipase D activity [see comments]. *Cell*, **75**: 1137–1144.
 11. CHAPMAN, E.R., AN, S., BARTON, N., and JAHN, R. 1994. SNAP-25, a t-SNARE which binds to both syntaxin and synaptobrevin via domains that may form coiled coils. *J. Biol. Chem.*, **269**: 27427–27432.
 12. CLARY, D.O., GRIFF, I.C., and ROTHMAN, J.E. 1990. SNAPS, a family of NSF attachment proteins involved in intracellular membrane fusion in animals and yeast. *Cell*, **61**: 709–721.
 13. COCKCROFT, S., THOMAS, G.M., FENSOME, A., GENY, B., CUNNINGHAM, E., GOUT, I., HILES, I., TOTTY, N.F., TRUONG, O., and HSUAN, J.I. 1994. Phospholipase D: a downstream effector of ARF in granulocytes. *Science*, **263**: 523–526.
 14. COSSON, P. and LETOURNEUR, F. 1994. Coatamer interaction with di-lysine endoplasmic reticulum retention motifs. *Science*, **263**: 1629–1631.
 15. COUVE, A. and GERST, J.E. 1994. Yeast Snc proteins complex with Sec9. Functional interactions between putative SNARE proteins. *J. Biol. Chem.*, **269**: 23391–23394.
 16. CUKIERMAN, E., HUBER, I., ROTMAN, M., and CASSEL, D. 1995. The ARF1 GTPase-activating protein: zinc finger motif and Golgi complex localization. *Science*, **270**: 1999–2002.
 17. DASCHER, C., OSSIG, R., GALLWITZ, D., and SCHMITT, H.D. 1991. Identification and structure of four yeast genes (SLY) that are able to suppress the functional loss of YPT1, a member of the RAS superfamily. *Mol. Cell Biol.*, **11**: 872–885.
 18. DE CAMILLI, P., EMR, S.D., MCPHERSON, P.S., and NOVICK, P. 1996. Phosphoinositides as regulators in membrane traffic. *Science*, **271**: 1533–1539.
 19. DEBELLO, W.M., BETZ, H., and AUGUSTINE, G.J. 1993. Synaptotagmin and neurotransmitter release [comment]. *Cell*, **74**: 947–950.
 20. DIAZ, R., MAYORGA, L.S., WEIDMAN, P.J., ROTHMAN, J.E., and STAHL, P.D. 1989. Vesicle fusion following receptor-mediated endocytosis requires a protein active in Golgi transport. *Nature*, **339**: 398–400.
 21. EDELMANN, L., HANSON, P.I., CHAPMAN, E.R., and JAHN, R. 1995. Synaptobrevin binding to synaptophysin: a potential mechanism for controlling the exocytotic fusion machine. *Embo. J.*, **14**: 224–231.
 22. ELFERINK, L.A., PETERSON, M.R., and SCHELLER, R.H. 1993. A role for synaptotagmin (p65) in regulated exocytosis. *Cell*, **72**: 153–159.
 23. GEPPERT, M., GODA, Y., HAMMER, R.E., LI, C., ROSAHL, T.W., STEVENS, C.F., and SÜDHOF, T.C. 1994. Synaptotagmin I: a major Ca²⁺ sensor for transmitter release at a central synapse. *Cell*, **79**: 717–727.
 24. GRAHAM, T.R. and EMR, S.D. 1991. Compartmental organization of Golgi-specific protein modification and vacuolar protein sorting events defined in a yeast sec 18 (NSF) mutant. *J. Cell Biol.*, **114**: 207–218.
 25. HARDWICK, K.G. and PELHAM, H.R. 1992. SED5 encodes a 39-kD integral membrane protein required for vesicular transport between the ER and the Golgi complex. *J. Cell Biol.*, **119**: 513–521.
 26. HARRISON, S.D., BROADIE, K., VAN DE GOOR, J., and RUBIN, G.M. 1994. Mutations in the *Drosophila* Rop gene suggest a function in general secretion and synaptic transmission. *Neuron*, **13**: 555–566.
 27. HAYASHI, T., MCMAHON, H., YAMASAKI, S., BINZ, T., HATA, Y., SÜDHOF, T.C., and NIEMANN, H. 1994. Synaptic vesicle membrane fusion complex: action of clostridial neurotoxins on assembly. *Embo. J.*, **13**: 5051–5061.
 28. HUNT, J.M., BOMMERT, K., CHARLTON, M.P., KISTNER, A., HABERMANN, E., AUGUSTINE, G.J., and BETZ, H. 1994. A post-docking role for synaptobrevin in synaptic vesicle fusion. *Neuron*, **12**: 1269–1279.
 29. JACKSON, M.R., NILSSON, T., and PETERSON, P.A. 1990. Identification of a consensus motif for retention of transmembrane proteins in the endoplasmic reticulum. *Embo. J.*, **9**: 3153–3162.
 30. JONES, E.W. 1977. Proteinase mutants of *Saccharomyces cerevisiae*. *Genetics*, **85**: 23–33.
 31. KAISER, C.A. and SCHEKMAN, R. 1990. Distinct sets of SEC genes govern transport vesicle formation and fusion early in the secretory pathway. *Cell*, **61**: 723–733.
 32. LETOURNEUR, F., GAYNOR, E.C., HENNECKE, S., DEMOLLIERE, C., DUDEN, R., EMR, S.D., RIEZMAN, H., and COSSON, P. 1994. Coatamer is essential for retrieval of dilysine-tagged proteins to the endoplasmic reticulum. *Cell*, **79**: 1199–1207.
 33. LEVEQUE, C., HOSHINO, T., DAVID, P., SHOJI-KASAI, Y., LEYS, K., OMORI, A., LANG, B., EL FAR, O., SATO, K., MARTIN-MOUTOT, N., and *et al.* 1992. The synaptic vesicle protein synaptotagmin associates with calcium channels and is a putative Lambert-Eaton myasthenic syndrome antigen. *Proc. Natl. Acad. Sci. USA*, **89**: 3625–3629.
 34. LEWIS, M.J. and PELHAM, H.R. 1996. SNARE-mediated retrograde traffic from the Golgi complex to the endoplasmic reticulum. *Cell*, **85**: 205–215.
 35. MAKLER, V., CUKIERMAN, E., ROTMAN, M., ADMON, A., and CASSEL, D. 1995. ADP-ribosylation factor-directed GTPase-activating protein. Purification and partial characterization. *J. Biol. Chem.*, **270**: 5232–5237.
 36. MALHOTRA, V., ORCI, L., GLICK, B.S., BLOCK, M.R., and ROTHMAN, J.E. 1988. Role of an N-ethylmaleimide-sensitive transport component in promoting fusion of transport vesicles with cisternae of the Golgi stack. *Cell*, **54**: 221–227.
 37. NEWMAN, A.P., SHIM, J., and FERRO-NOVICK, S. 1990. BET1, BOS1, and SEC22 are members of a group of interacting yeast genes required for transport from the endoplasmic reticulum to the Golgi complex. *Mol. Cell Biol.*, **10**: 3405–3414.
 38. NILSSON, T., JACKSON, M., and PETERSON, P.A. 1989. Short cytoplasmic sequences serve as retention signals for transmembrane proteins in the endoplasmic reticulum. *Cell*, **58**: 707–718.
 39. NOVICK, P., FIELD, C., and SCHEKMAN, R. 1980. Identification of 23 complementation groups required for post-translational events in the yeast secretory pathway. *Cell*, **21**: 205–215.
 40. ORCI, L., MALHOTRA, V., AMHERDT, M., SERAFINI, T., and ROTHMAN, J.E. 1989. Dissection of a single round of vesicular transport: sequential intermediates for intercisternal movement in the Golgi stack. *Cell*, **56**: 357–368.
 41. OSTERMANN, J., ORCI, L., TANI, K., AMHERDT, M., RAVAZZOLA, M., ELAZAR, Z., and ROTHMAN, J.E. 1993. Step-wise assembly of functionally active transport vesicles. *Cell*, **75**:

- 1015–1025.
42. OYLER, G.A., HIGGINS, G.A., HART, R.A., BATTENBERG, E., BILLINGSLEY, M., BLOOM, F.E., and WILSON, M.C. 1989. The identification of a novel synaptosomal-associated protein, SNAP-25, differentially expressed by neuronal subpopulations. *J. Cell Biol.*, **109**: 3039–3052.
 43. PALADE, G. 1975. Intracellular aspects of the process of protein synthesis. *Science*, **189**: 347–358.
 44. PALLANCK, L., ORDWAY, R.W., and GANETZKY, B. 1995. A *Drosophila* NSF mutant [letter]. *Nature*, **376**: 25.
 45. PEARSE, B.M. 1988. Receptors compete for adaptors found in plasma membrane coated pits. *Embo. J.*, **7**: 3331–3336.
 46. PEVSNER, J., HSU, S.C., BRAUN, J.E., CALAKOS, N., TING, A.E., BENNETT, M.K., and SCHELLER, R.H. 1994. Specificity and regulation of a synaptic vesicle docking complex. *Neuron*, **13**: 353–361.
 47. PROTOPOPOV, V., GOVINDAN, B., NOVICK, P., and GERST, J.E. 1993. Homologs of the synaptobrevin/VAMP family of synaptic vesicle proteins function on the late secretory pathway in *S. cerevisiae*. *Cell*, **74**: 855–861.
 48. ROTHMAN, J.E. and WIELAND, F.T. 1996. Protein sorting by transport vesicles. *Science*, **272**: 227–234.
 49. SCHIAVO, G., GMACHL, M.J., STENBECK, G., SÖLLNER, T.H., and ROTHMAN, J.E. 1995. A possible docking and fusion particle for synaptic transmission. *Nature*, **378**: 733–736.
 50. SCHIMMOLLER, F., SINGER-KRUGER, B., SCHRODER, S., KRUGER, U., BARLOWE, C., and RIEZMAN, H. 1995. The absence of Emp24p, a component of ER-derived COPII-coated vesicles, causes a defect in transport of selected proteins to the Golgi. *Embo. J.*, **14**: 1329–1339.
 51. SCHUTZE, M.P., PETERSON, P.A., and JACKSON, M.R. 1994. An N-terminal double-arginine motif maintains type II membrane proteins in the endoplasmic reticulum. *Embo. J.*, **13**: 1696–1705.
 52. SERAFINI, T., ORCI, L., AMHERDT, M., BRUNNER, M., KAHN, R.A., and ROTHMAN, J.E. 1991. ADP-ribosylation factor is a subunit of the coat of Golgi-derived COP-coated vesicles: a novel role for a GTP-binding protein. *Cell*, **67**: 239–253.
 53. SERAFINI, T., STENBECK, G., BRECHT, A., LOTTSPEICH, F., ORCI, L., ROTHMAN, J.E., and WIELAND, F.T. 1991. A coat subunit of Golgi-derived non-clathrin-coated vesicles with homology to the clathrin-coated vesicle coat protein beta-adaptin. *Nature*, **349**: 215–220.
 54. SOGAARD, M., TANI, K., YE, R.R., GEROMANOS, S., TEMPST, P., KIRCHHAUSEN, T., ROTHMAN, J.E., and SÖLLNER, T. 1994. A rab protein is required for the assembly of SNARE complexes in the docking of transport vesicles. *Cell*, **78**: 937–948.
 55. SÖLLNER, T., BENNETT, M.K., WHITEHEART, S.W., SCHELLER, R.H., and ROTHMAN, J.E. 1993. A protein assembly-disassembly pathway *in vitro* that may correspond to sequential steps of synaptic vesicle docking, activation, and fusion. *Cell*, **75**: 409–418.
 56. SÖLLNER, T., WHITEHEART, S.W., BRUNNER, M., ERDJUMENT-BROMAGE, H., GEROMANOS, S., TEMPST, P., and ROTHMAN, J.E. 1993. SNAP receptors implicated in vesicle targeting and fusion [see comments]. *Nature*, **362**: 318–324.
 57. STAMNES, M.A., CRAIGHEAD, M.W., HOE, M.H., LAMPEN, N., GEROMANOS, S., TEMPST, P., and ROTHMAN, J.E. 1995. An integral membrane component of coatamer-coated transport vesicles defines a family of proteins involved in budding. *Proc. Natl. Acad. Sci. USA*, **92**: 8011–8015.
 58. SÜDHOF, T.C. 1995. The synaptic vesicle cycle: a cascade of protein-protein interactions. *Nature*, **375**: 645–653.
 59. SWEENEY, S.T., BROADIE, K., KEANE, J., NIEMANN, H., and O'KANE, C.J. 1995. Targeted expression of tetanus toxin light chain in *Drosophila* specifically eliminates synaptic transmission and causes behavioral defects. *Neuron*, **14**: 341–351.
 60. TANIGAWA, G., ORCI, L., AMHERDT, M., RAVAZZOLA, M., HELMS, J.B., and ROTHMAN, J.E. 1993. Hydrolysis of bound GTP by ARF protein triggers uncoating of Golgi-derived COP-coated vesicles. *J. Cell Biol.*, **123**: 1365–1371.
 61. TRIMBLE, W.S., COWAN, D.M., and SCHELLER, R.H. 1988. VAMP-1: a synaptic vesicle-associated integral membrane protein. *Proc. Natl. Acad. Sci. USA*, **85**: 4538–4542.
 62. VALE, R.D., SCHNAPP, B.J., MITCHISON, T., STEUER, E., REESE, T.S., and SHEETZ, M.P. 1985. Different axoplasmic proteins generate movement in opposite directions along microtubules *in vitro*. *Cell*, **43**: 623–632.
 63. VALE, R.D., SCHNAPP, B.J., REESE, T.S., and SHEETZ, M.P. 1985. Movement of organelles along filaments dissociated from the axoplasm of the squid giant axon. *Cell*, **40**: 449–454.
 64. WATERS, M.G., SERAFINI, T., and ROTHMAN, J.E. 1991. 'Coat-omer': a cytosolic protein complex containing subunits of non-clathrin-coated Golgi transport vesicles. *Nature*, **349**: 248–251.
 65. WEIDMAN, P.J., MELANCON, P., BLOCK, M.R., and ROTHMAN, J.E. 1989. Binding of an N-ethylmaleimide-sensitive fusion protein to Golgi membranes requires both a soluble protein(s) and an integral membrane receptor. *J. Cell Biol.*, **108**: 1589–1596.
 66. WHITEHEART, S.W., GRIFF, I.C., BRUNNER, M., CLARY, D.O., MAYER, T., BUHROW, S.A., and ROTHMAN, J.E. 1993. SNAP family of NSF attachment proteins includes a brain-specific isoform [see comments]. *Nature*, **362**: 353–355.
 67. WIELAND, F.T., GLEASON, M.L., SERAFINI, T.A., and ROTHMAN, J.E. 1987. The rate of bulk flow from the endoplasmic reticulum to the cell surface. *Cell*, **50**: 289–300.
 68. WILSON, D.W., WILCOX, C.A., FLYNN, G.C., CHEN, E., KUANG, W.J., HENZEL, W.J., BLOCK, M.R., ULLRICH, A., and ROTHMAN, J.E. 1989. A fusion protein required for vesicle-mediated transport in both mammalian cells and yeast. *Nature*, **339**: 355–359.
 69. YOSHIDA, A., OHO, C., OMORI, A., KUWAHARA, R., ITO, T., and TAKAHASHI, M. 1992. HPC-1 is associated with synaptotagmin and omega-conotoxin receptor. *J. Biol. Chem.*, **267**: 24925–24928.