

Pavlo O. Kasyanov, Valery S. Mel'nik, Anna M. Piccirillo

**LOCAL SUBDIFFERENTIALS  
AND MULTIVARIATIONAL INEQUALITIES  
IN BANACH AND FRECHET SPACES**

**Abstract.** Some functional-topological concepts of subdifferential and locally subdifferential maps in Frechet spaces are established. Multivariational inequalities with an operator of the pseudo-monotone type, connected with subdifferential maps, are considered.

**Keywords:** local subdifferential, multi-variational inequality, Frechet space.

**Mathematics Subject Classification:** 47J20, 49J40, 58E30, 54C05, 47H04.

## 1. INTRODUCTION AND NOTATION

Subdifferential maps play an important role in the non-smooth analysis and the optimization theory [1–3], in nonlinear boundary value problems for partial differential equations, the theory of control of the distributed systems [4, 5], as well as the theory of differential games and mathematical economy [6, 7]. For basic properties of such maps we refer the reader to [2, 3, 8]. In this paper we will generalize basic properties of subdifferentials and local subdifferentials known for Banach spaces to the case of Frechet spaces.

Let  $X$  be a Frechet space,  $X^*$  its topologically dual (adjoint) space. For  $x \in X$  and  $f \in X^*$ , as usual, the symbol  $\langle f, x \rangle$  stands for the bilinear pairing between  $X$  and  $X^*$ . Assume that  $X$  is endowed with the topology  $\tau$  generated by a family of seminorms  $\{\rho_i\}_{i=1}^{\infty}$  separating points of  $X$ . Recall that the topology  $\tau$  is Hausdorff and metrizable by the metric

$$d(x, y) = \sum_{i=1}^{\infty} 2^{-i} \frac{\rho_i(x - y)}{1 + \rho_i(x - y)}. \quad (1)$$

Observe that  $d(x+h, y+h) = d(x, y)$ ,  $d(\alpha x, \alpha y) < |\alpha|d(x, y)$  if  $|\alpha| > 1$  and  $d(\alpha x, \alpha y) \geq |\alpha|d(x, y)$  if  $|\alpha| \leq 1$ .

Let  $Y$  be a locally convex linear space and  $T : Y \rightarrow X$  be a linear continuous map. Recall that the adjoint (dual) transformation  $T^* : X^* \rightarrow Y^*$  is given by a formula  $\langle x^*, Ty \rangle = \langle T^*x^*, y \rangle$  for  $y \in Y$ ,  $x^* \in X^*$ . For the existence and uniqueness of such transformation, see [10].

Throughout the paper,  $F$  stands for the functional  $F : X \rightarrow \mathbb{R} \cup \{+\infty\}$  and the symbol  $\text{dom } F$  denotes the set  $\{x \in X \mid F(x) < +\infty\}$ .

Given a functional  $F$  and a convex body  $U$  such that  $\text{int}U \subset \text{dom}F$ , a *local subdifferential* of  $F$  at the point  $x_0 \in U \cap \text{dom}F$  is, by definition, the set

$$\partial F(x_0; U) = \{\xi \in X^* \mid \langle \xi, x - x_0 \rangle_X \leq F(x) - F(x_0) \text{ for all } x \in U\}$$

Observe that  $\partial F(x_0; U_1) \supset \partial F(x_0; U_2)$ , if  $U_1 \subset U_2$ . In particular,  $\partial F(x_0; X) = \partial F(x_0) \subset \partial F(x_0; U)$ . The last set is called the *subdifferential* of  $F$  at the point  $x_0$ .

## 2. RESULTS

**Proposition 1.** *Let a functional  $F : X \rightarrow \mathbb{R} \cup \{+\infty\}$  be given. Assume that there are a convex body  $U$  and a point  $x_0 \in \text{int}U$  such that  $\partial F(x_0; U) \neq \emptyset$ . Then the functional  $F$  is weakly lower semicontinuous at  $x_0$ . Moreover, if  $\partial F(x_0; U) \neq \emptyset$  for every  $x \in U$ , then  $F$  is convex on  $U$ .*

*Proof.* Let  $\{x_\alpha\}$  be a net converging to  $\{x_0\}$  and  $W \subset U$  be a neighborhood of  $\{x_0\}$ . Obviously there exists  $\alpha_0$  such that  $x_\alpha \in W$  for  $\alpha \succeq \alpha_0$ . Let  $x^* \in \partial F(x_0; U)$ . For  $\alpha \succeq \alpha_0$ , there is  $\langle x^*, x_\alpha - x_0 \rangle \leq F(x_\alpha) - F(x_0)$ . Passing with  $x_\alpha$  to  $x_0$ , we deduce that  $\varliminf_{\alpha} F(x_\alpha) \geq F(x_0)$ .

Now suppose that  $\partial F(x_0; U) \neq \emptyset$  for an arbitrary  $x_0 \in U$ . Fix  $x_0 \in U$ . For  $x^* \in \partial F(x_0; U)$  and  $x_1, x_2 \in U$ , there is  $F(x_1) - F(x_0) \geq \langle x^*, x_1 - x_0 \rangle$ , and  $F(x_2) - F(x_0) \geq \langle x^*, x_2 - x_0 \rangle$  for all  $x_1, x_2 \in U$ .

Let  $t \in [0, 1]$ . Adding the first inequality multiplied by  $t$  to the second one multiplied by  $1 - t$ , we obtain

$$tF(x_1) + (1 - t)F(x_2) \geq F(x_0) + \langle x^*, tx_1 + (1 - t)x_2 - x_0 \rangle.$$

Since  $U$  is a convex set we can take  $x_0 = tx_1 + (1 - t)x_2$ . □

**Proposition 2.** *Let  $X$  be a Frechet space,  $Y$  a locally convex linear space,  $F : X \rightarrow \mathbb{R} \cup \{+\infty\}$  and  $T : Y \rightarrow X$  a linear continuous map admitting an adjoint map  $T^*$ . Let  $U \subset X$  be a convex body and  $V = T^{-1}(U)$ . Then for every  $Tv \in \text{int}U$ , where  $v \in V$ , there is*

$$\partial(F \circ T)(v; V) = T^*(\partial F(T(v; V)))$$

*Proof.* Let  $x \in \partial F(Tv; U)$ . Obviously,

$$\langle x^*, x - Tv \rangle \leq F(x) - F(T(v)) \quad \text{for every } x \in U.$$

Taking  $x = T(y)$  with  $y \in V$ , we can rewrite the last inequality in the form

$$\langle x^*, Ty - Tv \rangle \leq F(T(y)) - F(T(v)) \quad \text{for every } y \in V$$

or

$$\langle T^*x^*, y - v \rangle \leq (F \circ T)(y) - (F \circ T)(v) \quad \text{for every } y \in V,$$

which means that  $T^*x^* \in \partial(F \circ T)(v; V)$ . Thus  $\partial(F \circ T)(v; V) \supset T^*(\partial F(T(v); V))$ .

To prove the inverse inclusion, take  $y^* \in \partial(F \circ T)(v; V)$ . Clearly

$$\langle y^*, y - v \rangle \leq (F \circ T)(y) - (F \circ T)(v) \quad \text{for every } y \in V.$$

Taking  $x^* \in X^*$  such that  $y^* = T^*x^*$ , we can rewrite the last inequality in the form

$$\langle x^*, Ty - Tv \rangle \leq F(T(y)) - F(T(v)) \quad \text{for every } y \in V$$

or

$$\langle x^*, x - Tv \rangle \leq F(x) - F(T(v)) \quad \text{for every } x \in U,$$

which means that  $y^* = T^*x^* \in T^*(\partial F(T(v); V))$  and this completes the proof.  $\square$

**Theorem 1.** *Let  $U$  be a convex body in  $X$ ,  $F : X \mapsto \mathbb{R} \cup \{+\infty\}$  be a convex functional on  $U$  and a lower semicontinuous functional on  $\text{int } U$  ( $\text{int } U \subset \text{dom } F$ ). Then for every  $x_0 \in \text{int } U$  and every  $h \in X$ , the quantity*

$$D_+F(x_0; h) = \lim_{t \rightarrow 0^+} \frac{F(x_0 + th) - F(x_0)}{t} \quad (2)$$

is finite and the following statements hold true:

- (i) *there exists a counterbalanced (cf. [12]) convex absorbing neighborhood of zero  $\Theta$  ( $x_0 + \Theta \subset \text{int } U$ ) such that for every  $h \in \Theta$*

$$F(x_0) - F(x_0 - h) \leq D_+F(x_0; h) \leq F(x_0 + h) - F(x_0); \quad (3)$$

- (ii) *the functional  $\text{int } U \times X \ni (x; h) \mapsto D_+F(x; h)$  is upper semicontinuous;*  
 (iii) *the functional  $D_+F(x_0; \cdot) : X \mapsto \mathbb{R}$  is positively homogeneous and semiadditive for every  $x_0 \in \text{int } U$ ;*  
 (iv) *there exist a neighborhood  $O(h_0)$  and a constant  $c_1 > 0$  such that for every  $x_0 \in \text{int } U$  and every  $h_0 \in X$ ,*

$$|D_+F(x_0; h) - D_+F(x_0; h_0)| \leq c_1 d(h, h_0) \quad \text{for every } h \in O(h_0).$$

*Proof.* First we introduce some auxiliary statements.

**Claim 1.** *The functional  $F$  is locally upper bounded on  $\text{int } U$ , that is for every  $x_0 \in \text{int } U$  there exist positive constants  $r$  and  $c$  such that  $F(x) \leq c$ , for each  $x \in B_r(x_0)$ , where  $B_r(x_0) = \{x \in X \mid d(x, x_0) < r\}$ .*

*Proof of Claim 1.* For arbitrary  $x_0 \in \text{int } U$  there exists  $\varepsilon_1 > 0$  such that  $B_{2\varepsilon_1}(x_0) \subset \text{int } U \subset \text{dom } F$ , hence  $\overline{B_{\varepsilon_1}(x_0)} \subset B_{2\varepsilon_1}(x_0) \subset \text{dom } F$ . Since  $F$  is lower semicontinuous, then for each  $n = 1, 2, \dots$  the set

$$A_n = \{x \in \overline{B_{\varepsilon_1}(x_0)} \mid F(x) \leq n\}$$

is closed in  $X$  and

$$\bigcup_{n=1}^{+\infty} A_n = \overline{B_{\varepsilon_1}(x_0)} \subset \text{dom } F.$$

Since the metric space  $(\overline{B_{\varepsilon_1}(x_0)}, d)$  is complete, due to the Baire Category Theorem, there exists  $n_0 \in \mathbb{N}$  such that  $\text{int}A_{n_0} \neq \emptyset$  in  $\overline{B_{\varepsilon_1}(x_0)}$ . We now prove that  $\text{int}A_{n_0} \neq \emptyset$  in  $X$ . Since  $\text{int}A_{n_0} \neq \emptyset$  in  $\overline{B_{\varepsilon_1}(x_0)}$ , we conclude that there exist  $x_1 \in \text{int}A_{n_0}$  and  $\varepsilon_2 > 0$  such that the following equality holds true:

$$A_{n_0} \supset \{x \in \overline{B_{\varepsilon_1}(x_0)} \mid d(x, x_1) < \varepsilon_2\} = \overline{B_{\varepsilon_1}(x_0)} \cap B_{\varepsilon_2}(x_1) \neq \emptyset.$$

Thus the following two cases are possible:

- 1)  $B_{\varepsilon_1}(x_0) \cap B_{\varepsilon_2}(x_1) \neq \emptyset$ ;
- 2)  $B_{\varepsilon_1}(x_0) \cap B_{\varepsilon_2}(x_1) = \emptyset$ ,  $\partial B_{\varepsilon_1}(x_0) \cap B_{\varepsilon_2}(x_1) \neq \emptyset$ .

In the first case, the set  $B_{\varepsilon_1}(x_0) \cap B_{\varepsilon_2}(x_1)$  is open in topology  $\tau$ ; therefore, there exist  $x_2 \in X$  and  $\varepsilon_3 > 0$  such that  $B_{\varepsilon_3}(x_2) \subset B_{\varepsilon_1}(x_0) \cap B_{\varepsilon_2}(x_1) \subset A_{n_0}$ . Thus, for each  $x \in B_{\varepsilon_3}(x_2)$ , there is  $F(x) \leq n_0$ . Hence  $x_2 \in \text{int}A_{n_0}$  in  $X$ .

In the second case, for an arbitrary  $x \in \partial B_{\varepsilon_1}(x_0) \cap B_{\varepsilon_2}(x_1)$ , there exists  $\{x_n\}_{n \geq 1} \subset B_{\varepsilon_1}(x_0)$  such that  $x_n \rightarrow x$  as  $n \rightarrow +\infty$ . Since  $x \in B_{\varepsilon_2}(x_1)$ , then there exists  $N$  such that for each  $n \geq N$ ,  $x_n \in B_{\varepsilon_2}(x_1)$ . Therefore,  $x_n \in B_{\varepsilon_1}(x_0) \cap B_{\varepsilon_2}(x_1) \neq \emptyset$ , and we may proceed further as in the first case. Thus  $\text{int}A_{n_0} \neq \emptyset$  in  $X$ .

Now we show that the functional  $F$  is upper bounded in some neighborhood of  $x_0$ . Let  $x_2 \neq x_0$ ,  $y = x_2 + \frac{x_0 - x_2}{1 - \lambda}$ , where  $\lambda = \frac{\varepsilon_1/d(x_2, x_0)}{1 + \varepsilon_1/d(x_2, x_0)}$ . Therefore,

$$y = x_0 + \frac{\varepsilon_1}{d(x_2, x_0)}(x_0 - x_2),$$

$$d(y, x_0) = d\left(\frac{\varepsilon_1}{d(x_0, x_2)}(x_0 - x_2), 0\right) < \frac{\varepsilon_1}{d(x_0, x_2)}d(x_0, x_2) = \varepsilon_1,$$

that is  $y \in B_{\varepsilon_1}(x_0) \subset \text{dom}F$ . For an arbitrary  $x \in B_{\lambda\varepsilon_3}(x_0)$ , we consider  $z = (x + \lambda x_2 - x_0)/\lambda = (x - (1 - \lambda)y)/\lambda$ . Since  $0 < \lambda < 1$ , we conclude that

$$d(z, x_2) = d\left(x_2 + \frac{x - x_0}{\lambda}, x_2\right) = d\left(\frac{x - x_0}{\lambda}, 0\right) < \frac{1}{\lambda}d(x, x_0) < \frac{\lambda\varepsilon_3}{\lambda} = \varepsilon_3,$$

hence  $z \in B_{\varepsilon_3}(x_2)$ , and  $F(z) \leq n_0$ . Due to convexity of  $F$ , there is

$$F(x) = F(\lambda z + (1 - \lambda)y) \leq \lambda F(z) + (1 - \lambda)F(y) \leq n_0 + (1 - \lambda)F(y).$$

From this we conclude that  $F$  is upper bounded in the neighborhood  $B_r(x_0)$  with  $r = \lambda\varepsilon_3$  and  $c = n_0 + (1 - \lambda)F(y)$ .  $\square$

**Claim 2.** *The functional  $F$  is locally Lipschitzean on  $\text{int}U$ , i.e., for every  $x_0 \in \text{int}U$  there exist  $r_1 > 0$  and  $c_1 > 0$  such that*

$$|F(x) - F(y)| \leq c_1 d(x, y) \quad \text{for all } x, y \in B_{r_1}(x_0).$$

*Proof of Claim 2.* The local upper boundedness of the functional  $F$  on  $\text{int } U$  follows from Claim 1. Therefore, for every  $x_0 \in \text{int } U$  there exist  $r > 0$  and  $c > 0$  such that  $F(x) \leq c$  for every  $x \in B_r(x_0)$ .

For an arbitrary  $x \in B_r(x_0)$  ( $x \neq x_0$ ) and  $t = \frac{d(x, x_0)}{r + d(x, x_0)}$ , we put

$$y = \frac{x_0 + (t-1)x}{t} = x_0 + \frac{1-t}{t}(x_0 - x),$$

where  $t \in (0, 1)$ . Then

$$d(y, x_0) = d\left(\frac{1-t}{t}(x_0 - x), 0\right) = d\left(\frac{r}{d(x, x_0)}(x_0 - x), 0\right) < \frac{r}{d(x, x_0)}d(x, x_0) = r,$$

i.e.,  $F(y) \leq c$ . Due to convexity of  $F$ ,

$$F(x_0) = F(ty + (1-t)x) \leq tF(y) + (1-t)F(x) \leq tc + (1-t)F(x),$$

or  $(1-t)F(x_0) \leq t(c - F(x_0)) + (1-t)F(x)$ . Hence

$$F(x_0) - F(x) \leq \frac{t}{1-t}(c - F(x_0)) = \frac{(c - F(x_0))}{r}d(x, x_0). \quad (4)$$

Now let  $z = \frac{x - (1-\tau)x_0}{\tau} = x_0 + \frac{x - x_0}{\tau}$ , where  $\tau = \frac{d(x, x_0)}{r} \in (0, 1)$ . Then

$$d(z, x_0) = d\left(\frac{x - x_0}{\tau}, 0\right) < \frac{1}{\tau}d(x, x_0) = r,$$

i.e.,  $F(z) \leq c$ , and since  $F$  is convex, we obtain

$$F(x) = F(\tau z + (1-\tau)x_0) \leq \tau F(z) + (1-\tau)F(x_0) \leq \tau c + (1-\tau)F(x_0)$$

or

$$F(x) - F(x_0) \leq \tau(c - F(x_0)) = \frac{c - F(x_0)}{r}d(x, x_0). \quad (5)$$

Relations (4) and (5) imply the following estimate

$$|F(x) - F(x_0)| \leq \frac{c - F(x_0)}{r}d(x, x_0). \quad (6)$$

Now we show that the Lipschitz condition holds true for  $F$  on  $B_{\varepsilon_1}(x_0)$  with  $\varepsilon_1 = r/3$ . Hence in view of (6), for all  $x_1, x_2 \in B_{3\varepsilon_1}(x_0)$   $F(x_1) \leq c$ ,  $F(x_2) \leq c$ . If  $x_1 \in B_{\varepsilon_1}(x_0)$ , then  $B_{2\varepsilon_1}(x_1) \subset B_r(x_0)$ , that is  $x_1 \in \text{int } U$ . Therefore, from (6) we obtain

$$|F(x) - F(x_1)| \leq \frac{c - F(x_1)}{2\varepsilon_1}d(x, x_1) \quad \text{for every } x \in B_{2\varepsilon_1}(x_1). \quad (7)$$

In particular, inequality (7) is valid for an arbitrary element of  $B_{\varepsilon_1}(x_0)$ . Further, due to (6)

$$\begin{aligned} -F(x_1) &\leq (c - F(x_0)) + |F(x_1) - F(x_0)| \leq \\ &\leq (c - F(x_0)) + \frac{c - F(x_0)}{r}d(x_1, x_0) < 2(c - F(x_0)). \end{aligned}$$

From the last relation, using (7), we finally obtain

$$|F(x_2) - F(x_1)| \leq \frac{c - F(x_0)}{\varepsilon_1} d(x_2, x_1) \quad \text{for all } x_1, x_2 \in B_{\varepsilon_1}(x_0),$$

i.e.,  $c_1 = \frac{c - F(x_0)}{\varepsilon_1}$ ,  $r_1 = \varepsilon_1$ . □

Now we continue to prove Theorem 1. Let  $x_0 \in \text{int } U$  and  $B_r(x_0) = x_0 + B_r(0)$ . Then due to Claims 1 and 2 the upper boundness and the Lipschitz condition for  $F$  on  $B_r(x_0)$  follow. We recall that, unlike in the case of a Banach space,  $B_r(0)$  is not absolutely convex, but at the same time there exists a convex absorbing counterbalanced set  $\Theta = \Theta(x_0)$  in a basis of topology  $\tau$ , such that  $\Theta \subset B_r(0)$ . Then  $F(x) \leq c$ , for every  $x \in x_0 + \Theta$ ,

$$|F(x_1) - F(x_2)| \leq c_1 d(x_1, x_2) \quad \text{for all } x_1, x_2 \in x_0 + \Theta. \quad (8)$$

For each  $u \in X$  there exists  $t = t(u) > 0$  such that  $t^{-1}u \in \Theta$  (if  $u \in \Theta$ , then we take  $t = 1$ ). So for each  $\tau \in (0, t^{-1}]$  the element  $\tau u \in \Theta$ , as  $t\Theta \subset \frac{1}{\tau}\Theta$ . Further, due to convexity of  $F$ , for every  $\tau_1, \tau_2 \in \mathbb{R}$  such that  $0 < \tau_1 \leq \tau_2 \leq t^{-1}$ , there follows:

$$\begin{aligned} F(x_0 + \tau_1 u) - F(x_0) &= F\left(x_0 \left(1 - \frac{\tau_1}{\tau_2}\right) + (x_0 + \tau_2 u) \frac{\tau_1}{\tau_2}\right) - F(x_0) \leq \\ &\leq \left(1 - \frac{\tau_1}{\tau_2}\right) F(x_0) + \frac{\tau_1}{\tau_2} F(x_0 + \tau_2 u) - F(x_0) = \\ &= \frac{\tau_1}{\tau_2} (F(x_0 + \tau_2 u) - F(x_0)). \end{aligned}$$

Hence the function  $\tau \mapsto \frac{F(x_0 + \tau u) - F(x_0)}{\tau}$  monotonely decreases as  $\tau \rightarrow 0+$ .

For each  $u \in \Theta$ , the quantity  $D_+ F(x_0; u)$  is finite. In fact,  $\alpha u \in \Theta$  for every  $\alpha$  such that  $|\alpha| \leq 1$ , therefore

$$\begin{aligned} D_+ F(x_0; u) &= \inf_{\tau > 0} \frac{F(x_0 + \tau u) - F(x_0)}{\tau} \leq \\ &\leq F(x_0 + u) - F(x_0) < +\infty \text{ as } x_0 + u \in B_r(x_0) \subset \text{int } U \subset \text{dom } F. \end{aligned}$$

On the other hand, for every  $\tau \in (0, 1)$   $x_0 = \frac{1}{1+\tau}(x_0 + \tau u) + \frac{\tau}{1+\tau}(x_0 - u)$ , i.e.,  $-u \in \Theta$ , and moreover, for each  $\tau \in (0, 1)$ ,

$$-\infty < F(x_0) - F(x_0 - u) \leq \frac{F(x_0 + \tau u) - F(x_0)}{\tau}.$$

Thus, for each  $u \in \Theta$ ,

$$-\infty < F(x_0) - F(x_0 - u) \leq D_+ F(x_0; u) \leq F(x_0 + u) - F(x_0) < +\infty,$$

i.e.,  $D_+ F(x_0; u) \in \mathbb{R}$  for every  $u \in \Theta$ . The validity of (3) follows from these facts.

From (2) we immediately obtain

$$D_+ F(x_0; \alpha u) = \alpha D_+ F(x_0; u) \quad \text{for all } \alpha > 0 \text{ and } u \in X, \quad (9)$$

and since the set  $\Theta$  is absorbing, then for each  $u \in X$  there is  $\alpha > 0$  such that  $\alpha u \in \Theta$ . Then from (9) we obtain

$$D_+F(x_0; u) \in \mathbb{R} \quad \text{for all } x_0 \in \text{int } U \text{ and } u \in X.$$

Now taking  $t > 0$ , we consider the function

$$\text{int } U \times \Theta \ni (x, h) \mapsto F_t(x; h) = \frac{F(x + th) - F(x)}{t}. \quad (10)$$

**Claim 3.** For each pair  $(x_0; h_0) \in \text{int } U \times X$  there exists  $l > 0$  such that for each  $t \in (0, l)$  the function  $F_t(\cdot; \cdot)$  is continuous at the point  $(x_0; h_0)$ .

*Proof of Claim 3.* Let  $h_0 \in X$  be arbitrary,  $x_0 \in \text{int } U$  (the set  $\Theta = \Theta(x_0)$  is defined above), then there exists  $t_0 > 0$  such that  $h_0 \in t_0\Theta$ . For  $t \in (0, l)$ , putting  $l = \min(\frac{1}{2t_0}, 1)$ , we consider the function  $F_t(x; h)$  in a neighborhood of the point  $(x_0, h_0)$ :

$$|F_t(x; h) - F_t(x_0; h_0)| = \frac{1}{t} \left| [F(x_0 + th_0 + (x - x_0) + t(h - h_0)) - F(x_0 + th_0)] + [F(x_0) - F(x)] \right| \quad (11)$$

If we take  $x \in x_0 + \frac{1}{4}\Theta$ ,  $h \in h_0 + \frac{1}{4}\Theta$ , then

$$x_0 + th_0 \in x_0 + \frac{1}{2}\Theta \subset x_0 + \Theta, \quad t(h - h_0) \in \frac{1}{4}\Theta,$$

$$(x - x_0) + t(h - h_0) \in \frac{1}{2}\Theta, \quad x_0 + th_0 + (x - x_0) + t(h - h_0) \in x_0 + \Theta.$$

From (11) using (8), we derive

$$|F_t(x; h) - F_t(x_0; h_0)| \leq \frac{c_1}{t} (d(x + th, x_0 + th_0) + d(x, x_0)) \rightarrow 0$$

as  $x \rightarrow x_0$ ,  $h \rightarrow h_0$ . □

Claim 3 implies the upper semicontinuity of the map

$$\text{int } U \times X \ni (x; h) \mapsto D_+F(x; h) = \inf_{t>0} F_t(x; h) = \inf_{t \in (0, l)} F_t(x; h),$$

since it is a “pointwise infimum” of continuous functions. The positive homogeneity of  $D_+F(x_0; \cdot)$  is obvious. Now we show that this map is semiadditive. Indeed, for all  $v_1, v_2 \in X$

$$\begin{aligned} D_+F(x_0; v_1 + v_2) &= \inf_{t>0} \frac{F(x_0 + t(v_1 + v_2)) - F(x_0)}{t} = \\ &= \lim_{t \rightarrow 0^+} \frac{2F(\frac{x_0 + tv_1}{2} + \frac{x_0 + tv_2}{2}) - 2F(x_0)}{t} \leq \\ &\leq \lim_{t \rightarrow 0^+} \frac{F(x_0 + tv_1) - F(x_0)}{t} + \lim_{t \rightarrow 0^+} \frac{F(x_0 + tv_2) - F(x_0)}{t} = \\ &= D_+F(x_0; v_1) + D_+F(x_0; v_2). \end{aligned}$$

In order to complete the proof it suffices to show that the map  $D_+F(x_0; \cdot)$  satisfies (iv). From semiadditivity it follows that

$$|D_+F(x_0; h) - D_+F(x_0; h_0)| \leq \max\{D_+F(x_0; h - h_0), D_+F(x_0; h_0 - h)\} \leq c_1 d(h, h_0)$$

for any  $h \in h_0 + \frac{1}{4}\Theta$ . This completes the proof of Theorem 1.  $\square$

**Definition 1.** We call a set  $B \subset X^*$  bounded in the  $\sigma(X^*; X)$  topology ( $*$ -bounded), if  $\sup_{y \in B} |\langle y, x \rangle_X| < +\infty$  for each  $x \in X$ .

It is obvious that each bounded set in  $X^*$  is  $*$ -bounded.

**Definition 2.** A multivalued map  $A : X \rightrightarrows X^*$  is called:

- a)  $*$ -bounded, if for any bounded set  $B$  in  $X$  the image  $A(B)$  is  $*$ -bounded in  $X^*$ ;
- b)  $*$ -upper semicontinuous, if for any set  $B$  open in the  $\sigma(X^*, X)$  topology the set  $A_M^{-1}(B) = \{x \in X \mid A(x) \subset B\}$  is open in  $X$ ;
- c) upper hemicontinuous, if the function

$$X \ni x \mapsto [A(x), y]_+ = \sup_{d \in A(x)} \langle d, y \rangle_X$$

is upper semicontinuous for each  $y \in X$ .

Let us note that c) follows from b).

**Theorem 2.** Let  $U$  be a convex body and  $\text{int} U \subset \text{dom} F$ , where  $F : X \rightarrow \overline{\mathbb{R}}$  is a convex functional on  $U$  and a semicontinuous function on  $\text{int} U$ . Then:

- i)  $\partial F(x; U)$  is a nonempty convex compact set for every  $x \in \text{int} U$  in the  $\sigma(X^*; X)$  topology;
- ii)  $\partial F(\cdot; U) : U \rightrightarrows X^*$  is a monotone map (on  $U$ );
- iii) the map  $\text{int} U \ni x \mapsto \partial \varphi(x; U) \subset X^*$  is  $*$ -upper semicontinuous (on  $\text{int} U$ ) and

$$[\partial \varphi(x_0; U), h]_+ = D_+ \varphi(x_0; h) \quad \text{for all } h \in X \text{ and } x_0 \in \text{int} U. \quad (12)$$

*Proof.* First we prove condition ii). Let  $x_1, x_2 \in U$  and  $\xi_i \in \partial F(x_i; U)$ ,  $i = 1, 2$ . Then

$$F(x_2) - F(x_1) \geq \langle \xi_1, x_2 - x_1 \rangle_X, \quad F(x_1) - F(x_2) \geq \langle \xi_2, x_1 - x_2 \rangle_X.$$

Adding the first inequality to the second, we obtain

$$\langle \xi_1 - \xi_2, x_1 - x_2 \rangle_X \geq 0,$$

or

$$[\partial F(x_1; U), x_1 - x_2]_- \geq [\partial F(x_1; U), x_1 - x_2]_+ \quad \text{for all } x_1, x_2 \in U.$$

The last relation proves the monotonicity on  $U$ . Convexity and weak star closure are obvious. Let us prove nonemptiness. Let us set an arbitrary  $x, h \in \text{int} U$  and

consider the real convex function  $\varphi(t) = F(x + t(h - x))$  defined on  $[0, 1]$ . So there exist  $\varphi(t-)$ ,  $\varphi(t+)$  such that

$$\varphi(t-) \leq \varphi(t+) = \lim_{t \rightarrow +0} \frac{\varphi(t) - \varphi(0)}{t} = D_+\varphi(x; x - h),$$

or

$$D_-\varphi(x; x - h) \leq D_+\varphi(x; x - h),$$

where  $D_-\varphi(x; v) = -D_+\varphi(x; -v)$ . From Theorem 1 it follows that

$$\frac{\varphi(\alpha) - \varphi(0)}{\alpha} \leq \varphi(1) - \varphi(0) \quad \text{for every } \alpha \in (0, 1)$$

or

$$D_+F(x; h - x) \leq F(h) - F(x) \quad \text{for all } x, h \in \text{int } U. \quad (13)$$

**Claim 4.** For arbitrary  $x \in \text{int } U$  there exists  $\xi(x) \in X^*$  such that

$$D_-F(x; h) \leq \langle \xi(x), h \rangle_X \leq D_+F(x; h) \quad \text{for every } h \in X.$$

*Proof of Claim 4.* Let us fix  $h_0 \in X$  and consider the one-dimensional subspace  $X_0 = \{\alpha h_0 \mid \alpha \in \mathbb{R}\}$ . Let us choose an element  $\xi \in X^*$  satisfying the following condition

$$\langle \xi, \alpha h_0 \rangle_X = D_+F(x, \alpha h_0), \quad \alpha \geq 0$$

(We remark that since  $x$  is an interior point of  $U$ , then due to Theorem 1 for every  $h \in X$  there exists  $D_+F(x; h)$ ). It is possible to choose  $\xi$  in such a way, since  $X \ni h \mapsto D_+F(x; h)$  is a positively homogeneous functional. Further, taking into account the semiadditivity of  $X \ni h \mapsto D_+F(x; h)$ , we obtain

$$0 = D_+F(x; h - h) \leq D_+F(x; h) + D_+F(x; -h)$$

or

$$-D_+F(x; h) \leq D_+F(x; -h). \quad (14)$$

Then for  $\alpha < 0$ , from (14), the following relation follows:

$$\begin{aligned} \langle \xi, \alpha h_0 \rangle_X &= \alpha D_+F(x; h_0) = -|\alpha| D_+F(x; h_0) \leq \\ &\leq |\alpha| D_+F(x; -h_0) = D_+F(x; -|\alpha| h_0) = D_+F(x; \alpha h_0). \end{aligned}$$

Since  $\langle \xi, v \rangle_X \leq D_+F(x; v)$  for each  $v \in X_0$  and  $X \ni h \mapsto D_+F(x; h)$  is a continuous positively homogeneous semiadditive functional, then according to the Hahn-Banach Theorem there exists  $\zeta \in X^*$  such that  $\langle \zeta, h \rangle_X \leq D_+F(x; h)$  for each  $h \in X$  and  $\langle \zeta, h_0 \rangle_X = \langle \xi, h_0 \rangle_X$ . Hence we obtain  $\langle \zeta, -h \rangle_X \leq D_+F(x; -h)$  and

$$\langle \zeta, h \rangle_X = -\langle \zeta, -h \rangle_X \geq -D_+F(x; -h) = D_-F(x; h) \quad \text{for every } h \in X.$$

The last relation proves the required inequality.  $\square$

Claim 4 and inequality (13) guarantee the existence of  $\xi(x) \in X^*$  such that

$$\langle \xi(x), h - x \rangle_X \leq D_+F(x; h - x) \leq F(h) - F(x) \quad \text{for every } h \in U,$$

i.e.,  $\xi(x) \in \partial F(x, U)$ , and hereby the nonemptiness of  $\partial F(x, U)$  is proved.

**Claim 5.** For every  $x_0 \in \text{int } U$ , the following inequality holds true:

$$\partial\varphi(x_0; U) = \{p \in X^* \mid \langle p, h \rangle_X \leq D_+\varphi(x_0; h) \text{ for every } h \in X\}.$$

*Proof of Claim 5.* Let  $p \in \partial F(x_0; U)$ . Then there exists an open convex set  $V$  containing zero such that  $x_0 + V \subset \text{int } U$  and

$$\langle p, h \rangle_X \leq F(x_0 + h) - F(x_0) \quad \text{for every } h \in V.$$

Hence,

$$\langle p, h \rangle_X \leq \frac{F(x_0 + th) - F(x_0)}{t} \quad \text{for every } t \in (0, 1).$$

Due to Theorem 1,

$$\langle p, h \rangle_X \leq \inf_{t>0} \frac{F(x_0 + th) - F(x_0)}{t} = D_+F(x_0; h) \quad \text{for every } h \in V.$$

Since the set  $V$  is absorbing and functions

$$X \ni h \mapsto D_+F(x; h), \quad X \ni h \mapsto \langle p, h \rangle_X$$

are positively homogeneous, then

$$\langle p, h \rangle_X \leq D_+F(x_0; h) \quad \text{for every } h \in X.$$

On the other hand, let for every  $h \in X$  the relation  $\langle p, h \rangle_X \leq D_+F(x_0; h)$  hold true. Due to Theorem 1, there follows the existence of a counterbalanced convex absorbing neighborhood of zero  $\Theta$  ( $x_0 + \Theta \subset \text{int } U$ ) such that

$$D_+F(x_0; v) \leq F(x_0 + v) - F(x_0) \quad \text{for every } v \in \Theta.$$

Let us fix an arbitrary  $h \in U \cap \text{dom } F$ . Then there is  $\alpha \in (0, 1)$  such that  $\alpha(h - x_0) \in \Theta$ . Therefore,

$$\begin{aligned} \alpha \cdot \langle p, h - x_0 \rangle_X &= \langle p, \alpha(h - x_0) \rangle_X \leq D_+F(x_0; \alpha(h - x_0)) \leq F(x_0 + \alpha(h - x_0)) - \\ &\quad - F(x_0) \leq \alpha F(h) + (1 - \alpha)F(x_0) - F(x_0) = \alpha(F(h) - F(x_0)). \end{aligned}$$

Hence we obtain that  $\langle p, h - x_0 \rangle_X \leq F(h) - F(x_0)$  for each  $h \in U \cap \text{dom } F$ , and for this reason  $\langle p, h - x_0 \rangle_X \leq F(h) - F(x_0)$  for each  $h \in U$ . Hence  $p \in \partial F(x_0; U)$ .  $\square$

By Claim 5, it immediately follows that

$$[\partial F(x_0; U), h]_+ \leq D_+F(x_0; h) \quad \text{for every } h \in X,$$

that is, due to Claim 5,

$$\begin{aligned} & \{p \in X^* \mid \langle p, h - x_0 \rangle_X \leq [\partial F(x_0; U), h - x_0]_+ \text{ for every } h \in X\} \subset \\ & \subset \{p \in X^* \mid \langle p, h \rangle_X \leq D_+ F(x_0; h - x_0) \text{ for every } h \in X\} = \partial F(x_0; U). \end{aligned}$$

On the other hand, every element  $p \in \partial F(x_0; U)$  satisfies the condition

$$\langle p, h \rangle_X \leq [\partial F(x_0; U), h]_+ \text{ for every } h \in X,$$

which proves the inverse inclusion. Therefore, equality (12) holds.

Further, due to (12) and Theorem 1,  $\partial F(\cdot; U)$  is upper hemicontinuous on  $\text{int } U$ . Moreover, the boundedness of  $\partial F(x_0; U)$  follows from the estimate

$$[\partial F(x_0; U), h]_+ = D_+ F(x_0, h) \leq c_1 d(h, 0) \text{ for every } h \in \Theta,$$

where  $\Theta$  is absorbing. So, by virtue of the Banach-Alaoglu Theorem (cf. [10]),  $\partial F(x_0; U)$  is a compact set in the  $\sigma(X^*, X)$  topology. Under these conditions, upper hemicontinuity of the map  $\partial F(\cdot; U)$  and the Castaing Theorem (cf. [2]) imply \*-upper semicontinuity of  $\partial F(\cdot; U)$  on  $\text{int } U$ . This completes the proof of Theorem 2.  $\square$

**Theorem 3.** Let  $F_1, F_2 : X \rightarrow \overline{\mathbb{R}}$  and  $U = U_1 \cap U_2$ , where  $\text{int } U \neq \emptyset$ ,  $U_1, U_2$  are convex sets and

$$\partial F_1(x_1; U_1) \neq \emptyset, \partial F_2(x_2; U_2) \neq \emptyset \text{ for all } x_1 \in U_1, x_2 \in U_2.$$

Then  $\partial F(x; U) \neq \emptyset$  for every  $x \in U$ , where  $F = F_1 + F_2$ , and

$$\partial F(x; U) = \partial F_1(x; U) + \partial F_2(x; U) \text{ for every } x \in \text{int } U.$$

*Proof.* Suppose that  $x \in U$ . It is clear that

$$\partial F(x; U) \supset \partial F_1(x; U) + \partial F_2(x; U) \supset \partial F_1(x; U_1) + \partial F_2(x; U_2) \neq \emptyset.$$

In order to complete the proof, it is necessary to show that for every  $x \in \text{int } U$  and for every  $h \in X$  the following equality is fulfilled:

$$D_+ F(x; h) = D_+ F_1(x; h) + D_+ F_2(x; h). \quad (15)$$

Indeed, since functions  $F, F_1, F_2$  satisfy assumptions of Proposition 1, then all conditions of Theorem 2 hold true for them as well. Thus, due to equality (12) and [11, Proposition 1],

$$\begin{aligned} [\partial F(x; U), h]_+ &= D_+ F(x; h) = D_+ F_1(x; h) + D_+ F_2(x; h) = \\ &= [\partial F_1(x; U), h]_+ + [\partial F_2(x; U), h]_+ = \\ &= [\partial F_1(x; U) + \partial F_2(x; U), h]_+ \text{ for all } x \in \text{int } U \text{ and } h \in X. \end{aligned}$$

Hence

$$\partial F(x; U) = \partial F_1(x; U) + \partial F_2(x; U) \text{ for every } x \in \text{int } U.$$

Now we prove (15). For functions  $F, F_1, F_2$ , due to Proposition 1, Theorem 1 holds true. Consequently, for all  $x \in \text{int } U$  and  $h \in X$ , we obtain

$$\begin{aligned} D_+F(x; h) &= \lim_{t \rightarrow 0+} \frac{F(x + th) - F(x)}{t} = \\ &= \lim_{t \rightarrow 0+} \frac{F_1(x + th) - F_1(x) + F_2(x + th) - F_2(x)}{t} = \\ &= \lim_{t \rightarrow 0+} \frac{F_1(x + th) - F_1(x)}{t} + \lim_{t \rightarrow 0+} \frac{F_2(x + th) - F_2(x)}{t} = \\ &= D_+F_1(x; h) + D_+F_2(x; h). \end{aligned}$$

This completes the proof of Theorem 3.  $\square$

**Definition 3.** Suppose that  $U$  is a convex body. The functional  $F : X \mapsto \mathbb{R} \cup \{+\infty\}$  ( $\text{int } U \subset \text{dom } F$ ) is said to be upper bounded on  $\text{int } U$  if for every bounded set  $B \subset \text{int } U$  the image  $F(B)$  is upper bounded in  $\mathbb{R}$ .

The following result is new even in the case of  $X$  being a Banach space.

**Theorem 4.** Let  $F : X \mapsto \mathbb{R}$  be a convex lower semicontinuous functional. Then the following statements are equivalent:

- a)  $F$  is an upper bounded functional on  $X$  ;
- b) a multivalued map  $\partial F(\cdot) = \partial F(\cdot; X)$  is  $*$ -bounded on  $X$ .

*Proof.* The following statements are true.

**Claim 6.** If  $B$  is a bounded set in  $X$  and  $C$  is a  $*$ -bounded set in  $X^*$ , then the quantity  $\sup_{x \in B} \sup_{p \in C} |\langle p, x \rangle_X|$  is finite.

*Proof of Claim 6.* Let  $\rho(x) = \sup_{p \in C} |\langle p, x \rangle_X|$ .  $*$ -boundedness of  $C$  implies that the given functional is well defined on  $X$ . We remark that  $\rho(-x) = \rho(x)$  for  $x \in X$ . Moreover,  $\rho$  is convex positively homogeneous and lower semicontinuous as the supremum of convex positively homogeneous continuous functionals. Hence, due to Claim 2,  $\rho$  is continuous on  $X$ , i.e.,  $\rho$  is a continuous seminorm on  $X$ . By Theorem V.23 in [12], the boundedness of  $B$  in  $X$  implies that  $\sup_{x \in B} \sup_{p \in C} |\langle p, x \rangle_X| = \sup_{x \in B} \rho(x) < +\infty$ .  $\square$

**Definition 4.** Let  $X$  be a separable locally convex topological space,  $U \subset X$  be an unbounded convex body. Then the functional  $F : U \mapsto \mathbb{R} \cup \{+\infty\}$  is called coercive on  $U$  if  $F(x) \rightarrow +\infty$  as  $\rho(x) \rightarrow +\infty$ ,  $x \in U$ , where  $\rho$  an arbitrary continuous seminorm on  $X$ .

**Claim 7.** Let  $B \subset X$  be a nonempty set satisfying one of the two conditions:

- (i)  $B$  is bounded,
- (ii)  $F$  is coercive on  $B$ .

Then  $\inf_{x \in B} F(x) > -\infty$ .

*Proof of Claim 7.* For some integer  $n$ , we consider the following set:

$$A_n = \{x \in B \mid F(x) \leq n\} \neq \emptyset.$$

The boundedness of  $A_n$  follows from the boundedness of  $B$  or coercivity of  $F$ . Indeed, if the set  $A_n$  is unbounded, then there exists a continuous seminorm  $\rho$  and a sequence  $\{x_n\}_{n \geq 1} \subset B$  such that  $\rho(x_n) \rightarrow +\infty$ . Thus we obtain  $F(x_n) \rightarrow +\infty$ , and this fact contradicts the construction of  $A_n$ . Therefore, taking into account Theorem 2 and Claim 6 with  $C = \{p\}$ ,  $p \in \partial F(\bar{0})$ , we deduce that  $\inf_{x \in B} F(x) \geq F(\bar{0}) - \sup_{x \in B} |\langle p, x \rangle| > -\infty$ . This completes the proof.  $\square$

We continue with the proof of Theorem 4. Let the set  $B$  be bounded in  $X$ . First we assume that the multivalued map  $\partial F(\cdot)$  is  $*$ -bounded on  $X$ . Then, by definition of a subdifferential,

$$F(x_0) - F(x) \geq \langle p_x, x_0 - x \rangle_X \quad \text{for all } x \in B \text{ and } p_x \in \partial F(x).$$

Whence for all  $x \in B$  and  $p_x \in \partial F(x)$ , we obtain

$$\begin{aligned} F(x) &\leq F(x_0) + \langle p_x, x - x_0 \rangle_X \leq |F(x_0)| + \sup_{p \in \partial F(B)} |\langle p, x - x_0 \rangle_X| \leq \\ &\leq |F(x_0)| + \sup_{x \in x_0 + B} \sup_{p \in \partial F(B)} |\langle p, x \rangle_X|. \end{aligned}$$

Claim 6 and the fact that  $x_0 + B$  is the bounded set in  $X$  yield

$$\sup_{x \in x_0 + B} \sup_{p \in \partial F(B)} |\langle p, x \rangle_X| < +\infty.$$

Moreover, let the functional  $F$  be upper bounded. Then, due to Theorem 2, for every  $u \in X$  there is

$$\sup_{p \in \partial F(B)} |\langle p, u \rangle_X| = \sup_{x \in B} \sup_{p \in \partial F(x)} \langle p, u \rangle_X = \sup_{x \in B} [\partial F(x), u]_+ = \sup_{x \in B} D_+ F(x; u).$$

Further, from Theorem 1 we infer that

$$\sup_{x \in B} D_+ F(x; u) \leq \sup_{x \in B} (F(x + u) - F(x)) \leq \sup_{x \in B + u} F(x) - \inf_{x \in B} F(x) =: I.$$

Since  $B$ ,  $B + u$  are bounded sets in  $X$ , then (due to Claim 7 and the definition of an upper bounded functional) the quantity  $I$  is finite. Consequently,  $\sup_{p \in \partial F(B; U)} \langle p, u \rangle_X < +\infty$  for every  $u \in X$ . Hence, the set  $\partial F(B)$  is  $*$ -bounded.  $\square$

**Remark 1.** For an arbitrary multivalued map  $A : Y \subset X \rightrightarrows X^*$ ,  $coA$  and  $\overline{co}A$  stand for multivalued maps defined as follows:  $coA(y) := co(A(y))$ ,  $\overline{co}A(y) := \overline{co}(A(y))$  for every  $y \in Y$ .

**Remark 2.** Claim 7 holds true if  $X$  is reflexive, but  $F : X \rightarrow \mathbb{R} \cup \{+\infty\}$  is weakly lower semicontinuous.

**Corollary 1.** Let  $\varphi_1, \varphi_2 : X \mapsto \mathbb{R}$  be lower semicontinuous convex functionals upper bounded on  $X$ . Then  $\partial\varphi_1 + \partial\varphi_2 : X \rightrightarrows X^*$  is a  $*$ -bounded  $*$ -upper semicontinuous map with compact values in the  $\sigma(X^*, X)$  topology.

*Proof.* The map  $G = \partial\varphi_1 + \partial\varphi_2$  is upper hemicontinuous, since it is the sum of upper hemicontinuous maps. Also,  $\partial\varphi_i = \overline{co}\partial\varphi_i$  ( $i = 1, 2$ ). Now we prove that  $\overline{co}G = G$ . As  $coG = G$ , i.e.,  $\overline{co}G \supset \partial\varphi_1 + \partial\varphi_2 = G$ , it remains to prove the inverse inclusion. Let  $u \in \overline{co}G(y)$ , then there exists a net  $\{u_\alpha\} \in G(y)$  such that  $u_\alpha \rightarrow u$  in  $X^*$ , and  $u_\alpha = u'_\alpha + u''_\alpha$ , where  $u'_\alpha \in \partial\varphi_1(y)$ ,  $u''_\alpha \in \partial\varphi_2(y)$ . Since  $\partial\varphi_1(y)$ ,  $\partial\varphi_2(y)$  are compact sets in  $\sigma(X^*, X)$ -topology, we deduce that  $u = u' + u''$ ,  $u' \in \partial\varphi_1(y)$ ,  $u'' \in \partial\varphi_2(y)$ , i.e.,  $\overline{co}G(y) \subset G(y)$ .

Thus,  $G$  satisfies all conditions of the Castaing Theorem, whence  $*$ -upper semicontinuity of the map  $\partial\varphi_1 + \partial\varphi_2$  follows. The  $*$ -boundedness of the map  $\partial\varphi_1 + \partial\varphi_2$  follows from a similar statement for  $\partial\varphi_1$  and  $\partial\varphi_2$ .

For an arbitrary bounded set  $B$ , images  $\partial\varphi_1(B)$  and  $\partial\varphi_2(B)$  are  $*$ -bounded in  $X^*$ . Then

$$\begin{aligned} \sup_{g \in \partial\varphi_1(B) + \partial\varphi_2(B)} |\langle g, x \rangle_X| &= \sup_{g_1 \in \partial\varphi_1(B)} \sup_{g_2 \in \partial\varphi_2(B)} |\langle g_1 + g_2, x \rangle_X| \leq \\ &\leq \sup_{g_1 \in \partial\varphi_1(B)} |\langle g_1, x \rangle_X| + \sup_{g_2 \in \partial\varphi_2(B)} |\langle g_2, x \rangle_X| < \\ &< +\infty \quad \text{for every } x \in X, \end{aligned}$$

i.e.,  $\partial\varphi_1 + \partial\varphi_2$  is a  $*$ -bounded set in  $X^*$ . □

Let us define

$$\varphi(y) = \varphi_1(y) + \varphi_2(y) - \langle f, y \rangle_X, \quad (16)$$

where  $U$  is a nonempty convex set,  $f \in X^*$ ,  $\varphi_1 : X \mapsto \mathbb{R} \cup \{+\infty\}$  is a convex upper semicontinuous functional on  $X$  ( $\text{int dom } \varphi_1 \neq \emptyset$ ),  $\varphi_2 : X \mapsto \mathbb{R} \cup \{+\infty\}$  is a convex functional on  $U$  and  $\text{dom } \varphi_1 \subset \text{dom } \varphi_2$ .

The following results are true.

**Theorem 5.** Under the above assumptions, the following conditions are equivalent:

- 1)  $x_0 \in \text{int dom } \varphi_1 \cap U$ ,  $\varphi(x_0) = \inf_{x \in U} \varphi(x)$ ;
- 2)  $x_0 \in \text{int dom } \varphi_1 \cap U$ ,  $[\partial\varphi_1(x_0; U), x - x_0]_{++}$

$$+\varphi_2(x) - \varphi_2(x_0) \geq \langle f, x - x_0 \rangle_X \quad \text{for every } x \in U. \quad (17)$$

*Proof.* First we prove that 1)  $\Rightarrow$  2). Let a point  $x_0 \in \text{int dom } \varphi_1 \cap U$  satisfy condition 1). Then for all  $x \in U$  and all  $t \in [0, 1]$  there is

$$\begin{aligned} \varphi(x_0) &= \varphi_1(x_0) + \varphi_2(x_0) - \langle f, x_0 \rangle_X \leq \\ &\leq \varphi_1(x_0 + t(x - x_0)) + \varphi_2(x_0 + t(x - x_0)) - \langle f, x_0 + t(x - x_0) \rangle_X \leq \\ &\leq \varphi_1(x_0 + t(x - x_0)) + t\varphi_2(x) + (1 - t)\varphi_2(x_0) - t\langle f, x - x_0 \rangle_X. \end{aligned}$$

Hence,

$$\frac{\varphi_1(x_0 + t(x - x_0)) - \varphi_1(x_0)}{t} + \varphi_2(x) - \varphi_2(x_0) \geq \langle f, x - x_0 \rangle_X$$

or, passing to a limit as  $t \rightarrow +0$ ,

$$D_+\varphi_1(x_0; x - x_0) + \varphi_2(x) - \varphi_2(x_0) \geq \langle f, x - x_0 \rangle_X.$$

Then, due to relation (12), we arrive at inequality (17).

To prove the inverse implication, assume that inequality (17) holds. By the definition of  $\partial\varphi_1(x_0; U)$ , we obtain

$$\begin{aligned} \varphi_1(x) - \varphi_1(x_0) + \varphi_2(x) - \varphi_2(x_0) &\geq [\partial\varphi_1(x_0; U), x - x_0]_+ + \varphi_2(x) - \varphi_2(x_0) \geq \\ &\geq \langle f, x - x_0 \rangle_X \quad \text{for every } x \in U \end{aligned}$$

i.e.,  $\varphi(x) \geq \varphi(x_0)$ , which is equivalent to 1). This completes the proof of Theorem 5.  $\square$

**Remark 3.** In the literature, inequality (17) is called a variational inequality with a multivalued map. In Banach spaces, such maps are being actively studied.

**Theorem 6.** Let  $X$  be a reflexive space and the functional  $\varphi$  be of the form (16). Let it be coercive and satisfy all conditions of Theorem 5. Let  $U \subset \text{dom } \varphi = X$  be a closed convex set. If the functional  $\varphi_2$  is lower semicontinuous on  $U$ , then variational inequality (17) has at least one solution  $x_0 \in X$ .

*Proof.* The following statement is true. (It represents a generalization of the Weierstrass Theorem onto the case of Frechet spaces.)

**Claim 8.** Let  $X$  be a reflexive Frechet space,  $\varphi : X \rightarrow \mathbb{R} \cup \{+\infty\}$  weakly lower semicontinuous functional,  $B \subset \text{dom } \varphi$  a closed convex set. Moreover, suppose that one of the following conditions holds:

- a) set  $B$  is bounded in  $X$ ;
- b) the functional  $\varphi$  is coercive on  $B$ .

Then functional  $\varphi$  is lower bounded on  $B$  and reaches its exact lower bound  $d$ , and the set

$$K = \{x \in B \mid \varphi(x) = d\}$$

is weakly compact in  $X$ .

*Proof of Claim 8.* Due to Claim 7 and Remark 2, the functional  $\varphi$  is lower bounded. Therefore, there exists a net  $\{x_\alpha\}_\alpha \subset B$  such that

$$\lim_\alpha \varphi(x_\alpha) = d = \inf_{x \in B} \varphi(x) < +\infty.$$

The set  $\{x_\alpha\}_\alpha$  is bounded in  $X$  due to either the boundedness  $B$  or coercivity of  $\varphi$ . Hence, in virtue of the Banach-Alaoglu Theorem, there exists a subnet (which we also denote by  $\{x_\alpha\}_\alpha$ ) such that  $x_\alpha \rightarrow x_0$  in  $\sigma(X; X^*)$ -topology of the space  $X$ , and  $x_0 \in B$ , because the set  $B$  is closed in  $\sigma(X; X^*)$ -topology.

Hence, due to the lower semicontinuity of the functional  $\varphi$  in  $\sigma(X; X^*)$ -topology, we obtain

$$\varphi(x_0) \leq \varliminf_\alpha \varphi(x_\alpha) = \lim_\alpha \varphi(x_\alpha) = d,$$

i.e.,  $x_0 \in K$ .

Finally, let  $\{x_\alpha\}_\alpha \subset K$  be an arbitrary net. By the construction, the set  $K$  is bounded. Consequently, we may assume that  $x_\alpha \rightarrow x_0$  in  $\sigma(X; X^*)$ -topology. So,  $\varphi(x_0) \leq \underline{\lim}_\alpha \varphi(x_\alpha) = d$ , whence  $x_0 \in K$ . Claim 8 is proved.  $\square$

In our case,  $U \subset X = \text{dom}\varphi$  and it satisfies the conditions of Claim 8; therefore, the problem  $\varphi(x) \rightarrow \inf, x \in U$  has a solution  $x_0 \in X$ . In order to complete the proof it remains to use Theorem 5. This completes the proof of Theorem 6.  $\square$

### Acknowledgements

*The authors thank the referee for his/her constructive and useful recommendations.*

### REFERENCES

- [1] B.N. Pshenichniy, *Convex analysis and extremum problems*, Moscow: Nauka (1980), 317 [in Russian].
- [2] J.-P. Aubin, I. Ekeland, *Applied nonlinear analysis*, A Valley-Interscience Publication (1988), 512.
- [3] V.F. Dem'yanov, L.V. Vasiliev, *Non-differentiated optimization*, Moscow: Nauka (1981), 384 [in Russian].
- [4] J.L. Lions, *Quelques methodes de resolution des problemes aux limites non lineaires*, Paris: DUNOD GAUTHIER-VILLARS (1969), 587.
- [5] M.Z. Zgurovsky, V.S. Mel'nik, *Nonlinear analysis and control of infinite-dimensional systems*, Kiev: Naukova dumka (1999), 630 [in Russian].
- [6] J.-P. Aubin, H. Frankowska, *Set-valued analysis*, Birkhauser (1990), 461.
- [7] A.A. Chikrii, *Conflict controlled processes*, Cluver academic publishers (1997), 424.
- [8] A.D. Ioffe, V.M. Tichomirov *The extremal problems theory*, Moscow: Nauka (1974).
- [9] M.Z. Zgurovsky, V.S. Mel'nik, A.N. Novikov, *Applied methods of the analysis and control of nonlinear processes and fields*, Kiev: Naukova dumka (2004), 590 [in Russian].
- [10] W. Rudin, *Functional Analysis*, Mc Graw Hill, New York, 1973, 442.
- [11] V.S. Mel'nik, *About critical points of some classes multivalued maps*, J. Cybernetics and the system analysis, **2** (1997), 87–98.
- [12] M. Reed, B. Simon, *Methods of modern mathematical physics*, ACAD. PRESS INC. **1** (1980), 400.

Pavlo O. Kasyanov  
kasyanov@univ.kiev.ua

Kyiv Taras Shevchenko University  
Mishuga str. 1/4, app. 362, Kyiv, Ukraine

Valery S. Mel'nik  
Institute of Applied System Analysis  
Prospect Peremogy 37, build. 36, app. 203, Kyiv, Ukraine

Anna M. Piccirillo  
annamaria.piccirillo@unina2.it

Second University of Naples  
Aversa, Italy

*Received: November 12, 2005.*

*Revised: December 20, 2007.*

*Accepted: February 20, 2008.*