

PAPER • OPEN ACCESS

Simulation of two-dimensional currents to the depth and suspended sediment concentration in Aceh Besar Waters

To cite this article: I Setiawan *et al* 2019 *IOP Conf. Ser.: Earth Environ. Sci.* **284** 012026

View the [article online](#) for updates and enhancements.



IOP | ebooks™

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of every title for free.

Simulation of two-dimensional currents to the depth and suspended sediment concentration in Aceh Besar Waters

I Setiawan^{1*}, S M Yuni², S Purnawan¹, Y Ilhamsyah^{1,3}, R Wafdan⁴

¹Department of Marine Sciences, Faculty of Marine and Fisheries, Syiah Kuala University, Banda Aceh, 23111, Indonesia

²Department of Mathematics, Faculty of Mathematics and Natural Sciences, Syiah Kuala University, Banda Aceh 23111, Indonesia

³Applied Climatology, Graduate School of Bogor Agricultural University, Bogor Indonesia 16680

⁴Ocean Modeling Laboratory, Syiah Kuala University, Banda Aceh, 23111, Indonesia

Email: ichsansetiawan@unsyiah.ac.id

Abstract. Aceh Besar Waters are geographically located at 5.3° - 5.7°N and 95.1° - 95.6°E which border Indian Ocean, Andaman Sea, and Malacca Strait in the West, North, and East, respectively. The purpose of this study was to simulate the circulation of current and suspended sediment transport driven by southwest and northeast monsoon. Model discretization cover 0.5 x 0.5 arcminutes (925 x 925 m) of spatial resolution with time interval of 9 seconds. The parameters of the model are wind friction $C_d = 1.285 \times 10^{-3}$, bottom friction $r = 0.0025$, lateral eddy viscosity = 2500 m²/s. The primary data of estuary suspended sediment concentrations were collected in the coastal waters of Lhok Nga, Ulee Lheue, Lampulo, Alue Naga and Krueng Raya as input for the suspended sediment distribution model. The results of the study show that during southwest monsoon, the current in the western part of Aceh Besar Waters move northward and during northeast monsoon in the western and eastern part of the region, the current move toward south and west. Thus, the distribution of suspended sediments follows the direction of the current, where sediment concentrations are reduced when they are away from the coast.

1. Introduction

The Aceh Besar Waters is part of Aceh Waters that is geographically located in the northern tip of Sumatera Island at 5.3° - 5.7°N and 95.1° - 95.6°E. It is adjacent to Indian Ocean, Andaman Sea, and Malacca Strait. Study on current circulation have been carried out including circulation in the Indian Ocean [1, 2]; the Malacca Strait [3, 4]; the northern waters of Aceh [5]; and in the Andaman Sea [6].

The study of sediment dynamics has been carried out [7] regarding suspended sediment transport 3D modeling in tidal estuary waters; Sravanthi et al. [8] conducted a numerical application of drifting sediment transport in Central Kerala, West-coast of India. Thus, it is very important to examine the circulation and sediment transport in the coastal waters of Aceh Besar (figure 1). The purpose of this study is to obtain a current-driven southwest and northeast monsoon and suspended sediment transport driven by current circulation which results in a distribution of suspended sediment concentrations.






Figure 1. Bathymetric map (depth is in meters).

2. Material and methods

Hydrodynamics (equations 1-6) and advection-diffusion models (equation 7) are employed to simulate current circulation and suspended sediment transport driven by wind forcing [9, 10, 11] with a discretization of 0.5×0.5 arcminutes (925×925 m) with interval of 9 arcseconds, wind friction $C_d = 1.285 \times 10^{-3}$, bottom friction $r = 0.0025$, lateral eddy viscosity = $2500 \text{ m}^2/\text{s}$. A couple of representative seasonal circulation include northeast and southwest monsoon. Wind data were retrieved from European Centre for Medium-Range Weather Forecasts (ECMWF) during February and August 2016 (figure 2 and figure 3). Data on estuary sediment concentrations were collected in the coastal waters of Lhok Nga, Ulee Lheue, Lampulo, Alue Naga and Krueng Raya as inputs for suspended sediment distribution models. The concentrations of Lhok Nga, Ulee Lheue, Lampulo, Alue Naga and Krueng Raya were 260 mg/l , 680 mg/l , 590 mg/l , 700 mg/l , and 740 mg/l , respectively.

$$\frac{\partial u}{\partial t} + Adv_h(u) - fv = -g \frac{\partial \eta}{\partial x} + \frac{\tau_x^{wind} - \tau_x^{bot}}{\rho_o} + Diff_h(u) \quad (1)$$

$$\frac{\partial v}{\partial t} + Adv_h(v) + fu = -g \frac{\partial \eta}{\partial y} + \frac{\tau_y^{wind} - \tau_y^{bot}}{\rho_o} + Diff_h(v) \quad (2)$$

$$\frac{\partial \eta}{\partial t} + \frac{\partial(uh)}{\partial x} + \frac{\partial(vh)}{\partial y} = 0 \quad (3)$$

$$dv(\psi) = u \frac{\partial \psi}{\partial x} + v \frac{\partial \psi}{\partial y} \quad (4)$$

$$Diff(\psi) = \frac{\partial}{\partial x} \left(A_h \frac{\partial \psi}{\partial x} \right) + \frac{\partial}{\partial y} \left(A_h \frac{\partial \psi}{\partial y} \right) \quad (5)$$

Courant-Friedrich-Lewy (CFL) criterion for stability:

$$\Delta t \leq \frac{\min(\Delta x, \Delta y)}{\sqrt{2gh_{max}}} \quad (6)$$

$$\frac{\partial C}{\partial t} + Adv_h(C) = Diff_h(C) \quad (7)$$

Where t , $u(t, x, y)$, $v(t, x, y)$, ψ are time, magnitudes of ocean currents in the west-east and north-south directions, arbitrary parameters respectively. $Adv_h(u, v)$ is a horizontal component of advection, $Diff_h(u, v)$ is a component of horizontal diffusion. $-fv$ and $+fu$ are Coriolis forces. The gradient pressure force is $-g \partial \eta / \partial x$ and $-g \partial \eta / \partial y$. Whereas the sum of wind friction and bottom friction is $\frac{\tau_x^{wind} - \tau_x^{bottom}}{(\rho_0 h)}$ and $\frac{\tau_y^{wind} - \tau_y^{bottom}}{(\rho_0 h)}$. The depth of seawater is h and the maximum depth is h_{max} , while g is the gravitational force.




Figure 2. Wind vector during Northeast monsoon in the Aceh Besar waters (m/s).




Figure 3. Wind vector during Southwest monsoon in the Aceh Besar waters (m/s).

3. Results and discussion

Current circulation and sediment transport simulations are shown in figure 4 and 5. The ocean current direction on February 2016 (figure 4) follows the effects of wind and Coriolis forces. In the western part of Aceh Besar, the current move northward following topographical condition. Current velocity is high at offshore and decreases when approaching the coastal area. The current velocity in the west coast reached 0.2 m/s while near coast was around 0.1 m/s.

On August 2016 (figure 5) the current in the western part of Aceh Besar moved southward with magnitudes of 0.2 m/s. Meanwhile, the distribution of suspended sediments followed the direction of the current, as the concentration of sediments moved away from the coast, the sediment concentration decreased (figure 4 and 5).

The model in this study was in accordance with the results obtained in previous studies [1, 2, 12]. The influence of the monsoonal circulation was well-described especially in the western part of the Aceh waters.




Figure 4. Average current in vector (m/s) and 1 month suspended sediment concentration in shaded contour (mg/l).




Figure 5. Average current in vector (m/s) and 1 month suspended sediment concentration in shaded contour (mg/l).

4. Conclusion

The simulation of two-dimensional currents to the depth and suspended sediment concentration has been carried out in Aceh Besar Waters. During the southwest monsoon, the current in the western part of Aceh Besar Waters move northward whereas during northeast monsoon in the western and eastern part of the region, the current move toward south and west. The distribution of suspended sediments follows the direction of the current, where sediment concentrations are reduced when they are away from the coast.

Acknowledgments

Authors express gratitude to the Ministry of Research, Technology and Higher Education of Indonesia for financial assistance under ‘*Penelitian Dasar Unggulan Perguruan Tinggi*’, with contract number: 36/UN.11.2/PP/SP3/2018.

References

- [1] Diansky NA, Zalesny VB, Moshonkin SN, and Rusakov AS 2006 High resolution modeling of the monsoon circulation in the Indian Ocean *Oceanology* **46** 608-628 doi: 10.1134/S000143700605002X
- [2] Schott FA, Xie SP, and McCreary, JP 2009 Indian Ocean circulation and climate variability *Reviews of Geophysics* **47** pp 1-46
- [3] Rizal S 2002 Taylor problem-influences on the spatial distribution of real and virtual amphidromes *Continental Shelf Res.* **22** pp 2147-2158
- [4] Rizal S, Setiawan I, Iskandar T, Ilhamsyah Y, Wahid MA, and Musman M 2010 Currents simulation in the Malacca Straits by using three-dimensional numerical model *Sains Malaysiana* **4** pp 519-524
- [5] Setiawan I, Rizal S, Haditiar Y, Ilhamsyah Y, Purnawan S, Irham M, and Yuni SM 2018 Study of Current Circulation in the Northern Waters of Aceh *IOP Conf. Ser.: Earth Environ. Sci.* **176** 012016 doi: 10.1088/1755-1315/176/1/012016
- [6] Rizal S, Damm P, Wahid MA, Sündermann J, Ilhamsyah Y, Iskandar T, and Muhammad 2012 General circulation in the Malacca Strait and Andaman Sea: A numerical model study *American Journal of Environmental Sciences* **8** pp 479-488 doi: 10.3844/ajessp.2012.479.488
- [7] Chen WB, Liu WC, Hsu MH, and Hwang CC 2015 Modeling investigation of suspended sediment transport in a tidal estuary using a three-dimensional model *Applied Mathematical Modelling* **39** pp 2570-2586 doi: 10.1016/j.apm.2014.11.006
- [8] Sravanthi N, Ramakrishnan R, Rajawat AS, and Narayana AC 2015 Application of numerical model in suspended sediment transport studies along the Central Kerala West-coast of India *Aquatic Procedia* **4** pp 109-116 doi: 10.1016/j.aqpro.2015.02.016
- [9] Kämpf J 2009 *Ocean Modelling For Beginner Using Open-Source Software* Springer Verlag, Heidelberg p 175
- [10] Rivera, PC 1997 Hydrodynamics, sediment transport and light extinction off Cape Bolinao, Philippines *Thesis* p 268 Retrieved from : <http://en.scientificcommons.org/39658876>
- [11] Koutitas CG 1988 *Mathematical models in coastal engineering* Pentech Press p 166
- [12] Wyrтки K 1961 *Physical Oceanography of the Southeast Asian waters* Scripps Institution of Oceanography UC San Diego: Scripps Institution of Oceanography p 195 Retrieved from: <http://escholarship.org/uc/item/49n9x3t4>