

Deformed Richardson-Gaudin model

P Kulish¹, A Stolin² and L H Johannesson²

¹St. Petersburg Department of Steklov Mathematical Institute, Russian Academy of Sciences
 Fontanka 27, 191023 St. Petersburg, Russia

²University of Gothenburg, Box 100, SE-405 30 Gothenburg, Sweden

Abstract. The Richardson-Gaudin model describes strong pairing correlations of fermions confined to a finite chain. The integrability of the Hamiltonian allows the algebraic construction of its eigenstates. In this work we show that the quantum group theory provides a possibility to deform the Hamiltonian preserving integrability. More precisely, we use the so-called Jordanian r -matrix to deform the Hamiltonian of the Richardson-Gaudin model. In order to preserve its integrability, we need to insert a special nilpotent term into the auxiliary L -operator which generates integrals of motion of the system. Moreover, the quantum inverse scattering method enables us to construct the exact eigenstates of the deformed Hamiltonian. These states have a highly complex entanglement structure which require further investigation.

The Richardson-Gaudin model [1, 2] is an integrable spin- $\frac{1}{2}$ periodic chain with Hamiltonian

$$H = \sum_{j=1}^N \epsilon_j S_j^z + g \sum_{j,k=1}^N S_j^- S_k^+ \quad (1)$$

where g is a coupling constant and $S_l^\pm = S_l^x \pm iS_l^y$, with N copies of the Lie algebra $su(2)$ generators S_l^α ,

$$[S_l^\alpha, S_{l'}^\beta] = i\epsilon^{\alpha\beta\gamma} S_l^\gamma \delta_{ll'}, \quad \alpha, \beta = x, y, z$$

As shown by Cambiaggio *et al* [3], by introducing the fermion operators c_{lm}^\dagger and c_{lm} related to the $sl(2)$ generators by

$$S_l^z = 1/2 \sum_m c_{lm}^\dagger c_{lm} - 1/2, \quad S_l^+ = \frac{1}{2} \sum_m c_{lm}^\dagger c_{l\bar{m}}^\dagger = (S_l^-)^\dagger$$

the Richardson-Gaudin model in Eq. (1) gets mapped onto the pairing model Hamiltonian

$$H_P = \sum_l \epsilon_l \hat{n}_l + g/2 \sum_{l,l'} A_l^\dagger A_{l'} \quad (2)$$

Here c_{lm}^\dagger (c_{lm}) creates (annihilates) a fermion in the state $|lm\rangle$ (with $|l\bar{m}\rangle$ in the time reversed state of $|lm\rangle$) and

$$n_l = \sum_m c_{lm}^\dagger c_{lm}, \quad A_l^\dagger = (A_l)^\dagger = \sum_m c_{lm}^\dagger c_{l\bar{m}}^\dagger$$



are the corresponding number- and pair-creation operators. The pairing strengths $g_{ll'}$ are here approximated by a single constant g , with ϵ_l the single-particle level corresponding to the m -fold degenerate states $|lm\rangle$.

As it is well-known, the pairing model in Eq. (2) is central in the theory of superconductivity. Richardson's exact solution of the model [1], exploiting its integrability, has been important for applications in mesoscopic and nuclear physics where the small number of fermions prohibits the use of conventional BCS theory [4]. Moreover, its (pseudo)spin representation in the guise of the Richardson-Gaudin model, Eq. (1), provides a striking link between quantum magnetism and pairing phenomena, both central concepts in the physics of quantum matter.

The eigenstates of the Richardson-Gaudin Hamiltonian, eq. (1), can be constructed algebraically using the quantum inverse scattering method (QISM) [5, 6]. The main objects of this method are the classical r -matrix

$$r(\lambda, \mu) = \frac{4}{\lambda - \mu} \sum_{\alpha} S^{\alpha} \otimes S^{\alpha} \Big|_{s=\frac{1}{2}} \simeq \frac{1}{\lambda - \mu} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 2 & 0 \\ 0 & 2 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad (3)$$

where $h(\lambda)$, $X^+(\lambda)$, $X^-(\lambda)$ are the generators of the loop algebra $\mathcal{L}(sl(2))$ whereas the L -matrix is

$$L(\lambda) = \begin{pmatrix} h(\lambda) & 2X^-(\lambda) \\ 2X^+(\lambda) & -h(\lambda) \end{pmatrix}$$

The commutation relations (CR) of loop algebra generators are given in compact matrix form

$$[L_1(\lambda), L_2(\mu)] = -[r_{12}(\lambda, \mu), L_1(\lambda) + L_2(\mu)]$$

where

$$L_1(\lambda) = L(\lambda) \otimes \mathbb{I}, \quad L_2(\mu) = \mathbb{I} \otimes L(\mu)$$

and $r(\lambda, \mu)$ is the 4×4 c -number matrix in Eq. (3). A consequence of this form is the commutativity of transfer matrices,

$$t(\lambda) = \frac{1}{2} \text{tr}_0(L^2(\lambda)) \in \mathcal{L}(sl(2)), \quad [t(\lambda), t(\mu)] = 0 \quad (4)$$

The corresponding mutually commuting operators extracted from the decomposition of $t(\lambda)$ define a Gaudin model [2, 7]. However, to get Richardson Hamiltonian a mild change of the L -operator is necessary

$$L(\lambda) \rightarrow L(\lambda; c) := c h_0 + L(\lambda)$$

where $h_0 = \sigma_0^z$ in auxiliary space \mathbb{C}_0^2 of spin $1/2$. This transformation does not change the CR of matrix elements of this matrix $L(\lambda; c)$ due to the symmetry of the r -matrix (3):

$$[Y \otimes \mathbb{I} + \mathbb{I} \otimes Y, r(\lambda, \mu)] = 0, \quad Y \in sl(2)$$

The resulting transfer matrix obtains some extra terms

$$t(\lambda; c) = \frac{1}{2} \text{tr}_0 (L(\lambda; c))^2 = c^2 \mathbf{1} + c h(\lambda) + h^2(\lambda) + 2 (X^+(\lambda)X^-(\lambda) + X^-(\lambda)X^+(\lambda))$$

Let us consider a spin- $\frac{1}{2}$ representation on auxiliary space $V_0 \simeq \mathbb{C}^2$ and spin ℓ_k representations on quantum spaces $V_k \simeq \mathbb{C}^{\ell_k+1}$ with extra parameters ϵ_k corresponding to site $k = 1, 2, \dots, N$.

The whole space of quantum states is $\mathcal{H} = \otimes_1^N V_k$ and the highest weight vector (highest spin, "ferromagnetic state") $|\Omega_+\rangle$ satisfies

$$X^+(\lambda) |\Omega_+\rangle = 0, \quad h(\lambda) |\Omega_+\rangle = \rho(\lambda) |\Omega_+\rangle \quad (5)$$

where

$$\rho(\lambda) = \sum_{k=1}^N l_k / (\lambda - \epsilon_k)$$

It is useful to introduce notation for global operators of $sl(2)$ -representation $Y_{gl} := \sum_{k=1}^N Y_k$. To find the eigenvectors and spectrum of $t(\lambda)$ on \mathcal{H} one requires that vectors of the form

$$|\mu_1, \dots, \mu_M\rangle = \prod_{j=1}^M X^-(\mu_j) |\Omega_+\rangle$$

are eigenvectors of $t(\lambda)$,

$$t(\lambda) |\{\mu_j\}_{j=1}^M\rangle = \Lambda(\lambda; \{\mu_j\}_{j=1}^M) |\{\mu_j\}_{j=1}^M\rangle$$

provided that the parameters μ_j satisfy the Bethe equations:

$$2c + \sum_{k=1}^N \ell_k / (\mu_i - \epsilon_k) - \sum_{j \neq i}^M 2 / (\mu_i - \mu_j) = 0, \quad i = 1, \dots, M \quad (6)$$

The realization of the loop algebra generators on the space \mathcal{H} takes the form

$$h(\lambda) = \sum_{k=1}^N \frac{h_k}{\lambda - \epsilon_k}, \quad X^-(\lambda) = \sum_{k=1}^N \frac{X_k^-}{\lambda - \epsilon_k}, \quad X^+(\lambda) = \sum_{k=1}^N \frac{X_k^+}{\lambda - \epsilon_k} \quad (7)$$

The coupling constant g of (1) is connected with parameter $c = 1/g$ while the Hamiltonian (1) is obtained as operator coefficient of term $1/\lambda^2$ in the expansion of $t(\lambda; c)$ at $\lambda \rightarrow \infty$.

The quantum group theory provides a possibility to deform a Hamiltonian preserving integrability [8, 9]. Specifically, we use the so-called Jordanian r -matrix to quantum deform the Hamiltonian of Richardson-Gaudin model (1). We add to $sl(2)$ symmetric r -matrix (3) the Jordanian part

$$r^J(\lambda, \mu) = \frac{C_2^\otimes}{\lambda - \mu} + \xi (h \otimes X^+ - X^+ \otimes h)$$

with Casimir element C_2^\otimes in the tensor product of two copies of $sl(2)$,

$$C_2^\otimes = h \otimes h + 2(X^+ \otimes X^- + X^- \otimes X^+)$$

After Jordanian twist the r -matrix (14) is commuting with the generator X_0^+ only

$$\left[X_0^+ \otimes \mathbb{I} + \mathbb{I} \otimes X_0^+, r^{(J)}(\lambda, \mu) \right] = 0$$

Hence, one can add the term $cX_0^+ + L(\lambda, \xi)$ to the L -operator. This yields the twisted transfer-matrix

$$t^{(J)}(\lambda) = \frac{1}{2} \text{tr}_0(cX_0^+ + L(\lambda, \xi))^2 = cX^+(\lambda) + h(\lambda)^2 - 2h'(\lambda) + 2(2X^-(\lambda) + \xi)X^+(\lambda) \quad (8)$$

The corresponding commutation relations between the generators of the twisted loop algebra are explicitly given by

$$\begin{aligned} [h(\lambda), h(\mu)] &= 2\xi (X^+(\lambda) - X^+(\mu)), & [X^-(\lambda), X^-(\mu)] &= -\xi (X^-(\lambda) - X^-(\mu)) \\ [X^+(\lambda), X^-(\mu)] &= -\frac{h(\lambda) - h(\mu)}{\lambda - \mu} + \xi X^+(\lambda), & [X^+(\lambda), X^+(\mu)] &= 0 \\ [h(\lambda), X^-(\mu)] &= 2\frac{X^-(\lambda) - X^-(\mu)}{\lambda - \mu} + \xi h(\mu), & [h(\lambda), X^+(\mu)] &= -2\frac{X^+(\lambda) - X^+(\mu)}{\lambda - \mu} \end{aligned} \quad (9)$$

The realization of the Jordanian twisted loop algebra $\mathcal{L}_J(sl(2))$ with CR (9) is given similar to (7) with extra terms proportional to the deformation parameter ξ

$$h(\lambda) = \sum_{k=1}^N \left(\frac{h_k}{\lambda - \epsilon_k} + \xi X_k^+ \right), \quad X^-(\lambda) = \sum_{k=1}^N \left(\frac{X_k^-}{\lambda - \epsilon_k} - \frac{\xi}{2} h_k \right), \quad X^+(\lambda) = \sum_{k=1}^N \frac{X_k^+}{\lambda - \epsilon_k} \quad (10)$$

To construct eigenstates for the twisted model one has to use operators of the form [9, 10]

$$B_M(\mu_1, \dots, \mu_M) = X^-(\mu_1) (X^-(\mu_2) + \xi) \dots (X^-(\mu_M) + \xi(M-1))$$

acting by these operators on the ferromagnetic state $|\Omega_+\rangle$.

The deformed Richardson-Gaudin model Hamiltonian can now be extracted from the transfer-matrix $t^{(J)}(\lambda)$ as the operator coefficient in its expansion $\lambda \rightarrow \infty$.

According to (4) and (8) one can also extract quantum integrals of motion J_k using the realization (10). It would yield rather cumbersome expressions for J_k :

$$t^{(J)}(\lambda) = J_0 + \frac{1}{\lambda} J_1 + \frac{1}{\lambda^2} J_2 + \dots$$

The corresponding quantum deformed Hamiltonian reads

$$H \simeq J_2 = c \sum_{j=1}^N \epsilon_j X_j^+ + 2\xi \left\{ \left(\sum_{j=1}^N \epsilon_j h_j \right) X_{gl}^+ - h_{gl} \sum_{j=1}^N \epsilon_j X_j^+ \right\} + (h_{gl}^2 + 2h_{gl} + 4X_{gl}^- X_{gl}^+)$$

It is instructive to write down a simplified case without the Jordanian twist: $\xi = 0$. One thus obtains

$$J_0 = 0, \quad J_1 = X_{gl}^+, \quad J_2 \simeq \sum_{k=1}^N \epsilon_k X_k^+ + g/2 (h_{gl}^2 + 2h_{gl} + 4X_{gl}^- X_{gl}^+)$$

The case $\xi = 0$ can also be obtained by taken off from the inhomogeneous XXX spin chain. The model can be described by a 2×2 monodromy matrix [5]

$$T(\lambda) = \begin{pmatrix} A(\lambda) & B(\lambda) \\ C(\lambda) & D(\lambda) \end{pmatrix}$$

and entries of this matrix satisfy quadratic relations

$$R(\lambda, \mu)T(\lambda) \otimes T(\mu) = (I \otimes T(\mu)) (T(\lambda) \otimes I) R(\lambda, \mu) \quad (11)$$

If we multiply $T(\lambda)$ by a constant 2×2 matrix $M(\varepsilon)$ the resulting matrix $\tilde{T}(\lambda) = M(\varepsilon) \cdot T(\lambda)$ will satisfy the same relation (11). Choosing a triangular matrix $M(\varepsilon) = \begin{pmatrix} 1 & \varepsilon \\ 0 & 1 \end{pmatrix}$ the entries of monodromy matrices become simply related:

$$\tilde{A} = A + \varepsilon C, \quad \tilde{B} = B + \varepsilon D, \quad \tilde{C} = C, \quad \tilde{D} = D.$$

This choice of $M(\varepsilon)$ (of the same type as considered in [11]) permits us to use the same reference state $|\Omega_+\rangle \in \mathcal{H}$ (5) and \tilde{B} as a creation operator of the algebraic Bethe ansatz [5].

Bethe states are given by the same action of product operators $\tilde{B}(\mu_j) = B(\mu_j) + \varepsilon D(\mu_j)$ although operators $B(\mu_j)$ do not commute with $D(\mu_j)$:

$$D(\lambda)B(\mu) = \alpha(\lambda, \mu)B(\mu)D(\lambda) + \beta(\lambda, \mu)B(\lambda)D(\mu)$$

where

$$\alpha(\lambda, \mu) = (\lambda - \mu + \eta)/(\lambda - \mu), \quad \beta(\lambda, \mu) = -\eta/(\lambda - \mu)$$

For a 3 magnon state one gets due to B-D ordering

$$\begin{aligned} \prod_{j=1}^3 \tilde{B}(\mu_j) &= \prod_{j=1}^3 B(\mu_j) + \varepsilon \sum_{s=1}^3 \alpha(\mu_k, \mu_s) \alpha(\mu_s, \mu_l) B(\mu_k) B(\mu_l) D(\mu_s) \\ &\quad + \varepsilon^2 \sum_{s=1}^3 \alpha(\mu_k, \mu_s) \alpha(\mu_l, \mu_s) B(\mu_s) D(\mu_k) D(\mu_l) + \varepsilon^3 \prod_{j=1}^3 D(\mu_j) \end{aligned}$$

Similar formula is valid for M -magnon state. Hence, acting on ferromagnet state $|\Omega_+\rangle$, we obtain filtration of states with eigenvalues of S^z : $\frac{N}{2}, \frac{N}{2} - 1, \frac{N}{2} - 2, \frac{N}{2} - 3$.

More complicated deformations of the Richardson-Gaudin model can be obtained using r -matrices related to the higher rank Lie algebras [12]. The structure of the eigenstates of the transfer matrix and their entanglement properties [13] are under investigation.

Acknowledgments

This work was supported by RFBR grants 11-01-00570-a, 12-01-00207-a, 13-01-12405-ofi-M2 (P.K.), and by STINT grant IG2011-2028 (A.S. and H.J.). We would like to thank E. Damaskinsky for useful discussion.

References

- [1] Richardson R W 1963 *J. Math. Phys.* **6** 1034
- [2] Gaudin M 1983 *La fonction d'onde de Bethe* (Paris: Masson) chapter 13
- [3] Cambiaggio M C, Rivas A M F and Saraceno M 1997 *Nucl. Phys. A* **624** 157
- [4] Dukelsky J, Pittel S and Sierra G 2004 *Rev. Mod. Phys.* **76** 643
- [5] Faddeev L D 1998 How algebraic Bethe ansatz works for integrable models *Quantum symmetries, Proceedings of the Les Houches summer school, session LXIV* eds A Connes, K Gawedzki and J Zinn-Justi (North-Holland) p 149
- [6] Kulish P P and Sklyanin E K 1982 Quantum spectral transform method. Recent developments *Lecture Notes in Phys.* **151** (Springer-Verlag) p 61
- [7] Sklyanin E K 1999 *Lett. Math. Phys.* **47** 275
- [8] Kulish P P and Stolin A A 1997 *Czech. J. Phys.* **12** 207
- [9] Kulish P P 2002 *Twisted $sl(2)$ Gaudin model* Preprint PDMI 08/2002
- [10] Antonio N C and Manojlovic N 2005 *J. Math. Phys.* **46** 102701
- [11] Mukhin E, Tarasov V and Varchenko A 2010 *Contemp. Math.* **506** 187
- [12] Stolin A A 1991 *Math. Scand.* **69** 57
- [13] Dunning C, Links J and Zhou H-Q 2005 *Phys. Rev. Lett.* **94** 227002