

Jet Nuclear Modification Factor from the AdS/CFT Correspondence

R Morad and W A Horowitz

Department of Physics, University of Cape Town, Private Bag X3, Rondebosch 7701, SA

E-mail: razieh.morad@uct.ac.za and wa.horowitz@uct.ac.za

Abstract.

We define a novel definition of the holographic light hadron jet by a separation of scales from plasma to jet, which leads to the re-emergence of the late-time Bragg peak in the instantaneous jet energy loss rate. We use the holographic energy loss of light quark and our new jet prescription in AdS/CFT to calculate the nuclear modification factor of jet for a brick of plasma (both static and expanding plasma) and compare the AdS/CFT results with the experimental data for most central Pb-Pb collision at LHC at 2.76 TeV center-of-mass energy. Defining a renormalized AdS/CFT jets that we argue better reflect QCD physics, we find a surprisingly good agreement between our toy model which is the first fully strongly coupled calculations, and preliminary jet suppression data from heavy ion collisions at LHC.

1. Introduction

The spectacular measurements from the Relativistic Heavy Ion Collider (RHIC) and the Large Hadron Collider (LHC) provide compelling evidence that the matter produced in heavy ion collision is a deconfined state of QCD, Quark-Gluon Plasma (QGP) [1, 2, 3, 4], at temperatures above ~ 160 MeV which appears to be nearly perfect, with an extremely low viscosity-to-entropy ratio $\eta/s \sim 1/4\pi$ [5, 6]. While lattice QCD is the proper tool for understanding the static equilibrium thermodynamics of such strongly coupled plasma, it does not allow us to calculate its dynamics evolution on heavy-ion collision.

Recently, a novel tool called “the AdS/CFT correspondence” [7, 8, 9, 10, 11] provide valuable insight into the strongly coupled plasma. This conjecture state that the $\mathcal{N} = 4$ SYM theory in the large N_c and large 't Hooft coupling is dual to classical supergravity on ten-dimensional $AdS_5 \times S^5$ geometry [7]. In order to study the theory at finite temperature, one can add black hole (BH) to the geometry [8] which yields to the AdS-Sch metric. Fundamental quarks are described by open strings moving in the 10d geometry. In the large λ limit, the quantum fluctuation of string world sheet are suppressed and the dynamics of string are described by the classical string theory.

Jets are produced within the expanding fireball and probe the QGP. Analyses the energy loss of these energetic partons as they travel throw QGP may reveal extremely valuable information about the dynamics of the plasma and exhibit distinctive properties such as jet-quenching which can clearly be observed at RHIC [1, 2, 3, 4] and more recently LHC [12, 13, 14].

In this paper, we study the light quark jet energy loss in both gravitational background dual to static and expanding plasma. We propose a new prescription of jets in string theory based



on the separation of hard and soft sectors. We demonstrate that the light quark jet energy loss reveal the “Bragg peak” at late times. Then we compute the nuclear modification factor of jets R_{AA}^{jet} , renormalise the quantity, and compare the results with the preliminary CMS data [15].

2. Light quark jet energy loss

According to the AdS/CFT correspondence [10], the $\mathcal{N} = 4$ SYM theory at finite temperature is dual to a $10d$ black hole geometry with the AdS-Schwarzschild (AdS-Sch) metric as follows,

$$ds^2 = \frac{L^2}{u^2} \left[-f(u) dt^2 + d\mathbf{x}^2 + \frac{du^2}{f(u)} \right], \quad (1)$$

where $f(u) \equiv 1 - (u/u_h)^4$ is the blackening factor and L is the AdS curvature radius. Four dimensional Minkowski coordinates are denoted by x_μ and the coordinate u is an inverse radial coordinate. So, the boundary of the AdS-Sch spacetime is at $u = 0$ and the event horizon is located at $u = u_h$. The temperature of the equilibrium SYM plasma relates to the event horizon as $T \equiv \frac{1}{(\pi u_h)}$. World sheet coordinates are as σ^a where $\tau \equiv \sigma^0$ is denoted as the timelike world sheet coordinate, while the spatial coordinate is $\sigma \equiv \sigma^1$.

Fundamental representation quarks added to the $\mathcal{N} = 4$ SYM theory are dual to open strings moving in the $10d$ geometry. Addition of a $\mathcal{N} = 2$ hypermultiplet to the $\mathcal{N} = 4$ SYM theory is performed by adding D7 branes to the $10d$ geometry [16]. These branes extend along the radial coordinate from the boundary at $u = 0$ down to maximal coordinate at $u = u_m$ as well as fill the whole 4d Minkowski space. Also, they wrap on S^3 from the S^5 sphere. The bare mass M of quark is proportional to $1/u_m$ [17], so for massless quarks the D7 brane should fill the whole radial direction. Open strings that are attached to the D7 brane are dual to the quark-anti quark pairs on the field theory side. In the $5d$ geometry these strings can fall unimpeded toward the event horizon until their end points reach the radial coordinate u_m where the D7 brane ends. Since for sufficiently light or massless quarks $u_m > u_h$, open string end points can fall into the horizon.

We are interested in studying the back-to-back jets so we consider the configurations in which the two endpoints of string move away from each other as the total spatial momentum of the string vanishes. By choosing the appropriate frame, one half of the string has a large spatial momentum in x direction, and the other half of the string carries a large negative spatial momentum. We will limit our attention to strings which move in one direction in the R^3 space, x direction, so the embedding function of string $X^\mu(\tau, \sigma)$ will be a map to $(t(\tau, \sigma), x(\tau, \sigma), u(\tau, \sigma))$. So, the profile of an open string which is created at a point in space at time $t = t_c$ is given by

$$t(0, \sigma) = t_c, \quad x(0, \sigma) = 0, \quad u(0, \sigma) = u_c. \quad (2)$$

By these conditions, the string created at time t_c and by time evolution, the string evolves from a point into an extended object and the string endpoints fall toward the horizon.

The dynamics of string is governed by the Polyakov action

$$S_P = -\frac{T_0}{2} \int d^2\sigma \sqrt{-\eta} \eta^{ab} \partial_a X^\mu \partial_b X^\nu G_{\mu\nu}. \quad (3)$$

Variation of the Polyakov action with respect to the embedding functions X^μ lead to the equation of motion as

$$\partial_a \left[\sqrt{-\eta} \Pi_\mu^a \right] = \frac{1}{2} \sqrt{-\eta} \eta^{ab} \frac{\partial G_{\nu\rho}}{\partial X^\mu} \partial_a X^\nu \partial_b X^\rho, \quad (4)$$

where Π_μ^a are the canonical momentum densities associated to the string obtained from varying the action with respect to the derivatives of the embedding functions.

In order to study jets in AdS/CFT we need to define the proper objects in the dual string theory that corresponds to a jet, a slippery object even in field theory; jets are truly only defined by the algorithm used to measure them. Presumably the ideal way to compute jet observables in the dual theory is to compute the energy momentum tensor associated with a high-momentum probe and “run” a jet finding algorithm on the result. However, one can define a jet prescription in the AdS/CFT and calculate the rate of energy loss from the string profile itself.

The authors of [18] are motivated by the localization of the baryon density on the boundary which is of scale of order $\Delta x \sim 1/\pi T$ and defined jet as a part of string which is in the Δx spatial distance from the endpoint. We called this as the “ Δx – prescription” of jet [18].




Figure 1. Illustration of the Δx and Δu prescriptions of a jet in the string theory; see text for details.

In this paper, motivated by the separation of energy scales in, e.g., thermal field theory, we propose rather a Δu prescription which we believe will ultimately provide a closer approximation to the result of a more complete calculation, figure 1. Since the radial coordinate in the string theory sets an energy scale in the field theory, in our Δu prescription the portion of the string above some cutoff $u = u^*$ in the radial direction is considered part of the jet; the portion of the string below the cutoff is considered part of the thermalized medium. By choosing any value of u above the black hole horizon as the cutoff, we regain the natural result that a jet that is thermalized no longer has detectable energy or momentum.

We evaluate the energy loss rate of jet in the radial $\sigma = u$ parametrization for both prescriptions of jet and plot in figure 2(a,b). In order to define jet using the Δx – prescription, we choose $\sigma_\kappa(t)$ as $\Delta x = 0.3/\pi T$. Our result on the energy loss rate of light quark using the Δu – prescription shows a Bragg peak at late times which means the explosive transfer of quark energy to the plasma at late times and is consistence with the previous works [18].

Since the quark-gluon plasma produced at ultra-relativistic heavy ion collision is an expanding and cooling medium, we study the light quark jet energy loss in a time dependent gravity dual to the boost invariant flow [19]. This geometry is similar to the static black hole geometry, but the location of the horizon moves in the bulk as $\tau^{1/3}$ where τ is the proper time. Also the temperature of plasma cools as $T(\tau) \tau^{-1/3}$.

We calculate the energy loss rate of light quark using both Δx – prescription and Δu – prescription of jet. We consider the initial temperature of plasma in JP metric T_c the same




Figure 2. The instantaneous energy loss of a light quark jet as a function of time in the AdS-Sch (a,b) and JP metrics (c,d) in the Δx prescription and Δu prescription. The normalization constant $E_q = 100$ GeV is the initial energy of the jet, which has a virtuality of 175 GeV^2 , and $T = 350$ MeV is the temperature of the plasma.

as the temperature of plasma in AdS-Sch metric. The result is shown in figure 2(c,d) which demonstrates that the behavior of light quark energy loss in the JP metric is same as the AdS-Sch metric, but the distance that quark traveled before thermalizing increases.

3. Nuclear Modification Factor of Jet

We calculate an approximation of the nuclear modification factor R_{AA} for jet using our energy loss model of jet. We consider the contributions of both quark and gluon jets. We assume that the produced parton with initial energy p_T^i loses a fraction of its energy ϵ with probability $P(\epsilon|p_T^i, L, T)$ as the final energy of parton is given by $p_T^f = (1 - \epsilon)p_T^i$. Also, we suppose that the AdS energy loss is approximately independent of the initial energy [20], and the gluons lose their energy by the factor of 2 in the large N_c limit respect to the quarks.

We suppose that the production spectrum can be approximately by a power law [20], with slowly varying with respect to p_T , then we may find a simple equation for the jet nuclear modification factor as follows,

$$R_{AA}^{R \rightarrow jet}(p_T) = \left\langle \int d\epsilon P(\epsilon|p_T, L, T) \left(1 - \epsilon^R\right)^{n_R(p_T)-1} \right\rangle. \quad (5)$$

For an absolutely uniform nucleus that is a 1D line, the geometric average is carried out as an integral over a line of production points with a parton that propagates through the line. In this case, $R_{AA}^{R \rightarrow jet}(p_T)$ is obtained from the below line integral [20]

$$R_{AA}^{R \rightarrow jet}(p_T) = \int_0^{L_{max}} \frac{dl}{L_{max}} \left(1 - \epsilon^R(p_T, l, T)\right)^{n_R(p_T)-1}. \quad (6)$$

We calculate R_{AA} of jet by using our results of jet energy loss in both AdS-Sch and JP metric. Our results are shown in figure 3 (a). The purple curve shows the R_{AA} obtained from the AdS-Sch metric, while the blue curve is the R_{AA} obtained from the JP metric. The AdS/CFT results of jet energy loss show an over suppression of jets in both static and expanding plasma.

The point-like initial condition falling string that we consider here is dual to creation of a pair of quark-antiquark which fly away each other in the strongly coupled plasma, interact and lose their energy. We expect that jets produced in the pp collision do not lose their energy. So, we consider the falling string with the same initial conditions in AdS_5 metric. Our results show that the string falls in the empty AdS_5 . So, jet loses its energy in the vacuum! We calculate the R_{AA} for a falling string in AdS_5 metric and show in figure 3 (a) in red curve.




Figure 3. (a) Jet R_{AA} as a function of p_T in the most central Pb-Pb collision obtained via AdS/CFT in AdS_5 (red), JP (blue) and AdS-Sch (purple) metrics. (b) AdS/CFT jet R_{AA} as a function of p_T compared with the preliminary CMS data in different effective cone sizes for anti- k_T jets using the Bayesian unfolding method for most central Pb-Pb collision at the LHC with $\sqrt{s} = 2.76$ TeV per nucleon [15].

Since the definition of R_{AA} measured the difference between the medium and vacuum effects on jet, we define a renormalized R_{AA} in AdS/CFT as

$$R_{AA}^{\text{renorm}} = \frac{R_{AA}^{\text{medium}}}{R_{AA}^{\text{AdS}_5}} \quad (7)$$

We plot the renormalized R_{AA}^{renorm} for jets in both AdS-Sch and JP metric in figure 3 (b) and compare with the CMS preliminary data for the most central Pb-Pb collision at $\sqrt{s_{NN}} = 2.76$ GeV [15]. A suppression factor of 0.5 for high p_T jets is observed in central Pb-Pb collision in comparison to the pp collisions. Our results also show surprisingly agreement with CMS preliminary results on jet R_{AA} .

4. Conclusions

In this paper we have purposed a novel prescription of jet in the context of string theory and AdS/CFT correspondence. We have defined jet as a part of a falling string which lies above the radial coordinate u^* near the event horizon. Our motivation is defining the jet as hard partons. The instantaneous energy loss of light quark is identified with the energy flux from the point at $u(\tau, \sigma_\kappa) = u^*$. We have shown that using the Δu – prescription of jet, the light quark energy loss exhibit the Bragg peak again at late times in both static and expanding plasmas. This late time behavior of jet energy lost implies that after traveling substantial distances through the plasma, the thermalization of light quark ends with a large amount of energy transferring to the plasma which is similar to the energy loss rate of a fast charge particle moving through ordinary matter.

We considered a brick of plasma and calculated the nuclear modification factor of jet in both AdS-Sch and JP metric. We assumed that the temperature of plasma around 350 MeV at AdS-Sch metric and at initial time in JP metric. Our results show an aver suppression of jet of order of ten respect to the data. We investigated that it is because of the falling string setup at AdS space. In fact, R_{AA} of jet using the falling string in AdS_5 which is dual to jets in vacuum is not one, even though it is less than one. We introduced a renormalized R_{AA}^{renorm} by dividing the




Figure 4. The maximum stopping distance of jet versus the virtuality of quark Q^2 . Our result show a nontrivial dependency of the thermalization time to the initial condition of the string.

R_{AA} in the medium to the R_{AA} in the vacuum. Surprisingly, our ratio shows good agreement with the experimental data on the R_{AA}^{jet} of most central Pb-Pb collision at LHC figure 3 (b).

On the other hand, the light quark energy loss is highly depends on the initial conditions of falling string figure 4. The only way to determine the energy loss of a jet precisely in strongly coupled regime is solving the gravitational bulk-to-boundary problem. One can solve Einstein's equations for the perturbation in the $5d$ geometry due to the presence of the string and according to the bulk to boundary map interpret the near boundary behaviour of the metric perturbation as the perturbation in the SYM energy-momentum tensor by the presence of jet which will be left for the future work.

References

- [1] Adams J *et al.* (STAR Collaboration) 2005 *Nucl.Phys.* **A757** 102–183 (*Preprint nucl-ex/0501009*)
- [2] Adcox K *et al.* (PHENIX Collaboration) 2005 *Nucl.Phys.* **A757** 184–283 (*Preprint nucl-ex/0410003*)
- [3] Arsene I *et al.* (BRAHMS Collaboration) 2005 *Nucl.Phys.* **A757** 1–27 (*Preprint nucl-ex/0410020*)
- [4] Back B, Baker M, Ballintijn M, Barton D, Becker B *et al.* 2005 *Nucl.Phys.* **A757** 28–101 (*Preprint nucl-ex/0410022*)
- [5] Policastro G, Son D T and Starinets A O 2001 *Phys.Rev.Lett.* **87** 081601 (*Preprint hep-th/0104066*)
- [6] Kovtun P, Son D T and Starinets A O 2005 *Phys.Rev.Lett.* **94** 111601 (*Preprint hep-th/0405231*)
- [7] Maldacena J M 1999 *Int.J.Theor.Phys.* **38** 1113–1133 (*Preprint hep-th/9711200*)
- [8] Witten E 1998 *Adv.Theor.Math.Phys.* **2** 253–291 (*Preprint hep-th/9802150*)
- [9] Gubser S, Klebanov I R and Polyakov A M 1998 *Phys.Lett.* **B428** 105–114 (*Preprint hep-th/9802109*)
- [10] Aharony O, Gubser S S, Maldacena J M, Ooguri H and Oz Y 2000 *Phys.Rept.* **323** 183–386 (*Preprint hep-th/9905111*)
- [11] Casalderey-Solana J, Liu H, Mateos D, Rajagopal K and Wiedemann U A 2011 (*Preprint 1101.0618*)
- [12] Yin Z B (ALICE Collaboration) 2013 *Acta Phys.Polon.Supp.* **6** 479–484
- [13] Aad G *et al.* (ATLAS Collaboration) 2010 *Phys.Rev.Lett.* **105** 252303 (*Preprint 1011.6182*)
- [14] Chatrchyan S *et al.* (CMS Collaboration) 2011 *Phys.Rev.* **C84** 024906 (*Preprint 1102.1957*)
- [15] Collaboration C (CMS Collaboration) 2012
- [16] Karch A and Katz E 2002 *JHEP* **0206** 043 (*Preprint hep-th/0205236*)
- [17] Herzog C, Karch A, Kovtun P, Kozcaz C and Yaffe L 2006 *JHEP* **0607** 013 (*Preprint hep-th/0605158*)
- [18] Chesler P M, Jensen K, Karch A and Yaffe L G 2009 *Phys.Rev.* **D79** 125015 (*Preprint 0810.1985*)
- [19] Janik R A and Peschanski R B 2006 *Phys.Rev.* **D73** 045013 (*Preprint hep-th/0512162*)
- [20] Horowitz W A 2010 (*Preprint 1011.4316*)