

The three-center parameterized continuum wave function and Dyson orbitals for the determination of the triply differential cross section of the simple ionization of CO_2 by electron impact

O. Alwan¹, O. Chuluunbaatar^{2,3}, X. Assfeld⁴, and B.B. Joulakian¹

¹ Université de Lorraine, SRSMC (UMR CNRS 7565), 1 bld Arago, bat. ICPM 57078 Metz Cedex 3, France

² Joint Institute for Nuclear Research, Dubna, Moscow Region 141980, Russia

³ National University of Mongolia, Ulaanbaatar, Mongolia

⁴ Université de Lorraine, SRSMC (UMR CNRS 7565), 54506 Vandoeuvre-lès-Nancy Cedex, France

Synopsis (e,2e) ionization of the linear three center CO_2 molecules by fast electron impact.

The triple differential cross section of the simple (e, 2e) ionization of CO_2 is determined [1]. Its variation with the direction of the ejected electron is studied and compared to existing experimental [2] and a theoretical [3] results. The first-order perturbative Born procedure is applied where three-center Dyson type $1\pi_g$ orbital are employed for the bound electron. This orbital is obtained from coupled cluster results [4] by calculating the overlap between the N state of the target and the (N-1) state of the singly ionized ion [5] and constructed by linear combinations of Gaussian type orbitals. The ejected electron is described by a three-center continuum (ThCC) solution of the Schrödinger equation for a specific wave vector k satisfying the correct asymptotic boundary condition. This model given in the following form

$$\chi(\mathbf{k}, \mathbf{r}, \boldsymbol{\rho}) = \frac{\exp(i\mathbf{k}\mathbf{r})}{(2\pi)^{3/2}} M_{a1} F_1(i\alpha_a, 1, -i[kr_a + \mathbf{k}\mathbf{r}_a]) \times M_{b1} F_1(i\alpha_b, 1, -i[kr_b + \mathbf{k}\mathbf{r}_b]) M_{c1} F_1(i\alpha_c, 1, -i[kr_c + \mathbf{k}\mathbf{r}_c]), \quad (1)$$

is an extension of the two-center continuum model developed in the past [6, 7], and lately applied to the ionization of CO_2 [3]. Here

$$M_j = \exp\left(-\pi \frac{\alpha_j}{2}\right) \Gamma(1 - i\alpha_j), \quad j = a, c, b. \quad (2)$$

and $\alpha_j = -Z_j/k$ is the Sommerfeld parameter. Empirical values for the screening of the three nuclei of the target and for the Sommerfeld parameters of the three-center Coulomb continuum function are introduced.




Figure 1. The variation, in polar representation, of the TDCS in terms of the ejection angle θ_e of the ionization of the $1\pi_g$ level of CO_2 for the empirical parameter $Z_a = 0.3$ for the final state. The energy of the scattered electron $E_s = 500\text{eV}$, detected at an angle $\theta_s = -6^\circ$. The energy of the ejected electron $E_e = 37\text{eV}$.

References

- [1] O Alwan, O chuluunbaatar, X. Assfeld, A Naja and B.B Joulakian, 2014 *J. Phys. B: At. Mol. and Opt. Phys.* **47** 225201
- [2] Lahmam-Bennani A, Staicu Casagrande E M and Naja A. 2009 *J. Phys. B: At. Mol. and Opt. Phys.* **42** 235205
- [3] Chuluunbaatar O and Joulakian B B. 2010 *J. Phys. B: At. Mol. and Opt. Phys.* **43** 155201
- [4] Oana C M and Krylov A I 2007 *J. Chem. Phys.* **127** 234106
- [5] Kaplan I G, Barbiellini B and Bansil A. 2003 *Phys. Rev. B* **68** 235104
- [6] Joulakian B, Hanssen J, Rivarola R and Motassim A. 1996 *Phys. Rev. A* **54** 1473
- [7] Chuluunbaatar O, Joulakian B B, Tsookhuu Kh and Vinitisky S I. 2004 *J. Phys. B: At. Mol. and Opt. Phys.* **37** 2607

¹E-mail: osman.alwan@univ-lorraine.fr