

On existence of an x -integral for a semi-discrete chain of hyperbolic type

K Zheltukhin¹ and N Zheltukhina²

¹ Department of Mathematics, Middle East Technical University, Ankara, Turkey

² Department of Mathematics, Bilkent University, Ankara, Turkey

E-mail: zheltukh@metu.edu.tr

Abstract. A class of semi-discrete chains of the form $t_{1x} = f(x, t, t_1, t_x)$ is considered. For the given chains easily verifiable conditions for existence of x -integral of minimal order 4 are obtained.

1. Introduction

In the present paper we consider the integrable differential-difference chains of hyperbolic type

$$t_{1x} = f(x, t, t_1, t_x), \quad (1)$$

where the function $t(n, x)$ depends on discrete variable n and continuous variable x . We use the following notations $t_x = \frac{\partial}{\partial x} t$ and $t_1 = t(n+1, x)$. It is also convenient to denote $t_{[k]} = \frac{\partial^k}{\partial x^k} t$, $k \in \mathbb{N}$ and $t_m = t(n+m, x)$, $m \in \mathbb{Z}$.

The integrability of the chain (1) is understood as Darboux integrability that is existence of so called x - and n -integrals [1, 4]. Let us give the necessary definitions.

Definition 1 Function $F(x, t, t_1, \dots, t_k)$ is called an x -integral of the equation (1) if

$$D_x F(x, t, t_1, \dots, t_k) = 0$$

for all solutions of (1). The operator D_x is the total derivative with respect to x .

Definition 2 Function $G(x, t, t_x, \dots, t_{[m]})$ is called an n -integral of the equation (1) if

$$DG(x, t, t_x, \dots, t_{[m]}) = G(x, t, t_x, \dots, t_{[m]})$$

for all solutions of (1). The operator D is a shift operator.

To show the existence of x - and n -integrals we can use the notion of characteristic ring. The notion of characteristic ring was introduced by Shabat to study hyperbolic systems of exponential type (see [11]). This approach turns out to be very convenient to study and classify the integrable equations of hyperbolic type (see [12] and references there in).

For difference and differential-difference chains the notion of characteristic ring was developed by Habibullin (see [3]-[8]). In particular, in [4] the following theorem was proved



Theorem 3 (see [4]). *A chain (1) admits a non-trivial x -integral if and only if its characteristic x -ring is of finite dimension.*

A chain (1) admits a non-trivial n -integral if and only if its characteristic n -ring is of finite dimension.

For known examples of integrable chains the dimension of the characteristic ring is small. The differential-difference chains with three dimensional characteristic x -ring were considered in [6]. We consider chains with four dimensional characteristic x -ring, such chains admit x -integral of minimal order four. That is we obtain necessary and sufficient conditions for a chain to have a four dimensional characteristic x -ring. This conditions can be easily checked by direct calculations.

Note that if a chain (1) admits a nontrivial x -integral $F(x, t, t_1, \dots, t_k)$ and a non trivial n -integral $G(x, t, t_x, \dots, t_{[m]})$ its solutions satisfy two ordinary equations

$$F(x, t, t_1, \dots, t_k) = a(n),$$

$$G(x, t, t_x, \dots, t_{[m]}) = b(x)$$

for some functions $a(n)$ and $b(x)$. This allows to solve (1) (see [9]).

The paper is organized as follows. In Section 2 we derive necessary and sufficient conditions on function $f(x, t, t_1, t_x)$ so that the chain (1) has four dimensional characteristic ring and in Section 3 we consider some applications of the derived conditions.

2. Chains admitting four dimensional x -algebra.

Suppose F is an x -integral of the chain (1) then its positive shifts and negative shifts $D^k F$, $k \in \mathbb{Z}$, are also x -integrals. So, looking for an x -integral it is convenient to assume that it depends on positive and negative shifts of t .

To express x derivatives of negative shifts we can apply D^{-1} to the chain (1) and obtain

$$t_x = f(x, t_{-1}, t, t_x).$$

Solving the above equation for t_{-1x} we get

$$t_{-1x} = g(x, t_{-1}, t, t_x).$$

Let $F(x, t, t_1, t_{-1}, \dots)$ be an x -integral of the chain (1). Then on solutions of (1) we have

$$D_x F = \frac{\partial F}{\partial x} + t_x \frac{\partial F}{\partial t} + t_{1x} \frac{\partial F}{\partial t_1} + t_{-1x} \frac{\partial F}{\partial t_{-1}} + t_{2x} \frac{\partial F}{\partial t_2} + t_{-2x} \frac{\partial F}{\partial t_{-2}} + \dots = 0$$

or

$$D_x F = \frac{\partial F}{\partial x} + t_x \frac{\partial F}{\partial t} + f \frac{\partial F}{\partial t_1} + g \frac{\partial F}{\partial t_{-1}} + Df \frac{\partial F}{\partial t_2} + D^{-1}g \frac{\partial F}{\partial t_{-2}} + \dots = 0.$$

Define a vector field

$$K = \frac{\partial}{\partial x} + t_x \frac{\partial}{\partial t} + f \frac{\partial}{\partial t_1} + g \frac{\partial}{\partial t_{-1}} + Df \frac{\partial}{\partial t_2} + D^{-1}g \frac{\partial}{\partial t_{-2}} + \dots, \quad (2)$$

then

$$D_x F = K F.$$

Note that F does not depend on t_x but the coefficients of K do depend on t_x . So we introduce a vector field

$$X = \frac{\partial}{\partial t_x} \quad (3)$$

The vector fields K and X generate the characteristic x -ring L_x .

Let us introduce some other vector fields from L_x .

$$C_1 = [X, K] \quad \text{and} \quad C_n = [X, C_{n-1}] \quad n = 2, 3, \dots \quad (4)$$

and

$$Z_1 = [K, C_1] \quad \text{and} \quad Z_n = [K, Z_{n-1}] \quad n = 2, 3, \dots \quad (5)$$

Thus

$$C_1 = \frac{\partial}{\partial t} + f_{tx} \frac{\partial}{\partial t_1} + g_{tx} \frac{\partial}{\partial t_{-1}} + \dots$$

$$C_2 = f_{tx} \frac{\partial}{\partial t_1} + g_{tx} \frac{\partial}{\partial t_{-1}} + \dots$$

$$Z_1 = (f_{txx} + t_x f_{txt} + f f_{tx t_1} - f_t - f_{tx} f_{t_1}) \frac{\partial}{\partial t_1} + (g_{txx} + t_x g_{tx t} + g g_{tx t_1} - g_t - g_{tx} g_{t_1}) \frac{\partial}{\partial t_{-1}} + \dots$$

and so on.

It is easy to see that if $f_{tx} \neq 0$ then the vector fields X , K , C_1 and C_2 are linearly independent and must form a basis of L_x provided $\dim L_x = 4$. By Lemma 3.6 in [6], if $f_{tx} = 0$ and $(f_{txx} + t_x f_{txt} + f f_{tx t_1} - f_t - f_{tx} f_{t_1}) = 0$ then $\dim L_x = 3$. So in the case $f_{tx} = 0$ we may assume $(f_{txx} + t_x f_{txt} + f f_{tx t_1} - f_t - f_{tx} f_{t_1}) \neq 0$. Then the vector fields X , K , C_1 and Z_1 are linearly independent and must form a basis of L_x provided $\dim L_x = 4$. We consider this two cases separately.

In the rest of the paper we assume that the characteristic ring L_x is four dimensional.

Remark 4 *It is convenient to check equalities between vector fields using the automorphism $D(\cdot)D^{-1}$. Direct calculations show that*

$$DXD^{-1} = \frac{1}{f_x} X,$$

$$DKD^{-1} = K - \frac{f_x + t_x f_t + f f_{t_1}}{f_{tx}} X.$$

The images of other vector fields under this automorphism can be obtained by commuting DXD^{-1} and DKD^{-1} .

2.1. $f(x, t, t_1, t_x)$ is non linear with respect to t_x .

Let $f(x, t, t_1, t_x)$ be non linear with respect to t_x , $f_{tx} \neq 0$. Then the vector fields X , K , C_1 and C_2 form a basis of L_x . For the algebra L_x to be spanned by X , K , C_1 and C_2 it is enough that C_3 and Z_1 are linear combinations of X , K , C_1 and C_2 . From the form of the vector fields it follows that we must have

$$C_3 = \lambda C_2 \quad \text{and} \quad Z_1 = \mu C_2$$

for some functions μ and λ . The conditions for the above equalities to hold are given by the following theorem.

Theorem 5 *The chain (1) with $f_{tx} \neq 0$ has characteristic ring L_x of dimension four if and only if the following conditions hold*

$$D\left(\frac{f_{tx} t_x}{f_{tx} t_x}\right) = \frac{f_{tx} t_x f_{tx} - 3f_{tx}^2}{f_{tx} f_{tx}^2}. \quad (6)$$

$$D \left(\frac{f_{xt_x} + t_x f_{t_x t} + f f_{t_x t_1} - f_t - f_{t_x} f_{t_1}}{f_{t_x t_x}} \right) = \frac{f_{xt_x} + t_x f_{t_x t} + f f_{t_x t_1} - f_t - f_{t_x} f_{t_1}}{f_{t_x t_x}} f_{t_x} - (f_x + t_x f_t + f_{t_1}). \quad (7)$$

The characteristic ring is generated by the vector fields X, K, C_1, C_2 .

Proof. By Remark 4 we have

$$\begin{aligned} DC_2 D^{-1} &= \frac{1}{f_{t_x}^2} C_2 - \frac{f_{t_x t_x}}{f_{t_x}^3} C_1 + \frac{f_{t_x t_x} f_t}{f_{t_x}^4} X \\ DC_3 D^{-1} &= \frac{1}{f_{t_x}^3} C_2 - \frac{3f_{t_x t_x}}{f_{t_x}^4} C_2 - \frac{f_{t_x t_x t_x} f_{t_x} - 3f_{t_x t_x}^2}{f_{t_x}^5} C_1 + f_t \frac{f_{t_x t_x t_x} f_{t_x} - 3f_{t_x t_x}^2}{f_{t_x}^6} X \\ DZ_1 D^{-1} &= \frac{1}{f_{t_x}} Z_1 - \left(\frac{m f_{t_x} + p}{f_{t_x}^2} \right) \left(C_1 - \frac{f_t}{f_{t_x}} X \right), \end{aligned}$$

where $p = \frac{f_x + t_x f_t + f f_{t_1}}{f_{t_x}}$ and $m = \frac{-(f_{xt_x} + t_x f_{t_x t} + f f_{t_x t_1}) + f_t + f_{t_x} f_{t_1}}{f_{t_x}}$.

The equality $C_3 = \lambda C_2$ implies that

$$DC_3 D^{-1} = (D\lambda) DC_2 D^{-1}. \quad (8)$$

Substituting expressions for $DC_2 D^{-1}$ and $DC_3 D^{-1}$ into (8) and comparing coefficients of C_1 , C_2 and X we obtain that λ satisfies

$$\begin{aligned} \lambda &= f_{t_x} (D\lambda) + \frac{3f_{t_x t_x}}{f_{t_x}} \\ (D\lambda) &= \frac{f_{t_x t_x t_x} f_{t_x} - 3f_{t_x t_x}^2}{f_{t_x t_x} f_{t_x}^2}. \end{aligned}$$

We can find λ and $D\lambda$ independently and condition that $D\lambda$ is a shift of λ leads to (6). The equality $Z_1 = \mu C_2$ implies that

$$DZ_1 D^{-1} = (D\mu) DC_2 D^{-1}. \quad (9)$$

Substituting expressions for $DC_2 D^{-1}$ and $DC_3 D^{-1}$ into (9) and comparing coefficients of C_1 , C_2 and X we obtain that μ satisfies

$$\mu - \frac{f_x + t_x f_t + f f_{t_1}}{f_{t_x}} = \frac{(D\mu)}{f_{t_x}}$$

and

$$-(f_{xt_x} + t_x f_{t_x t} + f f_{t_x t_1} - f_t - f_{t_x} f_{t_1}) + \frac{f_x + t_x f_t + f f_{t_1}}{f_{t_x}} f_{t_x t_x} = -\frac{f_{t_x t_x} (D\mu)}{f_{t_x}}$$

We can find μ and $D\mu$ independently and condition that $D\mu$ is a shift of μ leads to (7). \square

Remark 6 Let $\dim L_x = 4$ and $f_{t_x} \neq 0$. Then the characteristic ring L_x have the following multiplication table

	X	K	C_1	C_2
X	0	C_1	C_2	μC_2
K	$-C_1$	0	λC_2	ρC_2
C_1	$-C_2$	$-\lambda C_2$	0	ηC_2
C_2	$-\mu C_2$	$-\rho C_2$	$-\eta C_2$	0

where $\rho = \lambda\mu + X(\lambda)$ and $\eta = X(\rho) - K(\mu)$.

Example 7 Consider the following chain

$$t_{1x} = \frac{tt_x - \sqrt{t_x^2 - M^2}(t_1 + t)}{t_1}$$

introduced by Habibullin and Zheltukhina [10]. We can easily check that the function

$$f(t, t_1, t_x) = \frac{tt_x - \sqrt{t_x^2 - M^2}(t_1 + t)}{t_1}$$

satisfies the conditions of Theorem 5. Hence the corresponding x -algebra is four dimensional. The chain has the following x -integral

$$F = \frac{(t_1^2 - t^2)(t_1^2 - t_2^2)}{t_1^2}.$$

2.2. $f(x, t, t_1, t_x)$ is linear with respect to t_x .

Let $f(x, t, t_1, t_x)$ be linear with respect to t_x , $f_{t_x t_x} = 0$. Then vector fields X , K , C_1 and Z_1 form a basis of L_x . The condition $f_{t_x t_x} = 0$ also implies that the vector field $C_2 = 0$, see [6]. For the algebra L_x to be spanned by X , K , C_1 and Z it is enough that Z_2 is a linear combination of X , K , C_1 and Z_1 . From the form of the vector fields it follows that we must have

$$Z_2 = \alpha Z_1$$

for some function α . The conditions for the above equality to hold given by the following theorem.

Theorem 8 The chain (1) with $f_{t_x t_x} = 0$ has the characteristic ring L_x of dimension four if and only if the following condition hold

$$D \left(\frac{K(m)}{m} - m + \frac{f_t}{f_{t_x}} \right) = \frac{K(m)}{m} + m - f_{t_1}. \quad (10)$$

where $m = \frac{-(f_{xt_x} + t_x f_{t_x t} + f f_{t_x t_1}) + f_t + f_{t_x} f_{t_1}}{f_{t_x}}$. The characteristic ring is generated by the vector fields X, K, C_1, Z_1 .

Proof. By Remark 4 we have

$$DZ_1 D^{-1} = \frac{1}{f_{t_x}} Z_1 - \left(\frac{m f_{t_x} + p}{f_{t_x}^2} \right) \left(C_1 - \frac{f_t}{f_{t_x}} X \right),$$

and

$$DZ_2 D^{-1} = \left(K \left(\frac{1}{f_{t_x}} \right) + \frac{\alpha + m}{f_{t_x}} \right) Z_1 + \left(K \left(\frac{m}{f_{t_x}} \right) + \frac{m f_t}{f_{t_x}^2} - pX \left(\frac{m}{f_{t_x}} \right) \right) \left(C_1 - \frac{f_t}{f_{t_x}} X \right)$$

The equality $Z_2 = \alpha Z_1$ implies that

$$DZ_2D^{-1} = (D\alpha)DZ_1D^{-1}. \quad (11)$$

Substituting expressions for DZ_1D^{-1} and DZ_2D^{-1} into (11) and comparing coefficients of C_1 , Z_1 and X we obtain that α and $D(\alpha)$ satisfy

$$K\left(\frac{1}{f_{t_x}}\right) + \frac{m}{f_{t_x}} + \frac{\alpha}{f_{t_x}} = \frac{D(\alpha)}{f_{t_x}}$$

$$K\left(\frac{m}{f_{t_x}}\right) + \frac{mf_t}{f_{t_x}^2} = \frac{mD(\alpha)}{f_{t_x}}$$

We can find α and $D(\alpha)$ independently and condition that $D(\alpha)$ is a shift of α leads to (10). \square

Remark 9 Let $\dim L_x = 4$ and $f_{t_{xx}} = 0$. Then the characteristic ring L_x have the following multiplication table

	X	K	C_1	Z_1
X	0	C_1	0	0
K	$-C_1$	0	Z_1	αZ_1
C_1	0	$-Z_1$	0	$X(\alpha)Z_1$
Z_1	0	$-\alpha Z_1$	$-X(\alpha)Z_1$	0

Example 10 Consider the following chain

$$t_{1x} = t_x + e^{\frac{t+t_1}{2}}$$

introduced by Dodd and Bullough [2]. We can easily check that the function

$$f(t, t_1, t_x) = t_x + e^{\frac{t+t_1}{2}}$$

satisfies the conditions of Theorem 8. Hence the corresponding x -algebra is four dimensional. The chain has the following x -integral

$$F = e^{\frac{t_1-t}{2}} + e^{\frac{t_1-t_2}{2}}$$

3. Applications

The conditions derived in the previous section can be used to determine some restrictions on the form of the function $f(x, t, t_1, t_x)$ in (1).

Lemma 11 Let the chain (1) have four dimensional characteristic x -ring. Then

$$f = M(x, t, t_x)A(x, t, t_1) + t_x B(x, t, t_1) + C(x, t, t_1), \quad (12)$$

where M , A , B and C are some functions.

Proof. Let $f_{t_x t_x} \neq 0$ (if $f_{t_x t_x} = 0$ then f obviously has the above form). Since characteristic x -ring has dimension four the condition (6) holds. It is easy to see that (6) implies that $\frac{f_{t_x t_x t_x}}{f_{t_x t_x}}$ does not depend on t_1 . Hence

$$X(\ln |f_{t_x t_x}|) = M_1(x, t, t_x) \quad \text{and} \quad \ln |f_{t_x t_x}| = M_2(x, t, t_x) + A_1(x, t, t_1).$$

The last equality implies (12). \square

We can also put some restrictions on the shifts of the function $f(x, t, t_1, t_x)$ in (1).

Lemma 12 *Let the chain (1) have four dimensional characteristic x -ring and $f_{t_x t_x} \neq 0$. Then*

$$Df = -H_1(x, t, t_1, t_2)t_x + H_2(x, t, t_1, t_2)f + H_3(x, t, t_1, t_2), \quad (13)$$

where H_1 , H_2 and H_3 are some functions.

Proof. Note that the shift operator D and the vector field X satisfy

$$DX = \frac{1}{f_{t_x}}XD. \quad (14)$$

The condition (6) can be written as

$$DX(\ln |f_{t_x t_x}|) = \frac{1}{f_{t_x}}X(\ln |f_{t_x t_x}| - \ln |f_{t_x}|^3)$$

Using (14) we get

$$\frac{1}{f_{t_x}}XD(\ln |f_{t_x t_x}|) = \frac{1}{f_{t_x}}X(\ln |f_{t_x t_x}| - \ln |f_{t_x}|^3)$$

which implies that

$$X\left(\ln \left|f_{t_x}^3 \frac{Df_{t_x t_x}}{f_{t_x t_x}}\right|\right) = 0 \quad \text{or} \quad X\left(f_{t_x}^3 \frac{Df_{t_x t_x}}{f_{t_x t_x}}\right) = 0.$$

Thus $Df_{t_x t_x} = H_1(x, t, t_1, t_2)\frac{f_{t_x t_x}}{f_{t_x}^3}$. Since $Df_{t_x t_x} = DX(f_{t_x})$ and $\frac{f_{t_x t_x}}{f_{t_x}^3} = -\frac{1}{f_{t_x}}X\left(\frac{1}{f_{t_x}}\right)$ we can rewrite previous equality using (14) as

$$X\left(Df_{t_x} + H_1(x, t, t_1, t_2)\frac{1}{f_{t_x}}\right) = 0$$

which implies

$$Df_{t_x} = -H_1(x, t, t_1, t_2)\frac{1}{f_{t_x}} + H_2(x, t, t_1, t_2).$$

Writing

$$DX(f) = -H_1(x, t, t_1, t_2)\frac{1}{f_{t_x}} + H_2(x, t, t_1, t_2)\frac{f_{t_x}}{f_{t_x}}$$

and applying (14) as before we get

$$X(Df + H_1(x, t, t_1, t_2)t_x - H_2(x, t, t_1, t_2)f) = 0.$$

The last equality gives (13). \square

Note that the equality (13) can be written as

$$t_{2x} = H_2(x, t, t_1, t_2)t_{1x} - H_1(x, t, t_1, t_2)t_x + H_3(x, t, t_1, t_2).$$

References

- [1] Darboux G 1915 *Leçons sur la théorie générale des surfaces et les applications géométriques du calcul infinitesimal* **2** (Paris: Gautier Villas)
- [2] R K Dodd and R K Bullough 1976 *Proc. R. Soc. London Ser. A* **351** 499
- [3] Habibullin I T 2005 *SIGMA Symmetry Integrability Geom.: Methods Appl.* **1**
- [4] Habibullin I T and Pekcan A 2007 *Theoret. and Math. Phys.* **151** 781790
- [5] Habibullin I 2007 *Characteristic algebras of discrete equations. Difference equations, special functions and orthogonal polynomials* (Hackensack NJ World Sci. Publ.) 249-257
- [6] Habibullin I Zheltukhina N and Pekcan A 2008 *Turkish J. Math.* **32** 277292
- [7] Habibullin I Zheltukhina N and Pekcan A 2008 *J. Math. Phys.* **49** 102702
- [8] Habibullin I Zheltukhina N and Pekcan A 2009 *J. Math. Phys.* **50** 102710
- [9] Habibullin I Zheltukhina N and Sakieva A 2010 *J. Phys. A* **43** 434017
- [10] Habibullin I and Zheltukhina N 2014 Discretization of Liouville type nonautonomous equations *Preprint* nlin.SI:1402.3692v1
- [11] Shabat A B and Yamilov R I 1981 Exponential systems of type I and Cartan matrices *Preprint* BBAS USSR Ufa
- [12] Zhiber A B, Murtazina R D, Habibullin I T and Shabat A B 2012 *Ufa Math. J.* **4** 17-85