

Probing ultrafast dynamics in a solid-density plasma created by an intense femtosecond laser

Amitava Adak¹, Dave Blackman², Gourab Chatterjee¹, Prashant Kumar Singh¹, Amit D. Lad¹, P. Brijesh¹, A. P. L. Robinson³, John Pasley^{2,3} and G. Ravindra Kumar^{*1}

¹Tata Institute of Fundamental Research, Dr. Homi Bhabha Road, Colaba, Mumbai-400005, India

²York Plasma Institute, University of York, Heslington, York, YO10 5DQ, United Kingdom

³Central Laser Facility, RutherfordAppleton Laboratory, Chilton, Didcot, OX10 0QX, United Kingdom

*Email: grk@tifr.res.in

Abstract. We report a study on the dynamics of a near-solid density plasma using an ultraviolet (266 nm) femtosecond probe laser pulse, which can penetrate to densities of $\sim 10^{22} \text{ cm}^{-3}$, nearly an order of magnitude higher than the critical density of the 800 nm, femtosecond pump laser. Time-resolved probe-reflectivity from the plasma shows a rapid decay (picosecond-timescale) while the time-resolved reflected probe spectra show red shifts at early temporal delays and blue shifts at longer delays. This spectral behaviour of the reflected probe can be explained by a laser-driven shock moving inward and a subsequent hydrodynamic free expansion in the outward direction.

1. Introduction

Ultra-intense lasers can create extremely hot, high-density plasma while interacting with a solid. Extreme states of matter, achieved on a laboratory table-top by such interactions are very interesting to study, the measurement of the equation of state similar to that of an astrophysical object being a striking example[1]. Relativistic energy electron beams produced at the plasma critical surface in such interactions are crucial to the success of fast ignition[2] of the fusion pellet in inertial confinement fusion(ICF). A lot of experimental and theoretical studies have been performed in recent decades for the basic understanding of intense short-pulse-laser matter interaction[3] and for applications like particle acceleration[4, 5]. The hot dense plasma created via various laser absorption mechanisms[6] evolves very rapidly (on femtosecond and picosecond time-scales) and these dynamics can be monitored using the pump-probe technique[7, 8, 9].

In this paper, we investigate the temporal dynamics of a highly dense electron layer($n_e = 10^{22} \text{ cm}^{-3}$) inside a plasma created on an aluminium-coated BK7 glass target by a laser at relativistic light intensities. Doppler spectrometry of the reflected probe enables the observation of the ultrafast motion of its critical surface (high density layer) inside the hot dense plasma. A numerical simulation is performed which reproduces the results of the experiment.






Figure 1. Experimental setup: T-target, BS-Beamsplitter, M-mirror, L-lens, SM-spectrometer, PD- Photo diode.

2. Experiment

The experiment (Fig. 1) was carried out with a chirped-pulse-amplification-based 20 TW laser system (800 nm, 40 fs, 10 Hz) at Tata Institute of Fundamental Research. An extra Pockels' cell was used to obtain a 10^{-6} nanosecond intensity contrast. The pump pulse was focussed on a solid slab at 45° angle of incidence to a focal spot of $17 \mu\text{m}$ (FWHM) to obtain intensities of $\sim 10^{18} \text{ W/cm}^2$. A small portion of the laser pulse (5 %) was extracted using a beam-splitter, up-converted to 266 nm, and focussed to a spot of $60 \mu\text{m}$ at the interaction point at near-normal incidence using a fused-silica lens. The reflected probe pulse was then split into two parts and fed to a photodiode and a high-resolution ultraviolet spectrometer. A delay line was introduced in the path of the probe to change the relative temporal delay between the pump and the probe. The focussed probe intensity was $\sim 10^{11} \text{ W/cm}^2$. Spatial and temporal overlap was achieved by looking at the reflected probe intensity from a plasma created on a dielectric slab at relatively lower pump intensity (10^{17} W/cm^2). We define the temporal zero where the reflectivity shows a sudden spike. In this experiment we observed the probe reflectivity and spectrum from a solid-density plasma on aluminium-coated BK-7 target.

3. Results & Discussions

Figure 2(a) shows reflection of the probe as a function of probe delay with respect to the pump from a super critical layer ($n_e=10^{22} \text{ cm}^{-3}$) in the plasma. Target: Al-coated 5 mm BK-7 glass target. The probe reflection shows an exponential decay ($\tau = 5 \text{ ps}$) as the plasma evolves after excitation by pump. The reflectivity of the normally incident probe can be written as [10]

$$R \propto \exp\left(-\frac{8\nu_{ei}^* L}{3c}\right) \quad (1)$$

where ν_{ei}^* is the effective electron ion collision frequency. L is the spatial scale length of plasma over which the probe gets absorbed. The collision frequency is a function of electron density




Figure 2. (a) Reflected probe intensity as a function of probe delay. The blue line is the exponential fit to the experimental data (blue asterisks). The decay time is measured to be 5 ps from the fit. (b) A similar plot for the velocity of the probe-critical-layer is shown. These velocities (black open circles) are calculated from the Doppler shifts of the ultraviolet probe measured from pump-probe Doppler spectrometry.

and temperature. In this context of fast time scale (few picosecond) probe reflectivity is mainly dependent on L , if we assume quasi-static values of density and temperature.

The velocity of the supercritical layer was measured by pump-probe Doppler spectrometry[8]. The velocity can be expressed as,

$$v_{exp} = -0.5c \frac{\Delta\lambda}{\lambda} \quad (2)$$

Where $\Delta\lambda$ is the Doppler shift from the experiment and $\lambda = 266$ nm in our case. Figure 2(b) shows the velocity of the probe-critical-layer, calculated from the Doppler shift at various probe delays. At initial few picoseconds, the probe critical-surface moves deeper into the plasma (negative velocity) riding on a non-relativistic shock. At subsequent times, the critical surface moves towards vacuum with the freely expanding plasma (positive velocity).

4. Simulation

Figure 3 shows the velocity of the probe critical-layer results from 1-D hybrid simulations. First, the HYADES code was run on a $500 \mu\text{m}$ silicate target. The output from this (ion, mass and electron density) was then interpolated onto a regular grid for use in an 1-D PIC code (ELPS) to find the hotspot formed by the pump laser. The code was run with 800 nm, 30 fs, $2 \times 10^{18} \text{W}/\text{cm}^2$ laser pulse. A density spike was observed and its motion was simulated by a Lagrangian hydro code. These 1-D simulations calculate a slightly higher velocity of the critical-layer of the 266 nm probe beam than the experimental observation. The sign reversal of the velocity is around 6 ps, which is close to the experimental result.




Figure 3. The simulated velocity of the probe-critical-layer as a function of the probe delay. The sign convention: outward motion corresponds to positive velocity.

5. Conclusions

In this study, we observe ultrafast motion of a supercritical (10^{22} cm^{-3}) electron layer in a plasma created by a high-intensity, femtosecond laser pulse on a solid target. We see laser-driven density pile-up and propagation of a non-relativistic shock inside the solid. 1-D hybrid HYADES-PIC-HYDRO simulations support the experiment results.

6. Acknowledgments

AA and ADL thank the ‘Strong Field Science’ (11P-1401) programme of the DAE, Government of India for support for participation in IFSA2013. GRK acknowledges support from a J.C. Bose Fellowship grant from DST, Government of India.

References

- [1] Paul McKenna *et. al.*, *Laser-Plasma Interactions and Applications* (Springer Switzerland, Heidelberg, New York, Dordrecht, London, 2013).
- [2] M. Tabak *et. al.*, *Phys. Plasmas* **1**, 1626 (1994).
- [3] P. Gibbon *et. al.*, *Plasma Phys. Control. Fusion* **38**, 769-793 (1996).
- [4] T. Tajima *et. el.*, *Phys. Rev. Lett.* **43**, 267 (1979).
- [5] V. Malka *et. el.*, *Science* **298**, 1596 (2002).
- [6] S. C. Wilks *et. el.*, *IEEE JOURNAL OF QUANTUM ELECTRONICS*, **33**, 1954 (1997).
- [7] A. S. Sandhu *et. al.*, *Phys. Rev. Lett.* **89**, 225002 (2002).
- [8] S. Mondal *et. al.*, *Phys. Rev. Lett.* **105**, 105002 (2010).
- [9] Y. Ping *et. al.*, *Phys. Rev. Lett.* **109**, 145006 (2012).
- [10] W. L. Kruer, *The physics of laser plasma interactions* (Boulder, Colorado, 2003).