

# Neutrino oscillations: from an historical perspective to the present status

**S Bilenky**

Joint Institute for Nuclear Research, Dubna, R-141980, Russia; TRIUMF 4004 Wesbrook Mall, Vancouver BC, V6T 2A3 Canada

**Abstract.** The history of neutrino mixing and oscillations is briefly presented. Basics of neutrino mixing and oscillations and convenient formalism of neutrino oscillations in vacuum is given. The role of neutrino in the Standard Model and the Weinberg mechanism of the generation of the Majorana neutrino masses are discussed.

## 1. Introduction. On the history of neutrino oscillations

Discovery of the neutrino oscillations in the atmospheric Super-Kamiokande [1], solar SNO [2] and reactor KamLAND [3] experiments was a first evidence in favor of a beyond the Standard Model physics in particle physics. Neutrino oscillations were further studied in the long baseline accelerator K2K [4], MINOS [5] and T2K [6] experiments. With the measurement of the small parameter  $\sin^2 \theta_{13}$  in the accelerator T2K [6], reactor Daya Bay [7], RENO [8] and Double Chooze [9] experiments investigation of neutrino oscillations enters into a new era, era of high precision measurements. The 2015 Nobel Prize to T. Kajita and A. McDonald "for the discovery of neutrino oscillations, which shows that neutrinos have mass" is a very important event for the neutrino community which will attract new people and give a great boost to the field.

Idea of neutrino oscillations was first proposed by B.Pontecorvo in 1957-58 soon after the theory of the two-component neutrino was proposed [10] and confirmed by the Goldhaber et al experiment [11]. B.Pontecorvo looked in the lepton world for a phenomena analogous to  $K^0 \rightleftharpoons \bar{K}^0$  oscillations. In the paper [12] he considered muonium ( $\mu^+e^-$ ) to antimuonium ( $\mu^-e^+$ ) transition. In this paper he mentioned a possibility of the neutrino oscillations. Special paper dedicated to neutrino oscillations was published by B.Pontecorvo in 1958 [13]. At that time only one type of neutrino was known. B.Pontecorvo assumed that in addition to the usual weak interaction *exist a much weaker interaction which does not conserve the lepton number*. Assuming maximum mixing (by the analogy with  $K^0 - \bar{K}^0$ ) he concluded that "...neutrino and antineutrino are particle mixtures, i.e. symmetrical and antisymmetrical combinations of two truly neutral Majorana particles  $\nu_1$  and  $\nu_2$ ...":

$$|\bar{\nu}_R\rangle = \frac{1}{\sqrt{2}}(|\nu_1\rangle + |\nu_2\rangle), \quad |\nu_R\rangle = \frac{1}{\sqrt{2}}(|\nu_1\rangle - |\nu_2\rangle) \quad (1)$$

Here  $|\bar{\nu}_R\rangle$  is the state of the right-handed antineutrino,  $|\nu_R\rangle$  is the state of right-handed neutrino, a particle which does not take part in the weak interaction (later B.Pontecorvo proposed the name sterile for such neutrinos),  $|\nu_{1,2}\rangle$  are states of Majorana neutrinos with



small masses  $m_{1,2}$ . As a result of the mixing (1), oscillations  $\bar{\nu}_R \leftrightarrow \nu_R$  (sterile) become possible. B.Pontecorvo discussed a possibility to check a hypothesis of neutrino oscillations in the reactor neutrino experiments. In 1958 the only known sources of neutrinos were reactors and the sun. B.Pontecorvo finished the paper [13] with the following remark "...effects of transformation of neutrino into antineutrino and vice versa may be unobservable in the laboratory because of large values of  $R$  (oscillation length) , but will certainly occur, at least, on an astronomic scale."

In 1962 the idea of neutrino masses and mixing was discussed by Maki, Nakagawa and Sakata [14]. Their proposal was based on the Nagoya model in which nucleons were considered as bound states of a vector boson and neutrino with definite mass. MNS assumed that the fields of the weak neutrinos  $\nu_e$  and  $\nu_\mu$  are connected with the fields of neutrinos with definite masses  $\nu_1$  and  $\nu_2$  (they called them true neutrinos) by the orthogonal transformation

$$\nu_e = \cos \theta \nu_1 + \sin \theta \nu_2, \quad \nu_\mu = -\sin \theta \nu_1 + \cos \theta \nu_2. \quad (2)$$

The phenomenon of neutrino oscillations was not considered in [14]. However, MNS discussed a possibility of "virtual transmutation" of  $\nu_\mu$  into  $\nu_e$ . They estimated a time of this transition and discussed how a possible  $\nu_\mu \rightarrow \nu_e$  transition would influence the interpretation of the results of the Brookhaven experiment [15],<sup>1</sup> which was going on at the time when the MNS paper was written.

In 1967 B.Pontecorvo published the second paper on neutrino oscillations [16]. In this paper he discussed flavor neutrino oscillations  $\nu_\mu \leftrightarrow \nu_e$  and also oscillations between flavor and sterile neutrinos ( $\nu_e \leftrightarrow \bar{\nu}_{eL}$  etc). In the paper [16] solar neutrino oscillations were considered. *Before the first results of the Davis solar neutrino experiment appeared* , B.Pontecorvo pointed out that because of neutrino oscillations the flux of the solar  $\nu_e$ 's could be two times smaller than the expected flux. Thus, he anticipated "the solar neutrino problem".

In the Gribov and Pontecorvo paper [17] it was suggested that only active left-handed neutrinos  $\nu_e$  and  $\nu_\mu$  and right-handed antineutrinos  $\bar{\nu}_e$  and  $\bar{\nu}_\mu$  exist in nature (no sterile neutrinos). It was assumed that exist a (miliweak) interaction which does not conserve lepton numbers. After the diagonalization of such an interaction the authors came to the mixing relation

$$\nu_{eL} = \cos \xi \phi_{1L} + \sin \xi \phi_{2L}, \quad \nu_{\mu L} = -\sin \xi \phi_{1L} + \cos \xi \phi_{2L}, \quad (3)$$

where  $\xi$  is the mixing angle and  $\phi_1$  and  $\phi_2$  are fields of the Majorana neutrinos with masses  $m_1$  and  $m_2$ . They calculated the probability of  $\nu_e$  to survive in vacuum. The case of the maximum mixing ( $\xi = \pi/4$ ), analogous to the  $K^0 - \bar{K}^0$  case, was considered as the most attractive one. Under this assumption the oscillations of solar neutrinos were discussed.

In the seventies and eighties idea of neutrino masses and oscillations was further developed in Dubna in the papers [18]. In addition to the Gribov-Pontecorvo scheme of the neutrino mixing, based on the Majorana mass term, neutrino mixing based on the Dirac mass term and the most general Dirac and Majorana mass term were considered. Possible reactor, accelerator, solar and atmospheric experiments on the search for neutrino oscillations were discussed. Our general point of view, which we advocated in our papers and in the first review on neutrino oscillations [19] was the following:

- (i) There are no principles which require that neutrinos be massless particles. It is plausible that neutrinos have small nonzero masses.
- (ii) Neutrino oscillations is an interference phenomenon. Search for neutrino oscillations is the most sensitive method to search for extremely small mass-squared differences.

<sup>1</sup> As it is well known, in this experiment it was discovered that  $\nu_\mu$  and  $\nu_e$  are different particles.

- (iii) Experiments with neutrinos from different sources are sensitive to different neutrino mass-squared differences. Experiments on the search for neutrino oscillations must be performed with neutrinos from all existing sources.<sup>2</sup>

## 2. Neutrino mixing

Neutrino oscillations are based on the mixing of neutrino fields

$$\nu_{lL}(x) = \sum_i U_{li} \nu_{iL}(x), \quad (4)$$

Here  $U$  is a unitary mixing matrix and  $\nu_i(x)$  is the field of neutrinos (Dirac or Majorana) with mass  $m_i$ .

The flavor neutrino fields  $\nu_{lL}(x)$  ( $l = e, \mu, \tau$ ) enter into the Standard Model CC and NC interactions

$$\mathcal{L}_I^{CC} = -\frac{g}{\sqrt{2}} j_\alpha^{CC} W^\alpha + \text{h.c.}, \quad \mathcal{L}_I^{NC} = -\frac{g}{2 \cos \theta_W} j_\alpha^{NC} Z^\alpha. \quad (5)$$

Here

$$j_\alpha^{CC} = \sum_{l=e,\mu,\tau} \bar{\nu}_{lL} \gamma_\alpha l_L, \quad j_\alpha^{NC} = \sum_{l=e,\mu,\tau} \bar{\nu}_{lL} \gamma_\alpha \nu_{lL} \quad (6)$$

are charged leptonic and neutral neutrino currents.

The neutrino mixing takes place if in the total Lagrangian there is a *mass term nondiagonal over flavor neutrino fields*. In the case of the charged particles (leptons and quarks) only Dirac mass terms are possible. Because the electric charges of neutrinos are equal to zero three different neutrino mass terms are possible (see [20, 21]).

### Dirac mass term

$$\mathcal{L}^D = - \sum_{l',l=e,\mu,\tau} \bar{\nu}_{l'L} M_{l'l}^D \nu_{lR} + \text{h.c.}, \quad (7)$$

where  $M^D$  is a complex, nondiagonal,  $3 \times 3$  matrix. After the diagonalization of the matrix  $M^D$  we have

$$\nu_{lL}(x) = \sum_{i=1}^3 U_{li} \nu_{iL}(x). \quad (8)$$

Here  $U$  is the unitary PNMS mixing matrix and  $\nu_i(x)$  is the Dirac field with the mass  $m_i$ . The Lagrangian  $\mathcal{L}^D$  conserves the total lepton number  $L = L_e + L_\mu + L_\tau$ . Neutrino  $\nu_i$  and antineutrino  $\bar{\nu}_i$  differ by the lepton number:  $L(\nu_i) = 1$ ,  $L(\bar{\nu}_i) = -1$ .

### Majorana mass term

$$\mathcal{L}^M = -\frac{1}{2} \sum_{l',l=e,\mu,\tau} \bar{\nu}_{l'L} M_{l'l}^M (\nu_{lL})^c + \text{h.c.}, \quad (9)$$

where  $M^M$  is a complex, nondiagonal, symmetrical  $3 \times 3$  matrix and  $(\nu_{lL})^c = C \bar{\nu}_{lL}^T$  is the conjugated field. The mass term (9) violates not only flavor lepton numbers but also the total lepton number  $L$ . After the diagonalization of the matrix  $M^M$  we have

$$\nu_{lL}(x) = \sum_{i=1}^3 U_{li} \nu_{iL}(x). \quad (10)$$

<sup>2</sup> As we know, after heroic efforts of many people this strategy led to the discovery of neutrino oscillations.

Here  $U$  is a unitary  $3 \times 3$  mixing matrix and

$$\nu_i(x) = \nu_i^c(x) \quad (11)$$

is the Majorana field with the mass  $m_i$  ( $\nu_i \equiv \bar{\nu}_i$ ).

The most general **Dirac and Majorana mass term**

$$\mathcal{L}^{\text{D+M}} = \mathcal{L}^{\text{M}} + \mathcal{L}^{\text{D}} - \frac{1}{2} \sum_{l', l=e, \mu, \tau} \overline{(\nu_{l'R})^c} M_{l'l}^{\text{R}} \nu_{lR} + \text{h.c.} \quad (12)$$

( $M^{\text{R}}$  is a complex symmetrical matrix) violates lepton numbers and require left-handed and right-handed neutrino fields. After the diagonalization of the mass term  $\mathcal{L}^{\text{D+M}}$  we find

$$\nu_{lL}(x) = \sum_{i=1}^6 U_{li} \nu_{iL}(x), \quad (\nu_{lR}(x))^c = \sum_{i=1}^6 U_{\bar{l}i} \nu_{iL}(x). \quad (13)$$

Here  $U$  is a unitary  $6 \times 6$  matrix and  $\nu_i(x) = \nu_i^c(x)$  is the field of a Majorana lepton with definite mass.

The mixing (13) open different possibilities: the seesaw possibility of the generation of small neutrino masses [22], a possibility of transitions of flavor neutrinos into sterile states etc.

Let us notice that the Dirac mass term can be generated by the standard Higgs mechanism. The Majorana and the Dirac and Majorana mass terms can be generated only by a beyond the SM mechanisms.

### 3. Flavor neutrino states

There exist different methods of the derivation (of the same) expression for transition probabilities. We will present here a method based on the notion of *the coherent flavor neutrino states* (see [21])

$$|\nu_l\rangle = \sum_i U_{li}^* |\nu_i\rangle, \quad l = e, \mu, \tau \quad (14)$$

Here  $|\nu_i\rangle$  is the state of neutrino (Dirac or Majorana) with mass  $m_i$ , momentum  $\vec{p}$  and energy  $E_i = \sqrt{p^2 + m_i^2} \simeq E + \frac{m_i^2}{2E}$  ( $E = p$ ), and  $|\nu_l\rangle$  is the state the flavor neutrino  $\nu_l$  which is produced together with  $l^+$  in a CC weak decay ( $\pi^+ \rightarrow \mu^+ + \nu_\mu$  etc) or produces  $l^-$  in a CC neutrino reaction ( $\nu_\mu + N \rightarrow \mu^- + X$  etc).

The relation (14) is valid if neutrino mass-squared differences are so small that in weak decays production of neutrinos with different masses can not be resolved. It follows from the Heisenberg uncertainty relation that this condition is satisfied in neutrino oscillation experiments with neutrino energies many orders of magnitude larger than neutrino masses.

The possibility to resolve small neutrino mass-squared differences is based on the time-energy uncertainty relation (see [24])

$$\Delta E \Delta t \gtrsim 1. \quad (15)$$

Here  $\Delta t$  is a time interval during which the state with the energy uncertainty  $\Delta E$  is significantly changed. In the case of neutrino beams from (15) we find

$$|\Delta m_{ki}^2| \frac{L}{2E} \gtrsim 1, \quad (16)$$

where  $L \simeq \Delta t$  is the distance between a neutrino source and neutrino detector. For "atmospheric" and "solar" mass-squared differences  $\Delta m_A^2 \simeq 2.4 \cdot 10^{-3} \text{ eV}^2$  and  $\Delta m_S^2 \simeq$

$7.5 \cdot 10^{-5} \text{ eV}^2$  the condition (16) is satisfied in the atmospheric Super-Kamiokande [1], long baseline accelerator K2K [4], MINOS [5], T2K [6], reactor KamLAND [3], Daya Bay [7], RENO [8] Double Chooze [9] and other neutrino oscillation experiments.

We will finish this section with a remark about the states of sterile neutrinos which (by definition) do not interact with leptons and quarks via the SM interaction. If in addition to the flavor neutrinos  $\nu_l$  sterile neutrinos  $\nu_s$  exist, their states *are determined* as follows

$$|\nu_s\rangle = \sum_{i=1}^{3+n_s} U_{si}^* |\nu_i\rangle, \quad s = s_1, s_1, \dots \quad (17)$$

where  $U$  is a unitary  $(3 + n_s) \times (3 + n_s)$  matrix. The states of active and sterile neutrinos (14) and (17) satisfy the condition

$$\langle \alpha' | \alpha \rangle = \delta_{\alpha'\alpha}, \quad \alpha' \alpha = e, \mu, \tau, s_1, s_1, \dots, s_{n_s}. \quad (18)$$

Neutrino oscillations is a direct consequence of the fact that flavor (and sterile) neutrinos are described by coherent states (14) and (17).

#### 4. Neutrino oscillations in vacuum

Let us assume that at the initial time  $t = 0$  a flavor neutrino  $\nu_\alpha$  was produced. In the general case of flavor and sterile neutrinos at the time  $t$  we have

$$|\nu_\alpha\rangle_t = e^{-iH_0 t} |\nu_\alpha\rangle = \sum_{i=1}^{3+n_s} |\nu_i\rangle e^{-iE_i t} U_{\alpha i}^* = \sum_{\alpha'} |\alpha'\rangle \left( \sum_{i=1}^{3+n_s} U_{\alpha' i} e^{-iE_i t} U_{\alpha i}^* \right). \quad (19)$$

Thus, for the  $\nu_\alpha \rightarrow \nu_{\alpha'}$  transition probability we find

$$P(\nu_\alpha \rightarrow \nu_{\alpha'}) = \left| \sum_{i=1}^{3+n_s} U_{\alpha' i} e^{-iE_i t} U_{\alpha i}^* \right|^2 \quad (20)$$

We will present here convenient expression for  $\overset{(-)}{\nu}_\alpha \rightarrow \overset{(-)}{\nu}_{\alpha'}$  transition probability (see [23]). From (20) we have

$$P(\nu_\alpha \rightarrow \nu_{\alpha'}) = \left| \sum_{i=1}^{3+n_s} U_{\alpha' i} e^{-2i\Delta_{pi}} U_{\alpha i}^* \right|^2 = \left| \delta_{\alpha'\alpha} - 2i \sum_i U_{\alpha' i} e^{-i\Delta_{pi}} \sin \Delta_{pi} U_{\alpha i}^* \right|^2 \quad (21)$$

where  $p$  is arbitrary, fixed index and

$$\Delta_{pi} = \frac{\Delta m_{pi}^2 L}{4E}, \quad \Delta m_{pi}^2 = m_i^2 - m_p^2. \quad (22)$$

Let us notice that in Eq. (21)

- $i \neq p$ ,
- we extract the common phase  $e^{-\frac{im_p^2 L}{2E}}$ ,
- we used the unitarity condition  $\sum_i U_{\alpha' i} U_{\alpha i}^* = \delta_{\alpha'\alpha}$ .

From (21) we find

$$\begin{aligned}
 P(\bar{\nu}_\alpha \rightarrow \bar{\nu}_{\alpha'}) &= \delta_{\alpha'\alpha} - 4 \sum_i |U_{\alpha i}|^2 (\delta_{\alpha'\alpha} - |U_{\alpha' i}|^2) \sin^2 \Delta_{pi} \\
 &+ 8 \sum_{i>k} [\text{Re}(U_{\alpha' i} U_{\alpha i}^* U_{\alpha' k}^* U_{\alpha k}) \cos(\Delta_{pi} - \Delta_{pk}) \\
 &\pm \text{Im}(U_{\alpha' i} U_{\alpha i}^* U_{\alpha' k}^* U_{\alpha k}) \sin(\Delta_{pi} - \Delta_{pk})] \sin \Delta_{pi} \sin \Delta_{pk}.
 \end{aligned} \tag{23}$$

Here +(-) sign refer to  $\nu_\alpha \rightarrow \nu_{\alpha'}$  ( $\bar{\nu}_\alpha \rightarrow \bar{\nu}_{\alpha'}$ ) transition.

From our point of view there are some advantages of the expression (23) with respect to the standard expression (for the standard expression see [25]).

- (i) Only independent mass-squared differences enter into this expression.
- (ii) The unitarity condition is fully explored in (23). As a result only independent terms enter into this expression.

We will consider now the most important case of the three-neutrino mixing. Usually neutrino masses are labeled in such a way that  $m_2 > m_1$  and solar ("small") mass-squared difference is determined as follows

$$m_2^2 - m_1^2 = \Delta m_{12}^2 \equiv \Delta m_S^2. \tag{24}$$

For the neutrino mass spectrum there are two possibilities:

- (i) Normal spectrum (NS) :  $\Delta m_S^2$  is the difference between square of masses of the lightest neutrinos. In this case  $m_3 > m_2 > m_1$ .
- (ii) Inverted spectrum (IS):  $\Delta m_S^2$  is the difference between square of masses of the heaviest neutrinos. In this case  $m_2 > m_1 > m_3$ .

We will determine the atmospheric ("large") neutrino mass squared difference in the following way

$$NS : \Delta m_A^2 = \Delta m_{23}^2, \quad IS : \Delta m_A^2 = |\Delta m_{13}^2|. \tag{25}$$

Let us notice that there exist different definition of this quantity in the literature

- (i) The Bari group [26] determines atmospheric mass-squared difference as follows

$$(\Delta m_A^2)' = \frac{1}{2} |\Delta m_{13}^2 + \Delta m_{23}^2| = \Delta m_A^2 + \frac{1}{2} \Delta m_S^2. \tag{26}$$

- (ii) The NuFit group [27] determines atmospheric mass-squared difference in the following way

$$(\Delta m_A^2)'' = \Delta m_{13}^2 \text{ (NS)} = |\Delta m_{23}^2| \text{ (IS)} = \Delta m_A^2 + \Delta m_S^2. \tag{27}$$

- (iii) In [28] the parameter  $\Delta m_{ee}^2$  were introduced. It is determined as followed

$$\Delta m_{ee}^2 = \cos^2 \theta_{12} \Delta m_{13}^2 + \sin^2 \theta_{12} \Delta m_{23}^2 \tag{28}$$

The parameter  $\Delta m_{ee}^2$  is connected with  $\Delta m_A^2$  and  $\Delta m_S^2$  by the relations

$$\Delta m_{ee}^2 = \Delta m_A^2 + \cos^2 \theta_{12} \Delta m_S^2 \text{ (NS)}, \quad |\Delta m_{ee}^2| = \Delta m_A^2 + \sin^2 \theta_{12} \Delta m_S^2 \text{ (IS)}. \tag{29}$$

As it is seen from (25), (26) (27) (29) different definitions of "large" mass-squared difference differ only by a few %. However, neutrino oscillation experiments enter now into precision era when neutrino oscillation parameters will be measured with % accuracy. We believe that the consensus in definition of "large" neutrino mass-squared difference must be found.

For the probability of the transition  $\bar{\nu}_l \rightarrow \bar{\nu}_{l'}$  ( $l, l' = e, \mu, \tau$ ) in the case of normal and inverted mass spectra from (23) we find, correspondingly, the following expressions

$$\begin{aligned}
 P^{NS}(\bar{\nu}_l \rightarrow \bar{\nu}_{l'}) &= \delta_{ll'} - 4|U_{l3}|^2(\delta_{ll'} - |U_{l'3}|^2) \sin^2 \Delta_A \\
 &- 4|U_{l1}|^2(\delta_{ll'} - |U_{l'1}|^2) \sin^2 \Delta_S - 8 [\text{Re} (U_{l'3}U_{l3}^*U_{l'1}^*U_{l1}) \cos(\Delta_A + \Delta_S) \\
 &\pm \text{Im} (U_{l'3}U_{l3}^*U_{l'1}^*U_{l1}) \sin(\Delta_A + \Delta_S)] \sin \Delta_A \sin \Delta_S,
 \end{aligned} \tag{30}$$

and

$$\begin{aligned}
 P^{IS}(\bar{\nu}_l \rightarrow \bar{\nu}_{l'}) &= \delta_{ll'} - 4|U_{l3}|^2(\delta_{ll'} - |U_{l'3}|^2) \sin^2 \Delta_A \\
 &- 4|U_{l2}|^2(\delta_{ll'} - |U_{l'2}|^2) \sin^2 \Delta_S - 8 [\text{Re} (U_{l'3}U_{l3}^*U_{l'2}^*U_{l2}) \cos(\Delta_A + \Delta_S) \\
 &\mp \text{Im} (U_{l'3}U_{l3}^*U_{l'2}^*U_{l2}) \sin(\Delta_A + \Delta_S)] \sin \Delta_A \sin \Delta_S.
 \end{aligned} \tag{31}$$

The transition probabilities (30) and (31) are the sum of atmospheric, solar and interference terms. Notice that expression (31) can be obtained from (30) by the change  $U_{l1} \rightarrow U_{l2}$  and  $(\pm) \rightarrow (\mp)$  in the last term.

The values of the oscillation parameters obtained from global analysis of existing data by the NuFit group [27] are presented in the Table 1.

**Table 1.** Values of neutrino oscillation parameters obtained in [27] from the global fit of existing data

Parameter	Normal Spectrum	Inverted Spectrum
$\sin^2 \theta_{12}$	$0.304^{+0.013}_{-0.012}$	$0.304^{+0.013}_{-0.012}$
$\sin^2 \theta_{23}$	$0.452^{+0.052}_{-0.028}$	$0.579^{+0.025}_{-0.037}$
$\sin^2 \theta_{13}$	$0.0218^{+0.0010}_{-0.0010}$	$0.0219^{+0.0011}_{-0.0010}$
$\delta$ (in $^\circ$ )	$(306^{+39}_{-70})$	$(254^{+63}_{-62})$
$\Delta m_S^2$	$(7.50^{+0.19}_{-0.17}) \cdot 10^{-5} \text{ eV}^2$	$(7.50^{+0.19}_{-0.17}) \cdot 10^{-5} \text{ eV}^2$
$\Delta m_A^2$	$(2.457^{+0.047}_{-0.047}) \cdot 10^{-3} \text{ eV}^2$	$(2.449^{+0.048}_{-0.047}) \cdot 10^{-3} \text{ eV}^2$

## 5. Neutrino and the Standard Model

After the discovery of the Higgs boson at LHC the Standard Model acquired the status of the theory of elementary particles in the electroweak range (up to  $\sim 300 \text{ GeV}$ ). The Standard Model is based on the following principles:

- Local gauge symmetry.
- Unification of the weak and electromagnetic interactions.
- Spontaneous breaking of the electroweak symmetry.

It was suggested in [30] that in the framework of these principles nature choose the simplest, most economical possibilities. The Standard Model started with the theory of the two-component neutrino. The two-component, massless, Weil neutrino is the *simplest possibility for the particle with spin 1/2*: only two degrees of freedom. The local  $SU_L(2)$  group with the lepton doublets

$$\psi_{eL}^{lep} = \begin{pmatrix} \nu'_{eL} \\ e'_L \end{pmatrix}, \quad \psi_{\mu L}^{lep} = \begin{pmatrix} \nu'_{\mu L} \\ \mu'_L \end{pmatrix}, \quad \psi_{\tau L}^{lep} = \begin{pmatrix} \nu'_{\tau L} \\ \tau'_L \end{pmatrix} \tag{32}$$

and corresponding quark doublets is the *simplest possibility* which allows to include charged leptons and quarks in addition to neutrinos.

In order to unify weak and electromagnetic interactions we need to enlarge the symmetry group: in electromagnetic currents of *charged particles* enter left-handed and right-handed fields. The simplest enlargement is the  $SU_L(2) \times U_Y(1)$  group where  $U_Y(1)$  is the group of the weak hypercharge  $Y$  determined by the Gell-Mann-Nishijima relation  $Q = T_3 + \frac{1}{2}Y$ . Neutrinos have no electromagnetic interaction. *Unification of the weak and electromagnetic interactions does not require right-handed neutrino fields.* The SM interaction of leptons, neutrinos and quarks with gauge vector bosons *is the minimal interaction* compatible with the local  $SU_L(2) \times U_Y(1)$  invariance.

The SM mechanism of the mass generation is the Brout-Englert-Higgs mechanism based on the assumption of the existence of scalar Higgs fields. In order to generate masses of  $W^\pm$  and  $Z^0$  bosons we need to have three (Goldstone) degrees of freedom. *Minimal possibility is a doublet of complex Higgs fields* (four degrees of freedom). With this assumption one scalar, neutral Higgs boson is predicted. This prediction is in a good agreement with existing LHC data.

Masses of  $W^\pm$  and  $Z^0$  bosons are given in the SM by the relations

$$m_W = \frac{1}{2}g v, \quad m_Z = \frac{1}{2}\sqrt{g^2 + g'^2} v = \frac{g}{2 \cos \theta_W} v, \quad (33)$$

where  $v = (\sqrt{2}G_F)^{-1/2} = 246$  GeV is the parameter which characterizes the scale of the electroweak symmetry breaking. Lepton and quark masses and mixing are due to  $SU_L(2) \times U_Y(1)$  invariant Yukawa interactions which generate Dirac mass terms. For the charged leptons we have

$$\mathcal{L}_Y^{lep} = - \sum_l m_l \bar{l} l, \quad (34)$$

where  $m_l = y_l v$  and  $y_l$  is the Yukawa constant. *Neutrinos in the minimal SM after spontaneous breaking of the electroweak symmetry remain two-component, massless, Weyl particles.*

## 6. The Weinberg mechanism of the neutrino mass generation

In the framework of the minimal SM neutrino masses and mixing can be generated only by a beyond the SM mechanism. The most general method which allows to describe effects of a beyond the SM physics is the method of the effective Lagrangian. The effective Lagrangian is a  $SU_L(2) \times U_Y(1)$  invariant, dimension five or more local operator built from SM fields. In order to built the effective Lagrangian which generate a neutrino mass term we must use the lepton doublets (32) and the Higgs doublet

$$\phi = \begin{pmatrix} \phi_+ \\ \phi_0 \end{pmatrix} \quad (35)$$

The only effective Lagrangian which generate the neutrino mass term has the form [29]

$$\mathcal{L}_I^{\text{eff}} = -\frac{1}{\Lambda} \sum_{l_1, l_2} (\bar{\psi}_{l_1 L}^{lep} \tilde{\phi}) Y_{l_1 l_2} (\tilde{\phi}^T (\psi_{l_2 L}^{lep})^c) + \text{h.c.}, \quad (36)$$

where the dimension  $M$  parameter  $\Lambda$  characterizes a scale of a beyond the SM physics ( $\Lambda \gg v$ ) and  $\tilde{\phi} = i\tau_2 \phi^*$  is the conjugated doublet. Let us stress that the Lagrangian (36) does not conserve the total lepton number.<sup>3</sup>

<sup>3</sup> The Lagrangian (36) can be generated (in the second order of the perturbation theory) by the seesaw interaction of the Higgs-lepton pair with a heavy Majorana right-handed lepton.

After spontaneous symmetry breaking from (36) we come to the Majorana mass term

$$\mathcal{L}^M = -\frac{1}{2} \frac{v^2}{\Lambda} \sum_{l_1, l_2} \bar{\nu}'_{l_1 L} Y_{l_1 l_2} (\nu'_{l_2 L})^c + \text{h.c.} = -\frac{1}{2} \sum_{i=1}^3 m_i \bar{\nu}_i \nu_i. \quad (37)$$

Here  $\nu_i = \nu_i^c$  is the field of the neutrino Majorana with the mass

$$m_i = \frac{v^2}{\Lambda} y_i = \frac{v}{\Lambda} (y_i v), \quad (38)$$

where  $y_i$  is a Yukawa coupling. In (38)  $y_i v$  is a "typical" fermion mass in SM. Thus, neutrino masses, generated by the effective Lagrangian (36), are suppressed with respect to "SM masses" by a factor

$$\frac{v}{\Lambda} = \frac{\text{scale of SM}}{\text{scale of a new physics}} \ll 1$$

The mechanism we have considered is, apparently, *the most economical and natural beyond the SM mechanism of the neutrino mass generation*. There are two general consequences of this mechanism.

- Neutrinos with definite masses  $\nu_i$  are Majorana particles.
- The number of neutrinos with definite masses is equal to the number of lepton-quark generations (three). This means that in this scheme there are no transitions of flavor neutrinos into sterile states.

The study of the lepton number violating neutrinoless double  $\beta$ -decay ( $0\nu\beta\beta$ -decay)

$$(A, Z) \rightarrow (A, Z + 2) + e^- + e^- \quad (39)$$

of some even-even nuclei is most sensitive way to investigate the Majorana nature of neutrinos with definite masses (see review [31]). The probability of the process (39) is proportional to square of the Majorana neutrino mass

$$m_{\beta\beta} = \sum_i U_{ei}^2 m_i \quad (40)$$

and is very small. It has the following general form

$$\frac{1}{T_{1/2}^{0\nu}} = |m_{\beta\beta}|^2 |M^{0\nu}|^2 G^{0\nu}(Q, Z). \quad (41)$$

Here  $M^{0\nu}$  is the nuclear matrix element and  $G^{0\nu}(Q, Z)$  is known phase factor.

Several experiments on the search for the  $0\nu\beta\beta$  of different nuclei are going on and are in preparation. Up to now the process was not observed. From the data of recent experiments EXO-200 [32], KamLAND-Zen [33] and GERDA [34] the following upper bounds were, correspondingly, obtained

$$|m_{\beta\beta}| < (1.9 - 4.5) \cdot 10^{-1} \text{ eV}, \quad (1.4 - 2.8) \cdot 10^{-1} \text{ eV}, \quad (2 - 4) \cdot 10^{-1} \text{ eV} \quad (42)$$

In future experiments on the search for  $0\nu\beta\beta$  decay the values  $|m_{\beta\beta}| \simeq \text{a few} \cdot 10^{-2} \text{ eV}$  are planned to be reached.

Indications in favor of transitions of flavor neutrinos into sterile states were obtained in the LSND [36] and MiniBooNE short baseline accelerator experiments and in the GALLEX and SAGE calibration experiments and in short baseline reactor experiments which were reanalyzed with a new reactor antineutrino flux (see recent review [35]). Many new short baseline source, reactor and accelerator neutrino experiments on the search for sterile neutrinos with masses  $\sim 1 \text{ eV}$  are in preparation (see [37]). There is no doubt that in a few years the sterile neutrino anomaly will be resolved.

## 7. Conclusion

Neutrino masses and mixing, discovered via the observation of neutrino oscillations, is a first particle physics evidence of a new beyond the SM physics. We discuss here briefly first proposals for neutrino oscillations and first steps in the development of the theory of neutrino oscillations. Then we consider basics of neutrino mixing and oscillations, convenient formalism for neutrino oscillations in vacuum and the definition of the atmospheric neutrino mass-squared difference. In the final part of the paper we discuss the important role which play two-component neutrino in the Standard Model and the most economical Weinberg mechanism of the generation of the Majorana neutrino masses.

## References

- [1] Fukuda Y *et al* [Super-Kamiokande Collaboration] 1999 *Phys. Rev. Lett.* **82** 2644
- [2] Ahmad Q *et al* [SNO Collaboration] 2002 *Phys. Rev. Lett.* **89** 011301
- [3] Eguchi K *et al* [KamLAND Collaboration] 2003 *Phys. Rev. Lett.* **90** 021802
- [4] Mariani C 2008 [K2K Collaboration] *AIP Conf. Proc.* **981**247
- [5] Timmons A 2015 *Preprint* arXiv:1511.06178 [hep-ex]
- [6] Abe K *et al* [T2K Collaboration] 2015 *Phys. Rev. D* **91** 7 072010
- [7] An F *et al* [Daya Bay Collaboration] 2015 *Phys. Rev. Lett.* **115** 11 111802 .
- [8] Kim S 2015 *Nucl. Part. Phys. Proc.* **265-266** 93
- [9] Abrahao T *et al* [Double Chooz Collaboration] 2015 *Preprint* arXiv:1510.08937 [hep-ex]
- [10] Landau L 1957 *Nucl. Phys.* **3** 127; Lee T D and Yang C N 1957 *Phys. Rev.* **105** 1671; Salam A 1957 *Nuovo Cim.* **5** 299
- [11] Goldhaber M, Grodzins L and Sunyar A 1958 *Phys. Rev.* **109** 1015
- [12] Pontecorvo B 1957 *Sov. Phys. JETP* **6** 429 [1957 *Zh. Eksp. Teor. Fiz.* **33** 549]
- [13] Pontecorvo B 1957 *Sov. Phys. JETP* **7** 172 [1957 *Zh. Eksp. Teor. Fiz.* **34** 247]
- [14] Maki Z, Nakagawa M and Sakata S 1962 *Prog. Theor. Phys.* **28** 870
- [15] Danby G, Gaillard J, Goulianos K, Lederman L, Mistry N, Schwartz M and Steinberger J 1962 *Phys. Rev. Lett.* **9** 36
- [16] Pontecorvo B 1968 *Sov. Phys. JETP* **26** 984 [1967 *Zh. Eksp. Teor. Fiz.* **53** 1717]
- [17] Gribov V and Pontecorvo B 1969 *Phys. Lett. B* **28** 493
- [18] Bilenky S and Pontecorvo B 1976 *Phys. Lett. B* **61** 248; Bilenky S and Pontecorvo B 1976 *Lett. Nuovo Cim.* **17** 569; Bilenky S and Pontecorvo B 1976 *Yad. Fiz.* **24** 603 [1976 *Sov. J. Nucl. Phys.* **24** 316]
- [19] Bilenky S and Pontecorvo B 1978 *Phys. Rept.* **41** 225
- [20] Bilenky S and Petcov S 1987 *Rev. Mod. Phys.* **59** 671
- [21] Bilenky S 2010 *Introduction to the physics of massive and mixed neutrinos* (Springer, Lecture Notes in Physics) vol 817
- [22] Minkowski P 1977 *Phys. Lett. B* **67** 421; Gell-Mann M, Ramond P and Slansky R 1979 in *Supergravity* p. 315 edited by van Nieuwenhuizen F and Freedman D, North Holland, Amsterdam; Yanagida Y 1979 Proc. of the *Workshop on Unified Theory and the Baryon Number of the Universe*, KEK, Japan; Glashow S 1979 *NATO Adv. Study Inst. Ser. B Phys.* **59** 687; Mohapatra R and Senjanović G 1981 *Phys. Rev. D* **23** 165
- [23] Bilenky S *Phys. Part. Nucl. Lett.* **12**, no 4 p 453 [arXiv:1502.06158 [hep-ph]].
- [24] Bilenky S, von Feilitzsch F and Potzel W J. *Phys. G* **38** (2011) 115002 [arXiv:1102.2770 [hep-ph]].
- [25] Giunti C and Kim C, "Fundamentals of Neutrino Physics and Astrophysics," Oxford, UK: Univ. Pr. (2007) 710 p.
- [26] Capozzi F, Fogli G, Lisi E, Marrone A, Montanino D and Palazzo A J. *Phys. Conf. Ser.* **598** (2015) 1, 012002.
- [27] Gonzalez-Garcia C, Maltoni M and Schwetz T *JHEP* **1411** (2014) 052 [ arXiv:1409.5439 [ hep-ph]]
- [28] Nunokawa H, Parke S and Zukanovich Funchal R *Phys. Rev. D* **72** (2005) 013009.
- [29] Weinberg S *Phys. Rev. Lett.* **43** (1979) 1566.
- [30] Bilenky S *Phys. Part. Nucl.* **46** (2015) no 4 p 475 [arXiv:1501.00232 [hep-ph]].
- [31] Bilenky S and Giunti C *Int. J. Mod. Phys. A* **30** (2015) no. 04n05 1530001 [arXiv:1411.4791[hep-ph]].
- [32] Albert J *et al* [EXO-200 Collaboration] *Nature* **510** (2014) 229 [arXiv:1402.6956 [nucl-ex]].
- [33] Asakura K *et al* [KamLAND-Zen Collaboration] *AIP Conf. Proc.* **1666** (2015) 170003[arXiv:1409.0077 [physics.ins-det]].
- [34] Wester T *AIP Conf. Proc.* **1686** (2015) 020026.
- [35] Gariazzo S, Giunti C, Laveder M, Li Y and Zavanin E arXiv:1507.08204 [hep-ph].

- [36] Aguilar-Arevalo A *et al* [LSND Collaboration] Phys. Rev. D **64** (2001) 112007 [hep-ex/0104049].
- [37] Lasserre T Phys. Dark Univ. **4**, 81 (2014), [arXiv:1404.7352 [hep-ex]].