

Time-Dependent Two-Particle Reduced Density Matrix Theory: Application to High-Harmonic Generation

Fabian Lackner^{*1}, Iva Březinová^{*2}, Takeshi Sato^{†‡}, Kenichi L. Ishikawa^{†‡}, Joachim Burgdörfer^{*}

^{*} Institute for Theoretical Physics, Vienna University of Technology, Wiedner Hauptstraße 8-10/136, 1040 Vienna, Austria, EU

[†] Photon Science Center, School of Engineering, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8656, Japan

[‡] Department of Nuclear Engineering and Management, School of Engineering, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8656, Japan

Synopsis We follow a new approach for calculating high-harmonic spectra for multi-electron atoms and molecules by propagating the two-particle reduced density matrix. Calculated results are in very good agreement with state-of-the-art many-body wavefunction-based benchmark calculations.

High-harmonic generation (HHG) is one of the fundamental processes in strong field physics whose applications range from attosecond metrology [1], tunable table-top XUV/Soft X-ray sources [2] to high precision spectroscopy [3] and orbital imaging [4]. On the atomic level the theoretical description of HHG is challenging because of the multi-electron nature of the underlying process. While simple models such as the single-active-electron approximation (SAE) or time-dependent Hartree-Fock (TDHF) are well suited to describe qualitative features of HHG, advanced theories capable of correctly treating electron correlation are needed for a quantitative description [5]. However, with increasing system size conventional wavefunction-based methods such as the multi-configurational time-dependent Hartree-Fock (MCTDHF) method soon become unfeasible due to their exponential scaling with particle number. One way to overcome this exponential barrier is to abandon a wavefunction-based description and propagate time-dependent reduced densities instead. Considerable success along this line has been achieved by time-dependent density functional theory (TDDFT) [6]. However, accurate calculations in TDDFT face the difficulty of unknown exchange-correlation functionals that are difficult to improve systematically. We propose to go beyond the limitations of TDDFT by propagating the time-dependent two-particle reduced density matrix (TD-2RDM). As a hybrid between the electron density and the many-body wavefunction the 2RDM fully includes two-particle correlations. We have implemented the TD-2RDM method to describe high-harmonic generation of fully three-dimensional multi-electron atoms [7]. To obtain accurate results for the electronic response we use an advanced closure scheme [8] that is suited to conserve all constants of motion associated with symmetries of the Hamiltonian. We benchmark the

performance of the TD-2RDM method by comparing it to a state of the art MCTDHF calculation [9] as well as to TDDFT calculations (see Fig.1). We find very good agreement between the TD-2RDM and the MCTDHF method while TDDFT within the local density approximation shows clear deviations indicating that the correct treatment of two-particle correlations is essential to obtain accurate HHG spectra.




Figure 1. HHG spectrum of beryllium subject to a 3-cycle laser pulse with intensity $I = 4.0 \times 10^{14} \text{ W/cm}^2$. MCTDHF(—), TD-2RDM(---), TDDFT(⋯).

References

- [1] M. Hentschel *et al.* 2001 *Nature* **414** 509
- [2] Ph. Zeitoun *et al.* 2004 *Nature* **431** 426
- [3] A. D. Shiner *et al.* 2001 *Nature Phys.* **7** 464
- [4] J. Itatani *et al.* 2004 *Nature* **432** 867
- [5] S. Sukiasyan *et al.* 2009 *Phys. Rev. Lett.* **102** 223002
- [6] X. M. Tong *et al.* 2015 *Phys. Rev. A* **92** 043422
- [7] F. Lackner *et al.* *Phys. Rev. A* in press
- [8] F. Lackner *et al.* 2015 *Phys. Rev. A* **91** 023412
- [9] T. Sato *et al.* 2016 *Phys. Rev. A* **94** 023405

¹E-mail: fabian.lackner@tuwien.ac.at

²E-mail: iva.brezinova@tuwien.ac.at

