

Article

Field Measurements and Satellite Remote Sensing of Daily Soil Surface Temperature Variations in the Lower Colorado Desert of California

Dana Coppernoll^{1,2} and Christopher Potter^{1*}¹ NASA Ames Research Center, Moffett Field, CA² Thomas Jefferson High School, Federal Way, WA

* Correspondence: chris.potter@nasa.gov; Tel.: +1 650 504 5164

Received: date; Accepted: date; Published: date

Abstract: The purpose of this study was to better understand the relationships between diurnal variations of air temperatures measured hourly at the soil surface, compared with the thermal infrared (TIR) emission properties of soil surfaces located in the Lower Colorado Desert of California, eastern Riverside County. Fifty air temperature loggers were deployed in January of 2017 on wooden stakes that were driven into the sandy or rocky desert soils at both Ford Dry Lake and the southern McCoy Mountains wash. Land Surface Temperature (LST) derived from Landsat satellite images was compared to measured air temperatures at 1 m and at the soil surface on 14 separate dates, until mid-September, 2017. Results showed that it is feasible to accurately derive estimate temperatures at the soil surface from hourly air temperatures recorded at 1 m above the surface. The study further correlated Landsat LST closely with site measurements of air and surface temperatures in these solar energy development zones of southern California, allowing interconversion with ground-based measurements for use in ecosystem change and animal population biology studies.

Keywords: Lower Colorado desert, surface temperature, thermal infrared, Landsat

1. Introduction

Diurnal and seasonal variations in land surface temperature can influence numerous ecological and biogeochemical processes in desert landscapes, including wildlife population dynamics, plant germination, microbial activity, evapotranspiration, and trace gas emissions. The effect of extreme hourly temperature variations on desert ecosystems range from limited photosynthesis rates and evapotranspiration fluxes, to altered microbial symbiotic relationships through nitrogen fixing bacteria in root nodules.

It is well documented that surface soil temperatures are a function of the downwelling electromagnetic radiation and of the albedo (reflectivity) of a soil surface. When energy from the sun encounters any matter, whether it is the atmosphere or the ground surface, it will go through the processes of transmission, reflection, and absorption. Solar irradiance that is reflected is mostly due to differing albedo levels across the Earth's surface from highly reflective ice and clouds to non-

reflective oceans and dark rocks. Slope, vegetative cover, water content, bulk density, and a host of other factors affect how much heat the soil absorbs or re-emits.

Variations in temperature at the soil surface has been shown to influence the survival and distribution of several noteworthy animal species in the deserts of southern California. Most, if not all, desert animals change their behavior based on outside temperatures. In summer, temperatures on Mojave Desert valley floors can exceed 50°C, much hotter than is generally tolerable by many animals. Consequently, desert animals are commonly more active at dawn and dusk, may operate entirely nocturnally, and take shelter in burrows or other cooler areas during the heat of the day.

Temperature regulation is important for Desert Tortoises (*Gopherus agassizii*), a federally protected species native to the Southwest United States. Tortoises have been noted to survive temperatures below freezing and above 40°C, however, most activity occurs between 26.1°C to 33.9°C. Temperature also affects incubation time, sex differentiation, and survival in Desert Tortoises, with higher temperatures indicating higher survival and shorter incubation periods, and sex differentiation occurring somewhere between 31°C–32°C. It is known that more frequent and prolonged droughts due to climate change can further decrease the habitable areas for tortoises, and some populations are already showing signs of density decreases due to climate change.

Fringe-Toed Lizards are also influenced strongly by air temperature. Optimal body temperature for these reptiles is around 37°C, and most lizards will seek out burrows or other shelter when temperatures exceed 48.9°C. As climate change and human interactions further affect and reduce these animals' habitats, identifying time periods and specific locations of potentially lethal temperatures for desert animals can be key to protecting these species.

In general, it would be advantageous for resource managers to be able to accurately infer variations of temperatures at the soil surface from hourly air temperatures recorded at 1–2 m above the surface (ambient), since such ambient air temperatures are measured commonly at weather station locations and have been maintained over relatively long periods of time. Although there have been numerous models developed to predict temperature at a few centimeters depth into the soil profile from ambient air temperatures, 10–16°C air temperatures at the soil surface, in the habitat zone within which desert animals must survive or traverse, have not been measured frequently in combination with ambient air temperatures across the desert landscape.

Moreover, it would be valuable to be able to develop correlations between measured temperatures at the soil surface (or at weather stations) to satellite remote sensing, which can provide surface emission temperatures in otherwise inaccessible remote locations. Areas that have few or no weather stations can often be difficult to monitor. Satellites offer an affordable method to monitor day-to-day temperature fluctuations across regions at a relatively high spatial resolution of 100 m. Nonetheless, it is not well documented where and when Land Surface Temperature (LST) derived from a satellite images diverge from measured surface or ambient air temperatures.

Previous research by Li et. al. (2017) showed a correlation between air temperature and LST from the Advanced Very High Resolution Radiometer (AVHRR) instrument on the National Oceanic and Atmospheric Administration (NOAA) polar-orbiting satellites with an R^2 value of 0.54, 0.42 and 0.62 over three years. Despite large differences in air temperature and LST values, it was shown that a linear model can be developed to enhance accuracy. Kawashima et. al. (2018) found strong correlations with R^2 values from 0.75 to 0.85 between air temperature measured by Automated Meteorological Data Acquisition System (AMeDAS) and LST measured by the Landsat Thematic

Mapper on winter nights in Japan. Liang and Chen [19] however, found no statistically significant correlations between surface air temperature from weather stations and Landsat TIR thermal infrared bands in the fall and spring in China.

The objective of this study was to quantify the relationship between air temperatures measured at the soil surface, ambient air temperatures measured at 1 m height, and Landsat surface temperature measurements over the same area and period of time in the Lower Colorado Desert of eastern Riverside County, California. We report on new algorithms and their levels of certainty to scale from satellite images down to the typical size of field plot for animal and plant research for surface temperature monitoring on hourly to daily time intervals.

2. Study Area

Field measurements of air and soil surface temperatures for this study were carried out in eastern Riverside County in the Lower Colorado Desert of California (Figure 1). This area is bounded on the west by the Laguna, Santa Rosa, and San Jacinto mountain ranges, on the east by the California–Arizona state line, on the north by the gradual transition to the Mojave Desert, and on the south by the California–Mexico border [20]. Low annual rainfall (50–300 mm) and high temperatures exceeding 45 °C in the summer make this area one of the most arid in North America.

Ford Dry Lake is the lowest point in Chuckwalla Valley of Riverside County, at 110 m above sea level. A Soil Climate Analysis Network (SCAN) weather station operated by the U.S. Department of Agriculture is located at Ford Dry Lake (33° 39' N, 115° 6' W) and was used to provide a climatology record for the study area (data available at www.wcc.nrcs.usda.gov/scan/). The southern McCoy Mountains are located 5–10 km to the east of Ford Dry Lake.

Two relatively new solar energy production facilities have been built in both Ford Dry Lake and in the southern McCoy Mountains washes. The Genesis Solar Energy Project completed in 2014 near Ford Dry Lake consists of two independent solar electric generating facilities with a nominal net electrical output of 125 megawatts (MW) each. Electrical power is produced using steam turbine generators fed from solar trough generators. The McCoy I Solar Energy Project is a 750-MW flat-panel photovoltaic facility completed in 2016 in the southern McCoy Mountain washes (locations of both facilities are shown in Figure 1).

The study sites are in the American Semi-Desert and Desert Province ecoregion, Sonoran Desert Subsection [21,22]. This part of the Sonoran Desert, in the Lower Colorado River basin, hosts many species of plants and animals. The main perennial vegetation cover type is creosote bush (*Larrea divaricata*) and white bursage (*Ambrosia deltoidea*), although ironwood (*Olneya tesota*), palo verde (*Cercidium floridum*) and ocotillo (*Fouquieria splendens*) are also found in the area.

Predators such as mountain lions (*Felis concolor*) and coyotes (*Canis latrans*) feed on desert bighorn sheep (*Ovis canadensis*), black-tailed jackrabbits (*Lepus californicus*) and round-tailed ground squirrels (*Spermophilus tereticaudus*). California leaf-nosed bats (*Macrotus californicus*) and ring-tailed cat (*Bassaris astutus*) also survive in the harsh environment. The Sonoran Desert is known for its great diversity in bird species, a few of which range into the Lower Colorado area: phainopepla (*Phainopepla nitens*), Costa's hummingbird (*Calypte costae*) and roadrunner (*Geococcyx californianus*). Many species of reptiles can be found, most notably the desert tortoise (*Gopherus agassizii*) and chuckwalla lizard (*Sauromalus ater*) [23].




Figure 1. Transect locations of temperature measurement stakes (black dots within circles) in eastern Riverside County, California. The southern McCoy Mountains are to the right center on the true-color satellite image, while Ford Dry Lake is on the westernmost section of the study area map.

3. Materials and Methods

3.1. Field Measurements of Air Temperature

A total of 50 pendant air temperature loggers were deployed in January of 2017, and attached to wooden stakes that were driven into the sandy or rocky desert soils at both Ford Dry Lake and the southern McCoy Mountains washes. Each wooden stake was spaced along a predominantly east-west transect, 100 m apart from the previous stake location (Figure 1). Two sensors were attached to each stake by plastic zip ties, one at 1 m height and the other resting directly on the soil substrate surface. The 1.0 m measurement height was used to approximate an average weather sampling height with free air circulation around the sensor, a height which can vary widely from weather station network to network. The sensors were not shielded from incoming solar radiation, so as to expose the temperature readings to the effects of direct sunlight and to mimic extreme surface heating conditions.

The locations of stakes ranged in elevation from 110 to 140 m above sea level. The western group of measurement locations was placed in sandy soils typical of the location of Genesis solar facility and the eastern group of measurement locations was placed in more rocky soils typical of the McCoy I solar facility landforms.

Pendant Temperature 8K Data Loggers were rated to measure temperatures from 20°C to 70°C (68°F to 158°F). Reported accuracy is $\pm 0.53^\circ\text{C}$, over the range of 0°C to 50°C ($\pm 0.95^\circ\text{F}$ from 32°F to 122°F). Resolution of the sensors is 0.14°C at 25°C (0.25°F at 77°F) with drift being less than $0.1^\circ\text{C}/\text{year}$ ($0.2^\circ\text{F}/\text{year}$). All temperature loggers were collected from the field site one year later (January, 2018) for data download and analysis.

3.2. Landsat Thermal Infra-Red Imaging

The Thermal Infrared Sensor (TIRS) is an instrument on the Landsat 8 satellite that collects images within the thermal range 10–12.5 μm . Landsat 8 Images are collected every 16 days, with each scene being 170 km north-south by 183 km east-west. The TIRS has a pixel size of 100 m which is resampled to 30 meters to match Landsat multispectral bands. Data were gathered from the Landsat 8 TIRS sensor for fourteen different flyover dates in 2017, all at approximately 10:16 local time (Table 1).

Prior to user download, calibration coefficients have been applied to Landsat digital numbers to derive top of the atmosphere (TOA) reflectance components using per-pixel solar angles derived from and 4 closest to center of focal plane. Brightness Temperature (BT) was derived from TOA radiance and two thermal constants as described online at landsat.usgs.gov/using-landsat-8-product. The BT of any 100 m resolution Landsat pixel without further atmospheric correction can be considered an approximation of kinetic LST in degrees Kelvin.

Table 1. Dates in 2017 for which Landsat scenes could be acquired. All days were verified to have minimal cloud cover over the ground temperature measurement locations.

Date	Rainfall events recorded at Ford Dry Lake SCAN site
1/18/2017	
2/3/2017	
2/19/2017	2/18/17 1 day before
3/7/2017	
3/23/2017	
5/10/2017	5/9/17 1 day before
5/26/2017	
6/11/2017	
6/27/2017	
7/13/17	
7/29/17	
8/14/17	8/10/17 4 days before
8/30/17	
9/15/17	9/8/17 7 days before

3.3. Statistical Analysis

Daily LST values from Landsat 8 images were compared to both ground surface temperature and air temperature readings from the sensors, using the mean difference (MD) standard deviation (SD) and linear least-squares regression correlation.

4. Results

4.1. Climatology

At the Ford Dry Lake SCAN station, diurnal temperature ranges (DTR) varied significantly over the year of 2017, with winter generally having the lowest DTR and summer having the highest DTR. The annual average DTR was 15.6 °C; the maximum was 25.5 °C on October 8th, and the minimum was 5.6 °C on both January 13th and February 18th. The hottest day occurred on June 20th, with ambient air temperature of 50.6 °C. The coldest day occurred on both December 23rd and 26th, with air temperature of 1.7 °C. During the 2017 measurement period, there were 15 total days in which measurable precipitation was recorded at the station, ranging from 0.3 to 24 mm (0.9 inches). Of those, five events showed rainfall totals of 5 mm or greater: one in February, one in May, two in August, and one in September.




Figure 2. Climatology from Ford Dry Lake SCAN station in 2017

Difference between surface and air temperature

The hourly ground surface and the ambient 1 m air temperature at each site were found to be closely correlated across the measurement locations (Table 2). Linear regression correlation values (R^2) between hourly T_{surface} and T_{air} measurements varied from 0.86 to 0.99. Throughout the year, the hourly mean difference (MD) between T_{surface} and T_{air} was consistently greater than zero. An estimated conversion factor using the average hourly MD was determined to be 5.8 °C, meaning that to convert ambient air temperature to ground surface air temperature, one should add 5.8 °C to air temperature measured at 1 m height.

Table 2. Mean difference (MD), standard deviation (SD), and linear regression correlation (R^2) between hourly T_{surface} and T_{air} at measurement locations over all days. Measurement locations 2, 5, 7, 11, 12, 17, 20, and 24 were unable to be used due to sensor malfunctions and missing data values.

Location No.	MD	SD	R^2
1	7.5	5.9	0.95
3	4.8	4.8	0.95
4	6.3	3.9	0.94

6	3.4	5.4	0.91
8	1.3	6.9	0.88
9	0.1	7.3	0.86
10	1.2	1.9	0.99
13	10.9	4.5	0.96
14	5.4	2.5	0.97
15	12.0	5.2	0.96
16	8.4	5.6	0.97
18	1.6	2.5	0.96
19	2.3	1.9	0.99
21	7.4	3.1	0.99
22	2.4	2.2	0.98
23	10.9	3.6	0.95
25	3.2	7.2	0.89
26	10.3	3.5	0.97
27	9.3	4.8	0.97
28	9.0	4.5	0.91

During summer months (June–Sep), we measured a larger disparity between surface and air temperatures at each measurement location than during winter months (Jan–May). For instance, the average difference for Jun/Jul between T_{surface} and T_{air} for all measurement locations was 7.5° C, compared to an Jan/Feb average difference of 2.4° C as illustrated in Figure 3.




Figure 3. Diurnal T_{surface} and T_{air} , for a selected measurement location for all available dates of Landsat image acquisition.

Comparisons between T_{air} and Landsat LST

Hourly T_{air} and the corresponding LST from Landsat at approximately 10:16 AM for each measurement location were closely correlated over the course of 2017 (Table 3). Linear regression R^2 values between hourly LST and T_{air} measurements varied from 0.83 to 0.93. The mean difference (MD) between T_{air} and LST was typically greater than zero, although the MDs for T_{air} and LST were smaller than the MDs measured between T_{surface} and T_{air} , comparing location by location from Table 2. The averaged conversion factor from the MDs for all measurement locations was 0.78°C, meaning that to convert T_{air} to Landsat LST, one would subtract 0.78°C from a measured T_{air} value.

An example scatterplot between T_{air} and LST values from 2017 Landsat imagery shown in Figure 4 demonstrated the strong linear relationship that was maintained from winter to summer months. A 1:1 ratio of actual measured T_{air} and LST (Figure 5) was observed most commonly during the months of March and July (Figure 4).

Table 3. Mean difference (MD), standard deviation (SD), and linear regression correlation (R^2) between measured T_{air} and LST using all dates of Landsat image acquisition.

Location No.	MD	SD	R^2
1	1.4	5.2	0.93
3	4.9	5.2	0.92
4	2.1	6.2	0.88
6	3.1	6.5	0.87
8	1.5	6.9	0.87
9	1.1	6.2	0.90
10	2.8	7.9	0.83
13	1.6	5.4	0.90
14	1.9	5.0	0.92
15	1.2	4.5	0.93
16	1.0	5.3	0.91
18	1.4	5.5	0.91
19	1.9	4.9	0.93
21	0.4	4.9	0.92
22	2.1	5.0	0.91
23	3.3	5.3	0.90
25	7.5	7.7	0.86
26	1.7	5.5	0.90
27	1.8	4.8	0.93
28	0.4	6.8	0.84




Figure 4. Scatter plot between T_{air} , and 2017 Landsat LST image data for measurement location 1 in Ford Dry Lake.




Figure 5. Slope and R^2 for the linear best-fit line between hourly T_{air} measurements and LST from Landsat imagery. The “best-fit” 1:1 ratio level is shown as a dashed horizontal line.

Comparisons between all three measurements

Across all measurement locations in the study area, hourly daily $T_{surface}$ was commonly higher than either maximum daily T_{air} , or LST (Figures 3 and 6) although there were a few exceptions to that pattern. Most notably, $T_{surface}$ was lower than T_{air} for 20% of the hourly observations collected in

2017, mainly during colder months (winter and autumn) and T_{surface} was lower than LST for 12 of hourly observations from summer and autumn Landsat flyover dates.




Figure 6. Comparison of T_{surface} , T_{air} , and LST at the hour of Landsat flyover for a selected measurement location.

5. Discussion

We observed a strong relationship between Landsat LST and air temperatures in the Lower Colorado Desert, allowing for satellite-derived estimates of surface temperatures to be interpreted with high confidence in remote locations where it is not possible or feasible to maintain weather stations. With respect to the results from continuous hourly air temperature measurements in 2017, the slightly better correlation of T_{air} than T_{surface} with LST may have been the result of freer air circulation at 1 m height than right at the ground surface, which would have explained the closer match of T_{air} with the 100 m resolution of Landsat TIR emission of heat from the land surfaces.

Desert animals survive and live in a variety of temperatures, but there are frequent instances of LST exceeding 50°C in Landsat image data (see example Figure 7) an extreme that would generally drive even the hardiest of animals underground or to shelter. There are several noteworthy features of LST pattern in Figure 7. First, the highest summer surface temperatures appear on the predominantly south-facing slopes at the rocky bases and lower washes of the Valen and McCoy Mountains, whereas the coolest summer surface temperatures appear on the predominantly north-facing slopes at the highest elevations of these mountains. Second, the solar energy absorbing facilities at Genesis and McCoy I sites remained several degrees Celsius cooler than the surrounding sandy washes of Ford Dry Lake and the southern McCoy Mountains.




Figure 7. June 2016 Landsat LST grayscale image, centered on the Valen and McCoy Wilderness Area in eastern Riverside County and the solar energy development zones shown in white outlines. LST ranges from 40°C in black shades to 50°C in light gray shades.

Being able to interpret near real-time spatial LST data for the whole of the Mojave and Lower Colorado Desert region can allow researchers to pinpoint areas that have shown a higher frequency of these critical temperature limits than others. This information could help guide new field surveys

and fine-tune existing conservation and habitat restoration plans. Desert Tortoise and Fringe-Toed Lizards are found throughout the Lower Colorado Desert, but specific habitat locations that can be favored by species—for example, alluvial washes or rocky hills—which can be used to narrow down optimal locations to monitor using LST images. Areas that have established populations of wildlife, like Joshua Tree National Park, are prime sites to monitor known breeding habitats and to better understand the stresses present for these animals in the ecosystem.

Dotter (2016) mapped out extensive coverages of well-developed desert rock pavements from Landsat imagery within solar energy development boundaries in eastern Riverside County, and specifically in the McCoy Mountain washes. Rock pavement surfaces display spectra that are strongly modified by the reflectance of surface varnish. Such varnishes decrease surface reflectance and albedo, and result in a steep positive slope in between Landsat bands 1 and 5, owing to a strong suppression of short wavelength radiance by the presence of oxidized metals—mainly Fe and Mg. In contrast, active washes that are more commonly dominated by sand—quartz and feldspar crystals—result in an increased reflectance—higher albedo—in all Landsat bands (27). Despite these detectable differences in reflectance properties between sandy and rocky desert soils, the hourly temperature records we collected in 2017 across the varied geomorphic surfaces of the study area were not statistically significant between measurement locations at any given time of the year. It may be the case that the consistently high mid-day solar irradiance in the Lower Colorado Desert dominates over any and all differences in soil albedo in determining surface air temperatures and DTR on a year-round basis.

6. Conclusions

Our understanding of how climate change and human activity together affect the natural world is presently limited in part by our ability to measure change at all necessary scales. Unlike physical monitoring stations, which are costly to install and maintain, satellite measurements can be potentially acquired every week at a relatively low cost to provide critical habitat information on regional surface conditions. This study has provided strong evidence that Landsat LST correlates closely with on-site measurements of air and surface temperatures in the deserts of southern California, and can be accurately interconverted with those ground-based measurements for use in ecosystem change studies.

Author Contributions: All authors have contributed equally to conceptualization, analysis, writing, and investigations.

Funding: This research was funded by the U.S. Department of the Interior Bureau of Land Management (BLM).

Acknowledgments: The authors wish to thank Mark Massar and James Seigand of the BLM Field Office Palm Springs for assistance in field sampling.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Paul, E. A. and Clark, F. C., 1996, *Soil Microbiology and Biochemistry*, Academic Press, pp. 12–32.

2. Sims, K. and Dunigan, J., 1984, Diurnal and seasonal variations in nitrogenase activity N_2 reduction of rice roots. *Soil Biology and Biochemistry*. 16:15–18.
3. Billel, D., 1998, *Environmental Soil Physics*, Academic Press pp. 340.
4. Lehnert, M., 2015, Factors affecting soil temperature as limits of spatial interpretation and simulation of soil temperature. *Acta Universitatis Palackianae Olomucensis . Geographica*. 45. 5–21.
5. Bai, J., T. A. Scott, J. Chen, R. A. Minnich, and A. C. Chang, 2011, Long-term variation in soil temperature of the Mojave Desert, southwestern USA, *Climate Research*, 46: 43–50.
6. Dudek ICF International 2012. Mojave Fringe-Toed Lizard (*Uma scoparia*). Retrieved from <https://www.drecp.org/documents/docs/baselinebiologyreport/10AppendixSpeciesProfiles/10aReptileAmphibian/MojaveFringetoedLizard.pdf>
7. Meyer, R. 2008. *Gopherus agassizii*. In: Fire Effects Information System, online. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory Producer. Available at: www.fs.fed.us/database/feis/animals/reptile/goag/all.html 2018, August 1.
8. Lewis-Inokur, V. and Inokur, R. M., 1995, Incubation temperature affects sexual differentiation, incubation time, and posthatching survival in desert tortoises *Gopherus agassizii*. *Canadian Journal of Zoology*, 73:11–2091–2097, <https://doi.org/10.1139/z95-246>
9. Lovich, J., Mackulic, C. J., Freilich, J., Agha, M., Austin, M., Meyer, K. J., Arundel, T. R., Hansen, T., Vamstad, M. S., Root, S. A. 2014. Climatic variation and tortoise survival: Was a desert species met its match? *Biological Conservation*, 169:214–224
10. Masfurther VR, Furman RD 1974. Soil temperature modeling using air temperature as a driving mechanism. *Trans ASAE*, 16:78 – 81.
11. Persaud N, Chang AC, 1983, Estimating soil temperature by linear filtering of measured air temperature. *Soil Sci Soc Am J* 47:841–847.
12. Pockock KL, Jeffers NR, Lindley DK, Adamson K, Hill CA 1977. Estimating woodland soil temperatures from air temperature and other climatic variables. *Agric. Meteorol.* 18:351– 372.
13. Toy T. J., Kuhaida A. J., Munson J. J., 1978. The prediction of mean monthly soil temperature from mean monthly air temperature. *Soil Sci.*, 126:181–189.
14. Mahrer J., 1980, A numerical model for calculating the soil temperature regime under transparent polyethylene mulches. *Agric. Meteorol.*, 22:227–234.
15. Cheng D., Hunt J. R., Running S. J. 1993. A daily soil temperature model based on air temperature and precipitation for continental applications. *Clim. Res.*, 2:183 –191.
16. Kang S, Kim S, H S, Lee D 2000. Predicting spatial and temporal patterns of soil temperature based on topography, surface cover and air temperature. *Forest Ecol. Manag.*, 136: 173–184.
17. Li, J., Luo, J., Dixon, J., and Ye, J., 2008, Applicability of Land Surface Temperature LST estimates from AVHRR satellite image composites in northern Canada. *Prairie Perspectives*. 11:119–130.
18. Kawashima, S., T. Ishida, M. Minomura, and T. Miwa, 2000: Relations between Surface Temperature and Air Temperature on a Local Scale during Winter Nights. *J. Appl. Meteor.*, 39, 1570–1579, [https://doi.org/10.1175/1520-0450\(2000\)39:1570:RSTAA\(2.0.CO;2](https://doi.org/10.1175/1520-0450(2000)39:1570:RSTAA(2.0.CO;2)
19. Xiong, J., and Chen, F. 2016. Relationship between temperatures from Landsat thermal infrared band retrievals and synchronous weather measurements. *2016 4th International Workshop on Earth Observation and Remote Sensing Applications (EORSA)*, 437–441.
20. Marks, J. 1950. Vegetation and soil relations in the Lower Colorado Desert. *Ecology*, 31, 176–193.
21. Bailey, R. J., 1994. Ecological classification for the United States. Washington, DC: USDA Forest Service.

22. Cleland, D.T., Freeouf, A., Keys, J., Nowacki, J., Carpenter, C.A., McNab, J. 2007. Ecological Subregions: Sections and Subsections for the conterminous United States. Retrieved from: <https://www.fs.usda.gov/treeearch/pubs/48672>.
23. Lovich, J. and J. R. 2011. Wildlife conservation and solar energy development in the Desert Southwest, United States. *Bioscience*, 61, 982–992.
24. U.S. Geological Survey 2018, Product Guide Landsat 8 Surface Reflectance Code LASRC Product, Department of the Interior,
25. J. J. Schott, R. J. Cook, S. J. Raoueno, N. J. Markham, J. L. Radocinski, R. J., 2014, Landsat 8 Thermal Infrared Sensor TIRS Vicarious Radiometric Calibration. *Remote Sens.* 6, 11607–11626.
26. Potter, C. 2016, Mapping changes in desert pavement surfaces of the Lower Colorado Desert of southern California using Landsat time series analysis, *International Journal of Advanced Remote Sensing and GIS*, 5: 1747–1754
27. J. K. K. and Anderson, R., 1998, The use of Landsat Thematic Mapper data for mapping and correlation of Quaternary geomorphic surfaces in the Southern Sierran Mountains, California, *International Journal of Remote Sensing*, 19 2345–2359.



© 2018 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license <http://creativecommons.org/licenses/by/4.0/>