



REVIEW ARTICLE



An overview of carbonic anhydrases and membrane channels of synoviocytes in inflamed joints

Min Jeong Ji  and Jeong Hee Hong 

Department of Physiology, College of Medicine, Gachon University, Lee Gil Ya Cancer and Diabetes Institute, Incheon, South Korea

ABSTRACT

The highly aggressive fibroblast-like synoviocytes (FLSs) are inflammatory mediators involved in synovial joint destruction. Membrane channels and transporters are essential components of the cell migration apparatus and are involved in various cellular functions. Although evidence is emerging that cell migration is a physiological/pathological process, the mechanism of highly dynamic synoviocytes linked to the membrane channels and carbonic anhydrases (CAs) in inflamed joints is only partially understood. In this review, topics covered will give a brief overview of CAs and the membrane channels of synoviocytes. We have also systematically focused on the role of FLS channels and transporters under various conditions, including rheumatoid arthritis (RA), to understand the pathophysiology of the migration of synoviocytes as inflammatory mediators in joints.

ARTICLE HISTORY

Received 4 July 2019
Revised 8 August 2019
Accepted 19 August 2019

KEYWORDS

Synoviocytes; migration;
carbonic anhydrases;
aquaporins; ion channels

1. Introduction

Rheumatoid arthritis (RA) is a common inflammatory autoimmune disease that induces diarthrodial joint inflammation¹. The fibroblast-like synoviocytes (FLSs), located in the synovium, mediate synovial joint destruction by releasing metalloproteinases (MMPs) and secreting cytokines, including interleukin (IL)-6, IL-1 β , IL-8, and tumour necrosis factor (TNF)- α in RA^{2–6}. Immune cells including macrophages, T cells, B cells, mast cells, and etc. are activated in RA and play crucial roles to secrete various cytokines and mediate inflammation of joint⁷. Moreover, TGF- β and platelet-derived growth factor (PDGF) levels were elevated in the RA synovial fluids^{8,9}. The inflamed synovium activates local FLS and induces the invasion of FLS¹⁰. Figure 1 represents the inflammatory mediators including immune cells and cytokines in pathogenesis of RA.

Cellular migration is not only an important physiological process related to wound healing, immune defence, and angiogenesis but also affects pathological processes, including tumour metastases and arthritis¹¹. Moreover, the maintenance of a polarised state is the basis for cell migration¹². Although the morphological polarisation of neuronal cells and epithelial cells is more critical than that of migrating cells, several mechanisms of the polarisation among classical polarised cells, such as neuronal and epithelial cells, and migrating cells follow similar principles. FLS are highly dynamic, and the chemotactic ability of the inflamed synovial fluid provides direction to the migration, invasion, and inflammatory/proliferative signalling events of pannus formation in the joint. To our knowledge, the membrane channels of FLS in migration have not been elucidated systematically. There is potential evidence for the involvement of carbonic anhydrases (CAs) and membrane channels in FLS migration in inflamed joints and has been discussed in the following sections.

2. Features of FLS

Normal synovial tissue is divided into two functional layers: a surface layer and sublining layer¹³. The surface layer is in direct contact with the intra-articular cavity, 1–2 layers deep consisting of two predominating cell types: macrophage-like synoviocytes (type A, MLS) derived from bone marrow, and FLS descended from the mesenchymal stem cells (type B)^{10,13}. Those two cell types are essential for maintaining joint homeostasis. The FLS is relatively richer than MLS and displays a variety of surface adhesion molecules, such as ICAM-1, CD90, and matrix proteins to help in the anchoring with extracellular matrix^{14,15}. The FLS contains endocrine and sensory functions and also possesses the epithelium-like nature¹⁶. While the typical feature of FLS is synovial hyperplasia during the inflammation, this review is designed to understand the dynamic role of FLS in the pathogenesis of RA and its associated membrane channels.

3. Carbonic anhydrases

Carbonic anhydrases are zinc metalloenzymes. The physiological role of CAs is related to membrane transporters and will be discussed in brief. CAs catalyse the reversible reactions of CO₂ and water to produce H⁺ and HCO₃[–]. They play a prominent role in the transport of CO₂ and protons across biological membranes, such as intercellular, intracellular, and extracellular spaces and are involved in diverse physiological functions, including pH regulation, fluids, and enzymes secretion and bone resorption^{17–19}. Several CA isozymes are expressed in mammals²⁰. Architecturally, CAs include cytosolic isoenzymes (CA I, CA II, CA III, CA VII, and CA VIII), membrane-associated isozymes (CA IV, CA IX, CA XII, and CA XIV), mitochondrial isoenzymes (CA VA and CA VB), and secreted CA isoenzyme (CA VI)²¹. CAs produce HCO₃[–], which fuels




Figure 1. Inflammatory mediators in pathogenesis of RA. Synovial fluid interacts with inflammatory cytokines and immune cells, which have role in inflammation and joint destruction in RA synovium. Inflammation and hyperplasia of FLS involve in pannus formation of joint.

the HCO_3^- transporter²². The $\text{Na}^+/\text{HCO}_3^-$ cotransporter, NBC1 cooperates with CA II and CA IV to modulate the intracellular pH²³. High H^+ concentration is found in the synovial fluid of RA patients, indicating that acidic pH reflects the pathophysiology of inflammation²⁴. CA and HCO_3^- -modulating transporters contribute to the modulation of synovial pH. The intensity of the inflammation process and ache-related symptoms in RA-affected patients inversely correlate with the tissue pH values^{24,25}. Tissue acidosis was found to be unfavourable for the progression of both antibody-mediated and cellular immunity processes²⁶. Although evidence indicates the importance of pH regulation of synovial tissues in various conditions, the precise role of the pH regulatory mechanism and its related transporters needs to be clarified. Beyond pH regulation, substantial evidence has indicated that CAs are also involved in bone resorption, hypoxia, and autoantibody formation. Expression of CA I was increased in the synovium of spondylitis and may accelerate calcification and bone resorption²⁷. The overexpression of CA IX and XII, associated with tumour hypoxia, was revealed in the inflamed synovium^{28,29}. Enhanced oxidative stress of erythrocytes in RA has addressed its correlation with CA II autoantibody formation³⁰.

Despite these differences in the role of CA, the regulatory role of CA in transporter machinery involves cell migration coordinated with cell adhesion molecules and ion transporters. The function of the CAs was to acidify the extracellular environment, thereby reducing cell adhesion and consequently increasing invasion and migration of tumour cells³¹. Especially, CA IX and CA XII were enhanced by hypoxic condition in tumour cells³². Hypoxia-inducible factor (HIF) affected the migration, cellular pH, and cell survival associated with tumour growth³³. The CA IX has been linked to cell–cell connections in the cell membrane, controlled by E-cadherin³⁴. It also regulated the cell migration by inhibiting E-cadherin associated with cell adhesion and interacting with the bicarbonate transporter, anion exchanger 2, in the leading edge regions in SiHa cells³¹. Bicarbonate transporters not only controlled the pH of the cells but also affected cell migration³⁵. The deficiency of SLC4A4 (NBCe1), an electrogenic $\text{Na}^+/\text{HCO}_3^-$ co-transporter, was influenced by cell migration by interfering with

the intracellular pH regulatory mechanism in MDA-MB 231 breast cancer cells³⁶. However, information on CAs on the RA FLS remains unclear. The verification of regulatory and migration role of CAs in FLS will provide a new scope for synovial physiology.

4. Membrane ion channels of FLS

4.1. Aquaporins

The aquaporins (AQPs) are water or small molecule-transporting channel proteins across the plasma membranes of various human tissues and cell types³⁷. Thirteen types of AQPs (AQP0–AQP12) from mammalian tissues have been cloned and sequenced³⁸. The AQPs are classified into two groups: water selective channel (orthodox AQPs) and water, glycerol, nitrate (AQP6), and urea channel (aquaglyceroporins; AQP 3, AQP7, and AQP9)³⁹. The permeability of AQPs is dependent on osmotic and hydrostatic gradients and pH values. Several investigations have shown the involvement of AQPs in cartilage damage in joint diseases like RA and osteoarthritis (OA). AQP1 is distributed in the articular cartilage and the synovium⁴⁰. AQP1 is also expressed in chondrocytes and synoviocytes of RA patients⁴¹. Up-regulated AQP1 found in the inflamed synovial tissues of RA patients might play a potential pathological role in hyarthrosis and joint swelling⁴². Acetazolamide, AQP1 inhibitor, was decreased AQP1 protein level via inhibition of NF- κ B activation and subsequent reduction of hind paw swelling in adjuvant-induced arthritis rats, suggesting that attenuation of AQP1 mediates anti-arthritis effect⁴². It is well-known that AQP4 possesses high water permeability than that of AQP1⁴³ and its role in the nervous system has been studied⁴⁴. AQP4 is over-activated in rat articular chondrocytes and high homologues of AQP4 between rat and human⁴⁵; however, the pathological role of AQP4 in RA is still unclear. AQP9 was strongly induced upon treatment with TNF- α in FLS and was also expressed in the RA and OA synovial tissues⁴¹. Although the pathological roles of AQP in the synovial tissues remain to be elucidated, experimental evidence has revealed that AQPs are

Table 1. AQPs in FLS.

AQP	Mechanism	Species	Ref.
AQP1	Hydrarthrosis and joint swelling Inhibiting NF- κ B pathway by AQP1 inhibitor	Adjuvant-injected arthritis rats	42
AQP4	Over-activated AQP4 in articular chondrocytes	Articular chondrocytes, adjuvant-injected arthritis rats	45
AQP9	Hydrarthrosis	HepG2, FLS from OA and RA patients	41

involved in the pathogenesis of hydrarthrosis and synovitis (Table 1).

4.2. K^+ channels

Ca^{2+} -activated potassium channel $K_{Ca1.1}$ (known as BK, Maxi-K, Slo1, or *KCNMA1*) is the only member of the $K_{Ca1.1}$ potassium channel family⁴⁶. The $K_{Ca1.1}$ channel consists of α -subunits and β -subunits comprising of four different isoforms (β_1 , β_2 , β_3 , and β_4)⁴⁷. The $K_{Ca1.1}$ was a major K^+ channel expressed in FLS plasma membrane in RA⁴⁸. Blocking the $K_{Ca1.1}$ channel in RA FLS by inhibiting the α -subunit interrupted Ca^{2+} homeostasis; the proliferation, migration, and the invasiveness of cells; and the cytokines and chemokines⁴⁸. The K^+ channels in the plasma membrane of cells play a critical role in regulating β_1 integrins by influencing Ca^{2+} homeostasis⁴⁹. The FLS cells express a variety of integrins, α_4 , α_5 , α_6 , and β_1 isotype⁵⁰. Blocking of $K_{Ca1.1}$ channel interrupted Ca^{2+} homeostasis, thus affecting integrin expression⁴⁹. Enhanced integrin ligation increased cytokine signalling and growth factor expression, thus leading to the expression of matrix MMPs⁵⁰. Blocking of $K_{Ca1.1}$ activity or its expression reduced the FLS proliferation and expression of pro-MMP2 and attenuated the subsequent FLS invasion. On the contrary, activated $K_{Ca1.1}$ or overexpression of the channel enhanced the invasiveness of FLS⁵¹. Regulation of $K_{Ca1.1}$ of FLS also affected the proliferation and migration of CCR7⁺ effector memory T cells, another major cell type implicated in the progression of RA⁵².

4.3. Acid-sensing ion channels

Acid-sensing ion channels (ASICs) mediate tissue acidosis by pH changes are known as voltage-insensitive, ligand-gated cation channels with protons^{53,54}. The ASICs are associated with inflammatory pain, and especially ASIC1 and ASIC3 contribute to the musculoskeletal pain⁵⁵. The ASIC3 is expressed in the sensory neurons that innervate the synovial joints by increasing the intracellular Ca^{2+} levels upon sensing a decrease of pH in the inflamed joint^{56,57}. Synovial inflammation and inflammatory cytokine levels were increased that led to joint destruction in ASIC3^{-/-} mice⁵⁵. FLS were activated with the decrease in pH; the acidic environment increased the intracellular Ca^{2+} levels by ASIC3⁵⁷. Activation of FLS in acidic pH mediates the accumulation of inflammatory cytokines. In addition, activation of ASIC3 by acidic pH evokes Ca^{2+} signalling, which lead to the apoptosis of FLS by phosphorylation of the MAP kinase ERK in synovial inflammation; thus, it could be a blockade of synovial proliferation⁵⁸. Activation of ASIC3 can be a therapeutic strategy for reducing inflammatory FLS level and subsequent disease progression in an inflamed joint.

4.4. Ca^{2+} signalling of FLS

Intracellular Ca^{2+} plays crucial roles in various physiological processes, including the flow of nerve impulses, muscle contraction, cell division, and hormone secretion⁵⁹. Enhanced Ca^{2+} -activated phosphatase calcineurin activity and Ca^{2+} release by

proinflammatory cytokine were observed in RA FLS, suggesting that dysregulated Ca^{2+} signalling involved in the pathogenesis of chronic arthritis⁶⁰. In addition, synovial fluid of patients with RA contains ATP⁶¹ and FLS expressed P2X7 receptor and functionally involved in ATP-dependent Ca^{2+} release and subsequently mediated IL-6 release⁶². Generally, the cytosol is surrounded by two major Ca^{2+} sources; the intracellular Ca^{2+} stores including sarco/endoplasmic reticulum (SR/ER), nucleus, golgi, and mitochondria and the extracellular media⁶³. The Ca^{2+} is released from intracellular stores or enters into the cells through the plasma membrane⁶⁴. The Ca^{2+} homeostasis is maintained by two types of membrane ATPase, the SR/ER Ca^{2+} -ATPase (SERCA) and plasma membrane Ca^{2+} -ATPase (PMCA). These pumps are involved in reduction of cytosolic Ca^{2+} , from cytosol to intracellular Ca^{2+} stores by the SERCA and to the extracellular space by the PMCA⁶⁵. Na^+/Ca^{2+} exchangers are also known to have a critical role in Ca^{2+} removing mechanism with Na^+ regulation^{66,67}. The ER also contains inositol-1,4,5-trisphosphate (IP_3) receptors (IP_3Rs) and ryanodine receptors (RyRs), which provide conduits for the rapid release of Ca^{2+} ⁶⁸. The agonist stimulation such as receptor activation leads to the generation of IP_3 , which releases to the cytosol and binds to the intracellular membranes to release Ca^{2+} from the intracellular stores of Ca^{2+} ⁶⁹. Although the Ca^{2+} signalling and its signalling proteins have been well established, the network of Ca^{2+} signalling in FLS needs to be clarified more extensively.

4.5. TRP channels

Although there is relatively low evidence in Ca^{2+} signalling network in FLS, studies of transient receptor potential (TRP) channels have been performed in various reports. The TRP channels have been known to be nonselective cation channels and play a critical role in inflammatory pain of arthritis^{70,71}. FLS express the TRP family proteins, including TRPC (TRPC-canonical) 1, TRPC5, TRPA (TRP-ankyrin) 1, TRPV (TRPV-vanilloid) 1, TRPV2, TRPV4, TRPM (TRPM-melastatin) 7, and TRPM8^{72–76}. We will discuss FLS-related TRP channel activation and will provide information on the following section. The detailed mechanism is summarised in Figure 2 and Table 2.

4.5.1. TRPC

The TRPC1 and TRPC5 were expressed in secretory FLS^{76,77}. The expression of oxidoreductase thioredoxin, a well-known oxidative stress marker, was increased in RA synovial fluids to counteract oxidative stress^{78–80}. The thioredoxin was considered as a costimulatory component with cytokines in FLS⁸¹ and can be secreted⁸². Extracellular reduced form of thioredoxin enhanced the activities of TRPC1 and TRPC5 channels as new extracellular targets⁷⁷. Inhibition of these channels by antibodies enhanced MMP secretion and suppressed the thioredoxin-mediated inhibitory effect on secretion⁷⁷. More recently, the study of TRPC5 KO mice and inhibition of TRPC5 channels by antagonist addressed the enhanced inflamed joint and hyperalgesia⁷⁶, suggesting that functional modulation of TRPC 1/5 could be considered as therapeutic targets for RA.

4.5.2. TRPA

The TRPA1 is a cold-sensitive and Ca^{2+} -permeable nonselective cation channel and plays an essential role in inflammation and pain⁸³. For the evidence of TRPA1 expression in FLS, mRNA of TRPA1 has been detected in SW982 human synoviocytes⁷⁴. Diphenyleneiodonium (DPI) as a TRPA1 activator induced Ca^{2+} signal in TRPA1-expressing FLS and pain response in ddY mice⁷⁵. More recently, it has been reported that proinflammatory FLS can be attenuated by TRPA1 activation. TNF-stimulated FLS enhanced protein level of TRPA1 and subsequent stimulation of TRPA1 enhanced the necrosis⁸⁴.

4.5.3. TRPV

The TRPV channels sense heat, protons, lipids, and osmolarity^{85,86}. The RA and OA patients possess pain linked to TRPV1⁷³. Capsaicin, an agonist of TRPV1, increased IL-6 mRNA and protein levels by promoting pro-inflammatory cytokines⁸⁷. Activation of TRPV1 enhanced mRNA level and protein level of IL-6 in FLS from RA and OA patient and application of TRPV1 antagonist could be therapeutic strategy to modulate nociception from arthritis⁷³. Activated FLS-mediated TNF- α secretion enhanced the expression of TRPV1 and TRPV4 in SW982 human synoviocytes⁸⁸. Especially, 4 α -phorbol-12,13-didecanoate (4 α -PDD), a selective TRPV4 agonist and hypotonic stimulation induced an increased intracellular Ca^{2+} level and decreased IL-8 secretion in RA⁸⁹. The enhanced TRPV2

expression was associated with invasion of FLS from rats using gene profile technique⁷². However, functional activation of TRPV2 by specific TRPV2 agonists, O1821 and LER13, dramatically reduced IL-1 β -mediated expression of the MMP2 and MMP3 proteins in FLS and reduced the severity of disease and genetic deletion of TRPV2 enhanced the invasiveness of FLS⁹⁰. Expression of TRPV2 involves in invasion mechanism and further functional stimulation of TRPV2 attenuates the invasiveness, suggesting that regulation of TRPV2 can be also novel therapeutic strategy of RA such as TRPC and TRPA channels.

4.5.4. TRPM

The Ca^{2+} entry through TRPM3 is involved in cell survival, death, growth, and differentiation⁹¹. Hyaluronan, known as the major component of the extracellular matrix, was increased in RA patients⁹². The increased secretion of hyaluronan from RA FLS was reduced by TRPM3 activator pregnenolone sulphate, activating TRPM3-mediated Ca^{2+} entry^{93,94}. The TRPM7 mediates a variety of functions, such as cell cycle, migration differentiation, and regulation of Ca^{2+} homeostasis and it is correlated with the oxidative stress-induced cell injury^{95–97}. It has been proposed that hypoxia and low glucose also lead to ER stress in RA joints⁹⁸. Inhibition of TRPM7 by Gd^{3+} and 2-aminoethoxydiphenyl borate (2-APB) induced RA FLS apoptosis by activating ER stress⁹⁹.

5. Future perspectives

We limited our review to the most relevant channels related to cell migration in RA FLS. Evidence related to cell migration by water and ion channels addressed the housekeeping functions. The migration or invasion is the major feature of cancer cell. The major consequences between the cancer cells and inflamed FLS are hypoxia and acidic circumstances¹⁰⁰. The cancer cells employ a circuit of ion transporters and enzymes to avoid the detrimental consequences of hypoxic and acidic tumour microenvironment. Alterations of CA IX and CA XII are associated with various cancers and considered oncogenic factors¹⁰¹. The overexpression of CA IX and XII in the inflamed synovium^{28,29} provides the several similarities of pathology between inflamed synovium and cancer (Figure 3). Such similarities between microenvironments including hypoxia, acidic pH, and enhanced CA IX and XII can be speculated, as RA FLS would share the migration mechanism with cancer cells. Thus, therapeutic options in cancer therapy can be expanded and exploited for the RA model.

Currently, experimental evidence for the involvement of CAs and FLS membrane channels in RA is limited. The physiological and pathological roles of ion channels and transporters in dynamic FLS migration have not yet been studied in detail. Here, we have summarised the studies on membrane channels and regulatory enzymes of RA-FLS with an aim to understand their migrated state. However, many questions regarding RA-FLS still need to be clarified. What are the exact molecular mechanisms by which ion transporter affects the FLS migration apparatus? What




Figure 2. Activators of TRP channels in RA FLS. The activators of FLS-associated TRP channels are summarised. The mechanism of action is represented in Table 2. Activation of TRP channels increases intracellular Ca^{2+} level and is involved in various functions including the reduced MMP secretion, joint destruction, enhancement of pain, and apoptosis of inflamed FLS. TRPV: transient receptor potential vanilloid channels; TRPC: transient receptor potential canonical channels; TRPM: transient receptor potential melastatin channels; TRPA: transient receptor potential ankyrin channels.

Table 2. TRP channels in FLS.

TRPs	Mechanism	Species	Ref.
TRPC1/TRPC5	Reduced MMP secretion and joint inflammation	Human FLS, mouse joint tissue	76,77
TRPV1	Promoted inflammation and joint destruction	SW982 human synovial cells	88
TRPV2	Reduced expression of the MMP2 and MMP3 proteins	FLS from DA (severe and erosive arthritis)	90
TRPV4	Reduced IL-8 production	FLS with RA and without RA, MH7A	89
TRPA1	Increased pain-related response	Human FLS, ddY mice	75
TRPM3	Decreased hyaluronan secretion	HIG-82 cells (FLS cell-line), joint tissue of RA patients	94
TRPM7	Activated ER stress, increased apoptosis of FLS	FLS from RA	99




Figure 3. Potential function of CAs in inflamed synovium. CA I was overexpressed in the synovium of the patients with ankylosing spondylitis²⁷. The overexpression of CA IX and XII was revealed in the inflamed synovium. Although determination of CA isoenzymes on the RA FLS remains unclear, there are several similarities between inflamed synovium and cancer. CA: carbonic anhydrase; pH_e : extracellular pH.

are the exact components of synovial fluid that mediate the FLS dynamics? What are the components affecting the differential expression of CAs and membrane channels in FLS? What is the combined mechanism of CAs as regulatory enzymes? Several membrane channels and transporters show tissue-specific expression. Thus, unravelling the mechanisms by which ion channels and transporters are positioned in and modulate the migration of activated FLS will be a rewarding pursuit for the coming years. The motivation of channel physiologists is also needed to develop potential therapeutics to counter the critical pathophysiological involvement of FLS migration in joints in RA.

Authors' contributions

JHH and MJJ conceptualized and designed the study. MJJ prepared and revised the manuscript critically for intellectual content. JHH approved the final version of the manuscript.

Disclosure statement

The authors declare that they have no conflicts of interest with the contents of this article.

Funding

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korean Government (MSIT; [NRF-2019R1F1A1046785]) and by the grant from the Gachon University Gil Medical Center Project [FRD 2018-07].

ORCID

Min Jeong Ji <http://orcid.org/0000-0002-3063-7833>
Jeong Hee Hong <http://orcid.org/0000-0003-3623-2201>

References

- Firestein GS. Evolving concepts of rheumatoid arthritis. *Nature* 2003;423:356–61.
- Xing R, Jin Y, Sun L, et al. Interleukin-21 induces migration and invasion of fibroblast-like synoviocytes from patients with rheumatoid arthritis. *Clin Exp Immunol* 2016;184:147–58.
- Hitchon CA, El-Gabalawy HS. The synovium in rheumatoid arthritis. *Open Rheumatol J* 2011;5:107–14.
- Abeles AM, Pillinger MH. The role of the synovial fibroblast in rheumatoid arthritis: cartilage destruction and the regulation of matrix metalloproteinases. *Bull NYU Hosp Jt Dis* 2006;64:20–4.
- Muller-Ladner U, Pap T, Gay RE, et al. Mechanisms of disease: the molecular and cellular basis of joint destruction in rheumatoid arthritis. *Nat Clin Pract Rheumatol* 2005;1:102–10.
- Koch AE, Kunkel SL, Burrows JC, et al. Synovial tissue macrophage as a source of the chemotactic cytokine IL-8. *J Immunol* 1991;147:2187–95.
- Yap HY, Tee SZ, Wong MM, et al. Pathogenic role of immune cells in rheumatoid arthritis: implications in clinical treatment and biomarker development. *Cells* 2018;7:161.
- Thornton SC, Por SB, Penny R, et al. Identification of the major fibroblast growth factors released spontaneously in inflammatory arthritis as platelet derived growth factor and tumour necrosis factor- α . *Clin Exp Immunol* 1991;86:79–86.
- Chu CQ, Field M, Abney E, et al. Transforming growth factor- β 1 in rheumatoid synovial membrane and cartilage/pannus junction. *Clin Exp Immunol* 1991;86:380–6.
- Bartok B, Firestein GS. Fibroblast-like synoviocytes: key effector cells in rheumatoid arthritis. *Immunol Rev* 2010;233:233–55.
- Schwab A, Fabian A, Hanley PJ, Stock C. Role of ion channels and transporters in cell migration. *Physiol Rev* 2012;92:1865–913.
- Schwab A. Function and spatial distribution of ion channels and transporters in cell migration. *Am J Physiol Renal Physiol* 2001;280:F739–F747.
- Mor A, Abramson SB, Pillinger MH. The fibroblast-like synovial cell in rheumatoid arthritis: a key player in inflammation and joint destruction. *Clin Immunol* 2005;115:118–28.
- Firestein GS. Invasive fibroblast-like synoviocytes in rheumatoid arthritis. Passive responders or transformed aggressors? *Arthritis Rheum* 1996;39:1781–90.
- Tu J, Hong W, Zhang P, et al. Ontology and function of fibroblast-like and macrophage-like synoviocytes: how do they talk to each other and can they be targeted for rheumatoid arthritis therapy? *Front Immunol* 2018;9:1467.
- Iwanaga T, Shikichi M, Kitamura H, et al. Morphology and functional roles of synoviocytes in the joint. *Arch Histol Cytol* 2000;63:17–31.
- Chegwidden WR, Dodgson SJ, Spencer IM. The roles of carbonic anhydrase in metabolism, cell growth and cancer in animals. *EXS* 2000;90:343–63.
- Missner A, Kugler P, Saparov SM, et al. Carbon dioxide transport through membranes. *J Biol Chem* 2008;283:25340–7.
- Swietach P, Vaughan-Jones RD, Harris AL. Regulation of tumor pH and the role of carbonic anhydrase 9. *Cancer Metastasis Rev* 2007;26:299–310.

20. Supuran CT, Scozzafava A, Casini A. Carbonic anhydrase inhibitors. *Med Res Rev* 2003;23:146–89.
21. Temperini C, Scozzafava A, Supuran CT. Carbonic anhydrase inhibitors. X-ray crystal studies of the carbonic anhydrase ii-trithiocarbonate adduct-an inhibitor mimicking the sulfonamide and urea binding to the enzyme. *Bioorg Med Chem Lett* 2010;20:474–8.
22. Morgan PE, Supuran CT, Casey JR. Carbonic anhydrase inhibitors that directly inhibit anion transport by the human $\text{Cl}^-/\text{HCO}_3^-$ exchanger, AE1. *Mol Membr Biol* 2004;21:423–33.
23. Alvarez BV, Loisel FB, Supuran CT, et al. Direct extracellular interaction between carbonic anhydrase IV and the human NBC1 sodium/bicarbonate co-transporter. *Biochemistry* 2003;42:12321–9.
24. Farr M, Garvey K, Bold AM, et al. Significance of the hydrogen ion concentration in synovial fluid in rheumatoid arthritis. *Clin Exp Rheumatol* 1985;3:99–104.
25. Steen KH, Steen AE, Reeh PW. A dominant role of acid pH in inflammatory excitation and sensitization of nociceptors in rat skin, in vitro. *J Neurosci* 1995;15:3982–9.
26. Lardner A. The effects of extracellular pH on immune function. *J Leukoc Biol* 2001;69:522–30.
27. Chang X, Han J, Zhao Y, et al. Increased expression of carbonic anhydrase I in the synovium of patients with ankylosing spondylitis. *BMC Musculoskelet Disord* 2010;11:279.
28. Margheri F, Ceruso M, Carta F, et al. Overexpression of the transmembrane carbonic anhydrase isoforms IX and XII in the inflamed synovium. *J Enzyme Inhib Med Chem* 2016;31:60–3.
29. Tafreshi NK, Lloyd MC, Proemsey JB, et al. Evaluation of CAIX and CAXII expression in breast cancer at varied O_2 levels: CAIX is the superior surrogate imaging biomarker of tumor hypoxia. *Mol Imaging Biol* 2016;18:219–31.
30. Alver A, Şentürk A, Çakırbay H, et al. Carbonic anhydrase ii autoantibody and oxidative stress in rheumatoid arthritis. *Clin Biochem* 2011;44:1385–9.
31. Svastova E, Witariski W, Csaderova L, et al. Carbonic anhydrase IX interacts with bicarbonate transporters in lamellipodia and increases cell migration via its catalytic domain. *J Biol Chem* 2012;287:3392–402.
32. Wykoff CC, Beasley NJ, Watson PH, et al. Hypoxia-inducible expression of tumor-associated carbonic anhydrases. *Cancer Res* 2000;60:7075–83.
33. Semenza GL. Targeting HIF-1 for cancer therapy. *Nat Rev Cancer* 2003;3:721–32.
34. Gumbiner BM. Cell adhesion: the molecular basis of tissue architecture and morphogenesis. *Cell* 1996;84:345–57.
35. Mount DB, Romero MF. The SLC26 gene family of multifunctional anion exchangers. *Pflugers Arch* 2004;447:710–21.
36. Parks SK, Pouyssegur J. The $\text{Na}^+/\text{HCO}_3^-$ co-transporter SLC4A4 plays a role in growth and migration of colon and breast cancer cells. *J Cell Physiol* 2015;230:1954–63.
37. Lee MD, King LS, Agre P. The aquaporin family of water channel proteins in clinical medicine. *Medicine (Baltimore)* 1997;76:141–56.
38. Wakayama Y. Aquaporin expression in normal and pathological skeletal muscles: a brief review with focus on AQP4. *J Biomed Biotechnol* 2010;2010:731569.
39. Liu H, Wintour EM. Aquaporins in development – a review. *Reprod Biol Endocrinol* 2005;3:18.
40. Trujillo E, Gonzalez T, Marin R, et al. Human articular chondrocytes, synoviocytes and synovial microvessels express aquaporin water channels; upregulation of aqp1 in rheumatoid arthritis. *Histol Histopathol* 2004;19:435–44.
41. Nagahara M, Waguri-Nagaya Y, Yamagami T, et al. TNF- α -induced aquaporin 9 in synoviocytes from patients with OA and RA. *Rheumatology (Oxford)* 2010;49:898–906.
42. Cai L, Chen WN, Li R, et al. Therapeutic effect of acetazolamide, an aquaporin 1 inhibitor, on adjuvant-induced arthritis in rats by inhibiting NF- κ B signal pathway. *Immunopharmacol Immunotoxicol* 2018;40:117–25.
43. Yang B, van Hoek AN, Verkman AS. Very high single channel water permeability of aquaporin-4 in baculovirus-infected insect cells and liposomes reconstituted with purified aquaporin-4. *Biochemistry* 1997;36:7625–32.
44. Xiao M, Hu G. Involvement of aquaporin 4 in astrocyte function and neuropsychiatric disorders. *CNS Neurosci Ther* 2014;20:385–90.
45. Cai L, Lei C, Li R, et al. Overexpression of aquaporin 4 in articular chondrocytes exacerbates the severity of adjuvant-induced arthritis in rats: an in vivo and in vitro study. *J Inflamm (Lond)* 2017;14:6.
46. Jenkinson DH. Potassium channels-multiplicity and challenges. *Br J Pharmacol* 2006;147Suppl.1:S63–S71.
47. Lippiat JD, Standen NB, Harrow ID, et al. Properties of BK(Ca) channels formed by bicistronic expression of hSloalpha and beta1-4 subunits in HEK293 cells. *J Membr Biol* 2003;192:141–8.
48. Hu X, Laragione T, Sun L, et al. KCa1.1 potassium channels regulate key proinflammatory and invasive properties of fibroblast-like synoviocytes in rheumatoid arthritis. *J Biol Chem* 2012;287:4014–22.
49. Tanner MR, Pennington MW, Laragione T, et al. KCa1.1 channels regulate β 1-integrin function and cell adhesion in rheumatoid arthritis fibroblast-like synoviocytes. *FASEB J* 2017;31:3309–20.
50. Lowin T, Straub RH. Integrins and their ligands in rheumatoid arthritis. *Arthritis Res Ther* 2011;13:244.
51. Tanner MR, Hu X, Huq R, et al. KCa1.1 inhibition attenuates fibroblast-like synoviocyte invasiveness and ameliorates disease in rat models of rheumatoid arthritis. *Arthritis Rheumatol* 2015;67:96–106.
52. Tanner MR, Pennington MW, Chauhan SS, et al. KCa1.1 and Kv1.3 channels regulate the interactions between fibroblast-like synoviocytes and T lymphocytes during rheumatoid arthritis. *Arthritis Res Ther* 2019;21:6.
53. Kweon HJ, Suh BC. Acid-sensing ion channels (ASICs): therapeutic targets for neurological diseases and their regulation. *BMB Rep* 2013;46:295–304.
54. Grunder S, Pusch M. Biophysical properties of acid-sensing ion channels (ASICs). *Neuropharmacology* 2015;94:9–18.
55. Sluka KA, Rasmussen LA, Edgar MM, et al. Acid-sensing ion channel 3 deficiency increases inflammation but decreases pain behavior in murine arthritis. *Arthritis Rheum* 2013;65:1194–202.
56. Waldmann R, Champigny G, Lingueglia E, et al. H(+)-gated cation channels. *Ann N Y Acad Sci* 1999;868:67–76.
57. Kolker SJ, Walder RY, Usachev Y, et al. Acid-sensing ion channel 3 expressed in type b synoviocytes and chondrocytes modulates hyaluronan expression and release. *Ann Rheum Dis* 2010;69:903–9.
58. Gong W, Kolker SJ, Usachev Y, et al. Acid-sensing ion channel 3 decreases phosphorylation of extracellular signal-

- regulated kinases and induces synoviocyte cell death by increasing intracellular calcium. *Arthritis Res Ther* 2014;16: R121.
59. Berridge MJ, Lipp P, Bootman MD. The versatility and universality of calcium signalling. *Nat Rev Mol Cell Biol* 2000;1: 11–21.
 60. Yoo SA, Park BH, Park GS, et al. Calcineurin is expressed and plays a critical role in inflammatory arthritis. *J Immunol* 2006;177:2681–90.
 61. Ryan LM, Rachow JW, McCarty DJ. Synovial fluid ATP: a potential substrate for the production of inorganic pyrophosphate. *J Rheumatol* 1991;18:716–20.
 62. Caporali F, Capecchi PL, Gamberucci A, et al. Human rheumatoid synoviocytes express functional P2X7 receptors. *J Mol Med (Berl)* 2008;86:937–49.
 63. Hannaert-Merah Z, Combettes L, Coquil JF, et al. Characterization of the co-agonist effects of strontium and calcium on myo-inositol trisphosphate-dependent ion fluxes in cerebellar microsomes. *Cell Calcium* 1995;18: 390–9.
 64. Bootman MD, Berridge MJ. The elemental principles of calcium signaling. *Cell* 1995;83:675–8.
 65. Brini M, Bano D, Manni S, et al. Effects of PMCA and SERCA pump overexpression on the kinetics of cell Ca^{2+} signalling. *EMBO J* 2000;19:4926–35.
 66. Floyd R, Wray S. Calcium transporters and signalling in smooth muscles. *Cell Calcium* 2007;42:467–76.
 67. Sommer B, Flores-Soto E, Gonzalez-Avila G. Cellular Na^{+} handling mechanisms involved in airway smooth muscle contraction. *Int J Mol Med* 2017;40:3–9.
 68. Mattson MP, LaFerla FM, Chan SL, et al. Calcium signaling in the ER: its role in neuronal plasticity and neurodegenerative disorders. *Trends Neurosci* 2000;23:222–9.
 69. Mikoshiba K, Furuichi T, Miyawaki A, et al. Structure and function of inositol 1,4,5-trisphosphate receptor. *Ann N Y Acad Sci* 1993;707:178–97.
 70. Wiesmann UN, DiDonato S, Herschkowitz NN. Effect of chloroquine on cultured fibroblasts: release of lysosomal hydrolases and inhibition of their uptake. *Biochem Biophys Res Commun* 1975;66:1338–43.
 71. Fernandes ES, Russell FA, Spina D, et al. A distinct role for transient receptor potential ankyrin 1, in addition to transient receptor potential vanilloid 1, in tumor necrosis factor alpha-induced inflammatory hyperalgesia and Freund's complete adjuvant-induced monoarthritis. *Arthritis Rheum* 2011;63:819–29.
 72. Laragione T, Brenner M, Li W, Gulko PS. Cia5d regulates a new fibroblast-like synoviocyte invasion-associated gene expression signature. *Arthritis Res Ther* 2008;10:R92.
 73. Engler A, Aeschlimann A, Simmen BR, et al. Expression of transient receptor potential vanilloid 1 (TRPV1) in synovial fibroblasts from patients with osteoarthritis and rheumatoid arthritis. *Biochem Biophys Res Commun* 2007;359: 884–8.
 74. Kochukov MY, McNearney TA, Fu Y, Westlund KN. Thermosensitive TRP ion channels mediate cytosolic calcium response in human synoviocytes. *Am J Physiol Cell Physiol* 2006;291:C424–32.
 75. Suzuki H, Hatano N, Muraki Y, et al. The NADPH oxidase inhibitor diphenyleneiodonium activates the human TRPA1 nociceptor. *Am J Physiol Cell Physiol* 2014;307:C384–94.
 76. Alawi KM, Russell FA, Aubdool AA, et al. Transient receptor potential canonical 5 (TRPC5) protects against pain and vascular inflammation in arthritis and joint inflammation. *Ann Rheum Dis* 2017;76:252–60.
 77. Xu SZ, Sukumar P, Zeng F, et al. TRPC channel activation by extracellular thioredoxin. *Nature* 2008;451:69–72.
 78. Kochukov MY, McNearney TA, Yin H, et al. Tumor necrosis factor-alpha (TNF-alpha) enhances functional thermal and chemical responses of TRP cation channels in human synoviocytes. *Mol Pain* 2009;5:49.
 79. Laragione T, Cheng KF, Tanner MR, et al. The cation channel TRPV2 is a new suppressor of arthritis severity, joint damage, and synovial fibroblast invasion. *Clin Immunol* 2015;158:183–92.
 80. Itoh Y, Hatano N, Hayashi H, et al. An environmental sensor, trpv4 is a novel regulator of intracellular Ca^{2+} in human synoviocytes. *Am J Physiol Cell Physiol* 2009;297: C1082–90.
 81. Stuhlmeier KM. Aspects of the biology of hyaluronan, a largely neglected but extremely versatile molecule. *Wien Med Wochenschr* 2006;156(21–22):563–568.
 82. Li X, Wang X, Wang Y, et al. Inhibition of transient receptor potential melastatin 7 (trpm7) channel induces ra flss apoptosis through endoplasmic reticulum (er) stress. *Clin Rheumatol* 2014;33(11):1565–1574.
 83. Burke-Gaffney A, Callister ME, Nakamura H. Thioredoxin: friend or foe in human disease? *Trends Pharmacol Sci* 2005;26:398–404.
 84. Maurice MM, Nakamura H, Gringhuis S, et al. Expression of the thioredoxin-thioredoxin reductase system in the inflamed joints of patients with rheumatoid arthritis. *Arthritis Rheum* 1999;42:2430–9.
 85. Jikimoto T, Nishikubo Y, Koshiba M, et al. Thioredoxin as a biomarker for oxidative stress in patients with rheumatoid arthritis. *Mol Immunol* 2002;38:765–72.
 86. Yoshida S, Katoh T, Tetsuka T, et al. Involvement of thioredoxin in rheumatoid arthritis: its costimulatory roles in the TNF-alpha-induced production of IL-6 and IL-8 from cultured synovial fibroblasts. *J Immunol* 1999;163:351–8.
 87. Rubartelli A, Bajetto A, Allavena G, et al. Secretion of thioredoxin by normal and neoplastic cells through a leaderless secretory pathway. *J Biol Chem* 1992;267:24161–4.
 88. Meseguer V, Alpizar YA, Luis E, et al. TRPA1 channels mediate acute neurogenic inflammation and pain produced by bacterial endotoxins. *Nat Commun* 2014;5:3125.
 89. Lowin T, Bleck J, Schneider M, Pongratz G. Selective killing of proinflammatory synovial fibroblasts via activation of transient receptor potential ankyrin (TRPA1). *Biochem Pharmacol* 2018;154:293–302.
 90. Liedtke W. Role of TRPV ion channels in sensory transduction of osmotic stimuli in mammals. *Exp Physiol* 2007;92: 507–12.
 91. Benham CD, Gunthorpe MJ, Davis JB. TRPV channels as temperature sensors. *Cell Calcium* 2003;33:479–87.
 92. Reilly CA, Johansen ME, Lanza DL, et al. Calcium-dependent and independent mechanisms of capsaicin receptor (TRPV1)-mediated cytokine production and cell death in human bronchial epithelial cells. *J Biochem Mol Toxicol* 2005;19:266–75.
 93. Grimm C, Kraft R, Schultz G, Harteneck C. Activation of the melastatin-related cation channel TRPM3 by D-erythro-sphingosine. *Mol Pharmacol* 2005;67:798–805.
 94. Engstrom-Laurent A, Hallgren R. Circulating hyaluronate in rheumatoid arthritis: relationship to inflammatory activity

- and the effect of corticosteroid therapy. *Ann Rheum Dis* 1985;44:83–8.
95. Ciurtin C, Majeed Y, Naylor J, et al. Trpm3 channel stimulated by pregnenolone sulphate in synovial fibroblasts and negatively coupled to hyaluronan. *BMC Musculoskelet Disord* 2010;11:111.
96. Cheng H, Feng JM, Figueiredo ML, et al. Transient receptor potential melastatin type 7 channel is critical for the survival of bone marrow derived mesenchymal stem cells. *Stem Cells Dev* 2010;19:1393–403.
97. Sun HS, Jackson MF, Martin LJ, et al. Suppression of hippocampal TRPM7 protein prevents delayed neuronal death in brain ischemia. *Nat Neurosci* 2009;12:1300–7.
98. Aarts M, Iihara K, Wei WL, et al. A key role for TRPM7 channels in anoxic neuronal death. *Cell* 2003;115:863–77.
99. Fantinelli JC, Orlowski A, Aiello EA, Mosca SM. The electrogenic cardiac sodium bicarbonate co-transporter (NBCe1) contributes to the reperfusion injury. *Cardiovasc Pathol* 2014;23:224–30.
100. Li X, Wang X, Wang Y, et al. Inhibition of transient receptor potential melastatin 7 (TRPM7) channel induces RA FLSs apoptosis through endoplasmic reticulum (ER) stress. *Clin Rheumatol* 2014;33:1565–74.
101. McDonald PC, Swayampakula M, Dedhar S. Coordinated regulation of metabolic transporters and migration/invasion by carbonic anhydrase ix. *Metabolites* 2018;8:20.