



ORIGINAL ARTICLE

Parental tobacco consumption and child development[☆]



Nadine F. Santos^{a,*}, Raquel A. Costa^{a,b}

^a Universidade de Trás-os-Montes e Alto Douro, Vila Real, Portugal

^b Universidade Europeia Laureate International Universities, Lisbon, Portugal

Received 3 June 2014; accepted 17 September 2014

Available online 27 February 2015

KEYWORDS

Psychomotor development;
Tobacco use;
Parenting

Abstract

Objective: To analyze the association between parental tobacco consumption and the prevalence of psychomotor development disorders in children between 6 and 22 months of age.

Method: One hundred and nine mothers, fathers, and their babies participated in the study. The sociodemographic and clinical conditions were assessed using questionnaires. Tobacco consumption was assessed using the Fagerström Test for Nicotine Dependence (FTND). Child development was evaluated using the Scale of Psychomotor Development in Early Childhood.

Results: There was a significant negative correlation between the father's morning smoking (FTND) and the child's language development quotient; $r = -0.41$, $p = 0.005$, $r^2 = 0.15$. The children of mothers without nicotine dependence had a higher mean language development quotient than children of mothers with nicotine dependence; $F(1, 107) = 5.51$, $p = 0.021$, $\eta_p^2 = 0.05$.

Conclusion: Parental smoking appears to have a detrimental effect on child development.
© 2015 Sociedade Brasileira de Pediatria. Published by Elsevier Editora Ltda. All rights reserved.

PALAVRAS-CHAVE

Desenvolvimento psicomotor;
Consumo de tabaco;
Parentalidade

Consumo de tabaco parental e desenvolvimento infantil

Resumo

Objetivo: Analisar a relação entre o consumo de tabaco parental e a prevalência de distúrbios no desenvolvimento psicomotor em crianças entre os seis e os vinte e dois meses de idade.

Método: Cento e nove mães, pais e seus bebês participaram no estudo. As circunstâncias sociodemográficas e clínicas foram avaliadas com recurso a questionários. O consumo de tabaco foi avaliado utilizando o Teste de Fagerström para a Dependência Tabágica (Heatherton, Kozlowski, Frecker, & Fagerström, 1991). O desenvolvimento infantil foi avaliado utilizando a Escala do Desenvolvimento Psicomotor da Primeira Infância (Brunet & Lézine, 1951).

[☆] Please cite this article as: Santos NF, Costa RA. Parental tobacco consumption and child development. J Pediatr (Rio J). 2015;91:366–72.

* Corresponding author.

E-mail: nadifernandessantos@gmail.com (N.F. Santos).

Resultados: Há uma correlação negativa significativa entre o fumo matinal (FTND) do pai e o quociente de desenvolvimento de linguagem da criança, $r = -0,41$, $p = 0,005$, $r^2 = 0,15$. As crianças de mães sem dependência tabágica têm em média um quociente de desenvolvimento de linguagem superior às crianças de mães com dependência tabágica, $F(1,107) = 5,51$, $p = 0,021$, $\eta_p^2 = 0,05$.

Conclusão: O consumo de tabaco parental parece ter um efeito prejudicial para o desenvolvimento da criança.

© 2015 Sociedade Brasileira de Pediatria. Publicado por Elsevier Editora Ltda. Todos os direitos reservados.

Introduction

Development in the first years of life is essential. Several environmental factors, such as the parental consumption of substances, can increase the likelihood of developmental difficulties in childhood,¹ especially in the emotional, educational,² social, behavioral, and psychological levels.³ Tobacco consumption, specifically—which has high prevalence in Portugal (22%)⁴—is an important public health problem, and has been associated with difficulties in self-regulation; increased excitability and activation in the neonatal period;⁵ lower birth weight;^{6,7} learning difficulties;⁶ lower volume of the frontal and cerebellar lobes—responsible for emotional functioning, impulse control, and attention;⁸ smaller head circumference;⁹ cognitive and language neurodevelopmental disorders;^{10,11} and childhood emotional and behavioral disorders.⁶ This association can be explained by the fact that during childhood, the brain continues to develop and is particularly sensitive to environmental pollutants,⁶ or by brain disorders resulting from exposure to nicotine during pregnancy.⁸

As environmental risk factors appear to be related to children's developmental disorders, especially motor, language, social, cognitive, behavioral, and psychological disorders, the study of psychomotor development associated to these factors is the aim of this study. Although there is a great deal of literature related to child development, the association between tobacco use by both parents and psychomotor development is still largely unknown. Therefore, this study differs from previous studies due to the fact that (1) it analyzes several aspects of child development - posture, language, visual-motor coordination, and social—rather than being limited to overall development; (2) most studies focus on the effects of this consumption on children's health, and/or during pregnancy and not after it; (3) both parents were considered.

Methods

Participants

Participants were recruited from four day care centers located in the city of Funchal-Madeira, Portugal, after authorization by the Direção Regional de Educação (Regional Education Board). Most of the participants were of Portuguese origin (94.3%) and white (98.0%).

Participation in the study was proposed to 124 mothers and 124 fathers; 87.9% agreed to participate, 9.3% refused

to participate due to lack of free time, and 2.8% were not interested in participating. Thus, the sample consisted of 109 mothers, 109 fathers, and 109 babies. The sample inclusion criterion was: (1) to be the mother/father of a child aged between 6 and 22 months; the exclusion criteria were: (1) illiteracy and (2) the existence of diseases in the babies. The study was performed during 2011 and the data collection phase lasted three months.

Tools

Sociodemographic and clinical data

A questionnaire was used to collect social and demographic data (age, gender, marital status, years of education, professional status, physical and psychological diseases, medical or psychological treatment, number of pregnancies, number of miscarriages, number of children, age of children, and children's physical and psychological diseases), as well as clinical information on the pregnancy and newborn (pregnancy planning, prenatal care, risk pregnancy, gestational age, type of delivery, type of anesthesia, Apgar score, weight and height, head circumference, reanimation, health problems at birth, current sleep pattern—monophasic (long periods of continuous sleep) vs. biphasic (alternation between periods of sleep and wakefulness), and type of feeding.

Nicotine dependence

Fagerström Test for Nicotine Dependence (FTND)^{12,13}

This test was developed to compensate for the psychometric limitations of the Fagerström Tolerance Questionnaire,¹² and aims to measure the nicotine dependence of an individual.¹³ It consists of six items related to smoking habits and behaviors, rated on a Likert scale ranging from 0-3 points. Higher results indicate a greater smoking dependence,¹⁴ in which a score of 0-3 indicates absence of nicotine dependence, 4-6 indicates moderate nicotine dependence, and ≥ 7 indicates severe nicotine dependence. The Portuguese version has acceptable psychometric properties, with Cronbach's $\alpha = 0.66$.¹³ Test-retest reliability was ensured by correlation values of the original scale of 0.99. Factor analysis showed the existence of two factors: (1) cigarette consumption—daily consumption patterns—and (2) morning smoking - degree of urgency to restore the level of nicotine after the nighttime abstinence.¹³ In the present study, this tool had an acceptable internal consistency, with Cronbach's $\alpha = 0.73$.

Psychomotor development

Scale of Psychomotor Development in Early Childhood¹⁵

This scale allows the evaluation of the child's developmental level (1-30 months) in each of the following areas: postural-child's movements such as rolling over, sitting, and walking; visual-motor coordination-manipulation of objects, visual-manual coordination, and solving problems; language-expression and understanding; and social-social and personal relations, especially adaptation to social situations, games, and relations with others.¹⁵

It consists of 150 items, which are distributed in levels (1-30). This scale allows the calculation of the Developmental Quotient ($DQ = DA \times 100/CA$), in which DA represents the developmental age and CA the chronological age. A $DQ \leq 75$ indicates lower development, whereas DQ of 76 to 100 indicates normal development, and > 100 demonstrates higher development.¹⁵

The scale validity criterion is 0.68, with a test-retest correlation coefficient of 0.85.¹⁵

Procedures

All evaluation procedures were performed and approved by the ethics committee of the institution. Mothers and fathers received an explanation on the purpose and procedures of the study, and after the informed consent was signed, they were asked to complete a socio-demographic/clinical questionnaire and the FTND^{12,13} (approximate duration of 20-25 minutes).

The Scale of Psychomotor Development in Early Childhood¹⁵ was used to assess the infant's psychomotor development. This assessment was always performed by the same investigator, appropriately trained and familiarized with the scale application procedures. Each baby was observed individually and only at a specific time (approximately 30 to 40 minutes), while the daycare professional was present, but away from the baby's visual range. The questions pertaining to each subtest were answered by the day care professional at the end of each observation.

Statistical procedures

Pearson's Correlation test was used to assess whether there was a correlation between the degree of nicotine dependence (FTND) of the mother and the father and the child's postural, visual-motor, language, social, and overall development quotient.¹⁶

The significance of nicotine dependence factor (FTND ≥ 7) of the mother and father on the composite of the variables child's postural, visual-motor, language, and social development quotient was evaluated by several Multivariate Analysis of Variance (MANOVAs).¹⁷ The significance of the difference between the child's overall development quotient according to nicotine dependence of the mother and father (FTND ≥ 7) was evaluated by using the t-test for independent samples.¹⁶

The chi-squared test was used to assess whether there was an association between child's overall developmental quotient (lower, normal, upper) and the nicotine dependence (FTND ≥ 7) of the mother and father.¹⁶

Effect sizes of each of the analyses performed and interpreted based on the classification proposed by Maroco,¹⁷ Kinnear and Gray,¹⁸ and Cohen.¹⁹

Results

Most pregnancies were planned (65.1%) and desired (97.2%). Most mothers had prenatal care (98.2%) and a normal pregnancy (79.8%), lasting 37 weeks or more (94.7%, $M = 39.34$, $SD = 1.70$). Most babies were born by vaginal delivery (62.3%), and 37.7% were born by cesarean section, with most mothers receiving epidural anesthesia (73.4%). Apgar score values ranged from 3-10 ($M = 8.96$, $SD = 1.06$) at the first minute of life (97.2% ≥ 7) and between 7-10 ($M = 9.74$, $SD = 0.57$) at the fifth minute. At birth, weight varied between 1.990 kg and 4.530 kg ($M = 3.31$, $SD = 0.45$), and most had a birth weight ≥ 2.500 kg (99.1%), whereas height ranged from 36.00-59.05 cm ($M = 48.85$, $SD = 3.06$), and head circumference ranged between 30-37 cm ($M = 34.64$, $SD = 1.22$). Most babies did not require reanimation at birth (97.2%) and had no health problems (95.4%). Most had a pattern of biphasic sleep (59.6%). Most mothers breastfed (81.7%) for a period of time ranging from one month to 22 months ($M = 7.29$, $SD = 5.29$).

Maternal age ranged between 19 and 45 years ($M = 33.17$, $SD = 5.88$) and paternal age ranged between 20 and 50 years ($M = 36.29$, $SD = 6.03$). The sociodemographic characteristics are shown in Table 1.

Babies were aged between 6 and 22 months, with most being aged ≥ 12 months ($M = 14.50$, $SD = 4.62$). Approximately half were males (50.5%; Table 1).

Association between Maternal and Paternal Nicotine Dependence (FTND) and Child Development Quotient

There was no statistically significant correlation between the mother's nicotine dependence and cigarette smoking (FTND) and the postural, visual-motor, language, social, and overall child development quotient, with small effect size ($r^2 \leq 0.1$). There was no statistically significant correlation between the mother's morning smoking (FTND) and the postural, visual-motor, language, social, and overall child development quotient, with small effect size ($r^2 \leq 0.1$). However, there was a negative and marginally significant correlation between the mother's morning smoking (FTND) and the child's language development quotient, with medium-sized effect size ($r^2 = 0.11$) (Table 2). Thus, a higher degree of maternal morning smoking was associated with a lower language development quotient.

There was no statistically significant correlation between nicotine dependence, the morning smoking and cigarette smoking (FTND) of the father, and the postural, visual-motor, social, and overall child development quotient, with small effect size ($r^2 \leq 0.1$). However, there was a negative and statistically significant correlation between paternal nicotine dependence, smoking and morning smoking (FTND), and the child's language development quotient, with small effect size ($r^2 \leq 0.1$), for nicotine dependence and medium-sized ($r^2 = 0.15$) for morning smoking (Table 2). Thus, higher paternal nicotine dependence, cigarette smoking and morning

Table 1 Sociodemographic data.

	Mother (n = 109) %	Father (n = 109) %
Age		
< 36	58.7	45.0
≥ 36	41.3	55.0
Years of schooling		
< 9	20.2	31.2
≥ 9	79.8	68.8
Marital status		
Married	82.6	82.6
Single	11.9	11.9
Separated/Divorced	5.5	5.5
Professional status		
Student	1.8	-
Employed	74.3	82.6
Unemployed	19.3	17.4
Homemaker	4.6	-
Physical diseases		
Yes	1.8	7.3
No	98.2	92.7
Psychological diseases		
Yes	0.9	-
No	99.1	100
Medical/psychological treatment		
Yes	7.3	5.5
No	92.7	94.5
Number of children		
1	40.4	40.4
2	35.8	38.5
3 or more	23.8	21.1
	M (SD)	
Number of previous pregnancies	2.23 (1.27)	
Number of previous miscarriages	0.27 (0.63)	

smoking were associated with a lower language development quotient.

Impact of Maternal and Paternal Nicotine Dependence (FTND) on the Child's Development Quotient

The maternal nicotine dependence factor (FTND < 7 vs. FTNP ≥ 7) had a statistically significant effect on the multivariate composite of child development quotient, $\lambda = 0.91$, $F(4, 104) = 2.59$, $p = 0.041$, $\eta_p^2 = 0.09$, with the eta value suggesting a medium-sized effect. Univariate analysis showed that children of mothers without nicotine dependence had a higher mean visual-motor development, $F(1, 107) = 4.61$, $p = 0.034$ and language quotient $F(1, 107) = 5.51$, $p = 0.021$, than children of mothers with nicotine dependence, with the eta value suggesting small effect size (Table 3). There were no significant differences between children of mothers with nicotine dependence vs. children of mothers with no nicotine dependence in terms of overall development

quotient, $t (107) = 1.16$, $p = 0.248$; however, Cohen's d-value suggests a medium-sized effect (Table 3). The children of mothers with no nicotine dependence had a higher mean overall development quotient than children of mothers with nicotine dependence.

Paternal nicotine dependence factor (FTND < 7 vs. FTNP ≥ 7) did not have a statistically significant effect on the multivariate composite of child development quotient, $\lambda = 0.99$, $F(4, 104) = 0.32$, $p = 0.865$, $\eta_p^2 = 0.01$, with the eta value suggesting a small effect size. There were no significant differences between children of fathers with nicotine dependence vs. children of fathers with no nicotine dependence in terms of overall development quotient, $t (107) = 0.16$, $p = 0.875$, with Cohen's d-value suggesting a small effect size (Table 3).

Association between Maternal and Paternal Nicotine Dependence (FTND) and Child Development Quotient

There was no significant association between the overall development quotient of the child and the maternal nicotine dependence (FTND < 7 vs. FTNP ≥ 7), $\chi^2(2) = 1.56$, $p = 0.459$, and the paternal dependence, $\chi^2(2) = 1.01$, $p = 0.605$, with small effect size (Table 4).

Discussion

This study was conducted to analyze the association between parental consumption of tobacco and child development. The results show that there was such an association, with parental daily consumption pattern, nicotine dependence, and the urgency of consumption after a nocturnal abstinence period demonstrating association with language difficulties; there was an especially significant association between maternal nicotine dependence and language, visual-motor, and global development. These results are especially important because they show that the effect of parental nicotine dependence is harmful not only during pregnancy, as other studies had demonstrated¹⁰ but also after delivery. In fact, this study does not clarify the differential effect of prenatal or postnatal maternal nicotine dependence on child development. However, when considering the effect of paternal smoking dependence on child development, it can be assumed that this refers to the postnatal period, which draws attention to the vulnerability of the newborn to passive exposure to parental smoking. The mechanisms that explain the effect of tobacco consumption during pregnancy on the fetus have been well studied in recent years, and there is some evidence indicating that the consumption of nicotine and other tobacco components influence gestational duration,²⁰ have toxic effects on fetal brain development during pregnancy,¹⁰ and are associated with fetal hypoxia, changes in the serotonin uptake, changes in the dopaminergic systems, and changes in DNA and RNA synthesis in the brain.¹⁷ Nicotine appears to target specific neurotransmitter receptors in the fetal brain, causing abnormalities in cell proliferation and differentiation, resulting in cell number deficits and changes in synaptic activity,²¹ impairing the fetal-placental development and metabolism²² with developmental implications

Table 2 Association between nicotine dependence, cigarette consumption, morning tobacco consumption, and child development quotient.

Development quotient	Nicotine dependence			Cigarette consumption			Morning smoking		
	r	p	r ²	r	p	r ²	r	p	r ²
MOTHER (n = 109)									
Postural	-0.17	0.399	-0.01	-0.13	0.514	-0.02	-0.26	0.196	0.03
Visual-Motor	-0.09	0.650	-0.03	-0.06	0.785	-0.04	-0.18	0.382	-0.01
Language	-0.29	0.143	0.05	-0.25	0.215	0.02	-0.37	0.055	0.11
Social	-0.10	0.621	-0.03	-0.05	0.792	-0.04	-0.23	0.241	0.02
Overall	-0.14	0.486	-0.02	-0.09	0.661	-0.03	-0.27	0.176	0.04
FATHER (n = 109)									
Postural	0.02	0.896	-0.02	0.01	0.943	-0.02	-0.02	0.899	-0.02
Visual-Motor	0.03	0.841	-0.02	0.01	0.955	-0.02	0.04	0.808	-0.02
Language	-0.31	0.035	0.08	-0.30	0.043	0.07	-0.41	0.005	0.15
Social	-0.22	0.144	0.03	-0.27	0.072	0.05	-0.16	0.298	0.00
Overall	-0.13	0.389	-0.01	-0.15	0.305	0.00	-0.15	0.320	0.00

r^2 , size effect coefficient determinant.

Pearson's correlation test.

Table 3 Parental nicotine dependence and child development quotient.

Development quotient	MOTHER				FATHER			
	Without nicotine dependence (FTND < 7) (n = 82) M (SD)	With nicotine dependence (FTND ≥ 7) (n = 27) M (SD)	p	η_p^2	Without nicotine dependence (FTND < 7) (n = 82) M (SD)	With nicotine dependence (FTND ≥ 7) (n = 27) M (SD)	p	η_p^2
Postural ^a	97.61 (16.54)	98.04 (17.49)	0.909	0.00	98.79 (17.36)	96.24 (15.81)	0.433	0.01
Visual-motor ^a	91.29 (12.20)	85.59 (11.19)	0.034	0.04	90.71 (12.71)	88.74 (12.19)	0.405	0.01
Language ^a	84.55 (16.36)	76.07 (15.96)	0.021	0.05	83.16 (15.94)	81.48 (17.60)	0.604	0.00
Social ^a	92.20 (12.96)	88.00 (12.67)	0.145	0.02	92.33 (13.42)	89.54 (12.55)	0.269	0.01
Overall ^b	91.50 (14.27)	88.00 (11.14)	0.248	0.27	90.81 (15.24)	90.39 (11.12)	0.875	0.03

FTND, Fagerström Test for Nicotine Dependence; M, mean; SD standard deviation; η_p^2 , eta squared-size effect for MANOVA; d, Cohen's d-value-size effect for t-test.

^a MANOVA.

^b t-test for independent samples.

Table 4 Association between parental nicotine dependence and child development quotient.

Development Quotient	MOTHER				FATHER			
	Without nicotine dependence (FTND < 7) (n = 82) %	With nicotine dependence (FTND ≥ 7) (n = 27) %	V	Without nicotine dependence (FTND < 7) (n = 63) %	With nicotine dependence (FTND ≥ 7) (n = 46) %	V		
			.12					.10
Lower DQ ^a	7.3	7.4		7.9	6.5			
Normal ^a DQ ^a	70.7	81.5		69.9	78.3			
Upper DQ ^a	22.0	11.1		22.2	15.2			

DQ, development quotient; FTND, Fagerström test for nicotine dependence; V, Cramer's V-size Effect for the chi-squared test.

^a χ^2 test.

for the newborn.²³ As a consequence, there are significant changes in the brain physiology responsible for basic perceptual skills;²⁴ loss of central nervous system cells in the postnatal period; reduction of the frontal lobe and cerebellar volumes responsible for attention, emotion, and impulse control;⁸ as well as a smaller head circumference in newborns.⁹ These physiological alterations may be responsible for developmental disorders found in these children, such as self-regulatory difficulties, increased excitability,⁵ decrease in cognitive functions, and learning and memory deficits.^{10,11} Overall developmental difficulties observed in this study, with emphasis on language and visual-motor skills, are consistent with previous studies that indicate an association between parental consumption of tobacco and child development disorders,²⁵ especially psychomotor disorders.²⁶ Regarding the effect of tobacco consumption by the parents, the effect is caused by exposure after birth, which indicates the newborn's specific vulnerability to exposure to toxins released by tobacco consumption.²⁷ Because of the implications, parental smoking during the perinatal period is an important public health problem, and therefore intervention programs in perinatal health that target smoking cessation, not only for future mothers but also for future fathers, should be considered a priority to ensure a better quality of life for families.

Despite the limitations of this study—especially because it did not consider a detailed analysis of the behavior associated with tobacco consumption or regarding parental consumption history—it should be considered an important contribution to the study of the effect of parental smoking, as it helps to better understand the effect of consumption not only by mothers, but also by fathers, thus encompassing both parents in order to analyze the individual effect on psychomotor development.

Future studies should assess not only the differential effect of consumption by each parent, but also the differential effect of prenatal and postnatal tobacco consumption on child development.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgments

The authors would like to thank the Regional Director of Education for allowing the study to be carried out in Região Autónoma da Madeira (RAM) day care centers. The authors are also very grateful to the heads of educational establishments, teachers, and kindergarten assistants, and the parents who allowed the observation of the psychomotor development of their babies and spent part of their precious time to answer the questionnaire.

This study was developed as part of the project “The role of genotype-environment interaction on the resilience and vulnerability to developmental and mental health problems in the first 18 months of life” (PTDC/PSI-PCL/119152/2010) funded by the Fundação para a Ciência e Tecnologia (Foundation for Science and Technology).

References

- Willrich A, Azevedo CC, Fernandes JO. Desenvolvimento motor na infância: influência dos fatores de risco e programas de intervenção. *Rev Neurocienc.* 2008;17:51–6.
- Blanchard KA, Sexton CC, Morgenstern J, McVeigh KH, McCrady BS, Morgan TJ, et al. Children of substance abusing women on federal welfare: implications for child well-being and TANF policy. *J Hum Behav in the Social Env.* 2005;12:89–110.
- Wells K. Substance abuse and child maltreatment. *Pediatr Clin North Am.* 2009;56:345–62.
- Bridgehead International Europe quitting - progress and pathways. London: Pfizer; 2011March. Report No.USC1184.
- Stroud L, Paster R, Papandonatos G, Niaura R, Salisbury A, Battle C, et al. Maternal smoking during pregnancy and newborn neurobehavior: effects at 10 to 27 days. *J Pediatr.* 2009;154:10–6.
- Agrawal A, Scherrer J, Grant J, Sartor C, Pergadia M, Duncan A, et al. The effects of maternal smoking during pregnancy on offspring outcomes. *Prev Med.* 2010;50:13–8.
- Chertok I, Luo J, Anderson R. Association between changes in smoking habits in subsequent pregnancy and infant birth weight in West Virginia. *Matern Child Health J.* 2011;15:249–54.
- Ekblad M, Korkeila J, Parkkola R, Lapinleimu H, Haataja L, Lehtonen L, et al. Maternal smoking during pregnancy and regional brain volumes in preterm infants. *J Pediatr.* 2010;156:85–90.
- Kallen K. Maternal smoking during pregnancy and infant head circumference at birth. *Early Hum Dev.* 2000;58:197–204.
- Kiechl-Kohlendorfer U, Ralser E, Pupp U, Reiter G, Griesmaier E, Trawöger R. Smoking in pregnancy: a risk factor for adverse neurodevelopmental outcome in preterm infants? *Acta Paediatr.* 2010;99:1016–9.
- Wehby G, Prater K, McCarthy A, Castilla E, Murray J. The impact of maternal smoking during pregnancy on early child neurodevelopment. *J Hum Cap.* 2011;5:207–54.
- Heatherton TF, Kozlowski LT, Frecker RC, Fagerström KO. The Fagerström Test for Nicotine Dependence: a revision of the Fagerström Tolerance Questionnaire. *British Journal of Addiction.* 1991;86:1119–27.
- Ferreira PL, Quintal C, Lopes I, Taveira N. Teste de dependência à nicotina: validação linguística e psicométrica do teste de Fagerström. *Dependência Tabágica.* 2009;27:37–56.
- Fagerström KO, Kunze M, Schoberberger R, Breslau N, Hughes JR, Hurt RD, et al. Nicotine dependence versus smoking prevalence: comparisons among countries and categories of smokers. *Tobacco Control.* 1996;5:52–6.
- Brunet O, Lézine I. Le développement psychologique de la première enfance. Paris: Presses Universitaires de France; 1951.
- Martins C. Manual de análise de dados quantitativos com recurso ao IBM SPSS: saber decidir, fazer, interpretar e redigir. Braga: Psiquilibrios; 2011.
- Maroco J. Análise estatística com utilização do SPSS. Lisboa: Edições Sílabo; 2007.
- Kinnear PR, Gray CD. PASW statistics 17 made simple. New York: Psychology Press; 2010.
- Cohen J. Statistical power analysis for the behavioral sciences. New Jersey: Lawrence Erlbaum Associates, Inc; 1988.
- Mello PR, Pinto GR, Botelho C. Influência do tabagismo na fertilidade, gestação e lactação. *J Pediatr (Rio J).* 2001;77:257–64.
- Slotkin TA. Fetal nicotine or cocaine exposure: which one is worse? *JPET.* 1998;285:931–45.
- Jauniaux E, Biernaux V, Gerlo E, Gulbis B. Chronic maternal smoking and cord blood amino acid and enzyme levels at term. *Obstet Gynecol.* 2001;97:57–61.
- Cornelius M, Day N. Developmental consequences of prenatal tobacco exposure. *Curr Opin Neurol.* 2009;22:121–5.

24. Key A, Ferguson M, Molfese F, Peach K, Lehman C, Molfese V. Smoking during pregnancy affects speech-processing ability in newborn infants. *Environ Health Perspect.* 2007;115: 623–9.
25. Poole-Di Salvo E, Liu YH, Brenner S, Weitzman M. Adult household smoking is associated with increased child emotional and behavioral problems. *J Dev Behav Pediatr.* 2010;31:107–15.
26. Andraca I, Pino P, La Parra A, Rivera F, Castillo M. Factores de riesgo para el desarrollo psicomotor em lactentes nacidos em óptimas condiciones biológicas. *Rev Saude Publica.* 1998;32:138–47.
27. Lando HA, Hipple BJ, Muramoto M, Klein JD, Prokhorov AV, Ossip DJ, et al. Tobacco is a global paediatric concern. *Bull World Health Organ.* 2010;88:2.