

Effect of Picloram, Additives and Plant Growth Regulators on Somatic Embryogenesis of *Phyla nodiflora* (L.) Greene.

Abdul Bakrudeen Ali Ahmed^{1,3*}, Adhikarla Suryanarayana Rao², Mandali Venkateswara Rao¹ and Rosna Mat Taha³

¹Department of Plant Science; School of Life Sciences; Bharathidasan University; 620 024, Tiruchirappalli – India.

²Department of Biotechnology; School of Life Sciences; Bharathidasan University; 620 024; Tiruchirappalli – India. ³Institute of Biological Sciences; Faculty of Science; University of Malaya; 50603; Kuala Lumpur - Malaysia

ABSTRACT

The present study describes the plant regeneration via somatic embryogenesis in suspension culture derived from the leaf and stem explants of *Phyla nodiflora*. The medium type, plant growth regulators, complex extract (coconut milk and malt extract) and anti-oxidant (activated charcoal, ascorbic acid, Polyvinylpyrrolidone and citric acid) markedly influenced the embryo regeneration of *P. nodiflora*. MS with 2,4-D and activated charcoal (10 mg/L) gave the highest stimulation of embryogenic callus growth. Optimized callus was transferred into suspension culture, which showed the globular, heart shaped embryos in MS with 2,4-D + BA + picloram (0.1 mg/L), coconut milk (10 ml/L), citric acid (100 mg/L) on 6th subcultures. Further development stages such as torpedo and cotyledonary stage embryos and fostered maturation of embryos were observed at 8th and 10th subculture. However, the high frequency embryo germination and plantlet (45 plants/20 mg cotyledonary stages embryos) formation was obtained in half-strength MS medium without growth regulators from cotyledonary embryos. All the plantlets established in the field exhibited morphological characters similar to those of the mother plant.

Key words: Somatic embryogenesis (SEs); picloram; ascorbic acid; coconut milk; plant growth regulators; Murashige and Skoog medium

INTRODUCTION

Phyla nodiflora L. Greene (= *Lippia nodiflora* (L.) Miq.) belongs to Verbenaceae family, distributed in India, Ceylon, Baluchistan, South Africa and Central America (Terblanche and Kornelin, 1996). Its leaves are eaten in Ceylon and taken as tea in the Philippines. It is an aromatic, runner plant with scanty roots and cures adenopathy, chronic indolent ulcers, diuretic and aphrodisiac and is also used for the treatment of heart diseases, ulcers, bronchitis, fevers, and colds (Kirtikar and Basu, 1975). The plant is also used for the boils, indigestion in

children, and by the women after the delivery (Nadkarni, 1954; Chopra *et al.*, 1956). Ravikanth *et al.*, (2000) reported that the anticancer compounds (halleridone and hallerone) from *P. nodiflora*.

It also has the alkaloids and shows the significant analgesic, anti-inflammatory and anti-pyretic activities (Costa *et al.*, 1989, Forestieri *et al.*, 1996), anti-cancer, anti-tumor, anti-malarial, antifungal, cytotoxic activity (Nishino *et al.*, 1988), antioxidant and free radical scavenging effect (Ashokkumar *et al.*, 2008), antioxidant

* Author for correspondence: bakru24@yahoo.co.in

activities (Shukla *et al.* 2009) and cure multiple skin disease (Abbasi *et al.*, 2010).

The commercial exploitation of the medicinal plants for the production and conventional propagation is hampered due to their poor seed viability, low rate of germination and poor rooting ability of the vegetative cuttings. The alternative propagation methods could be beneficial for accelerating the large scale multiplication and the conservation of the medicinal plant. Since, less *in vitro* studies have been done in this genus, latest studies on the propagation of *Lippia junelliana* (Juliani *et al.*, 1999) and *Lippia alba* (Gupta *et al.*, 2001). Direct shoot propagation using different explants has been implemented (Bhatt *et al.*, 2002; Ahmed *et al.*, 2005). Somatic embryogenesis is an alternative method for the large-scale propagation method. However, for this, there is lack of information for the embryo induction process (Dodeman *et al.*, 1997). The aim of this work was to study the germination capability and development of somatic embryos (SEs) from *Phyla nodiflora*.

MATERIAL AND METHODS

Plant material and inoculation

Phyla nodiflora young plants were collected and maintained in Department of plant science garden, Bharathidasan University, Tiruchirappalli, Tamil Nadu, India. Two years old young leaf and stem explants were washed in running tap water, surface disinfected in a solution of HgCl_2 (0.1 %, w/v) containing distilled water for 2 min and finally rinsed with sterile distilled water for several times. The explants were inoculated into the MS (Murashige and Skoog, 1962); SH (Schenk and Hildebrandt, 1972), WPM (Lloyd and McCown, 1980) and B5 (Gamborg *et al.*, 1968) medium.

Embryogenic callus induction and maturation

The leaf and stem explants were inoculated into the medium with plant growth regulators and antioxidant. Both the explants were cultured on different media for embryogenic callus induction: namely MS, SH, B5 and WPM supplemented with sucrose (3 %, w/v), combination with 2,4-D: 0.2 - 1.4 mg/L, NAA: 0.2 - 1.4 mg/L with ascorbic acid (10 mg/L) respectively. For SEs maturation, auxins with picloram (0.01 - 0.2 mg/L), cytokinins

(BA: 0.1 - 2.0 mg/L), (KN: 0.1 - 2.0 mg/L), coconut milk (10 ml/L) and citric acid (50 -150 mg/L) were tested in the suspension culture. The media pH was adjusted to 5.8 (1N NaOH / HCl) and after adding the agar (0.8 %, w/v) autoclaved at 121 °C for 15 min. The callus tissue was weighed and placed into 125 ml Erlenmeyer flask containing 60 ml basal medium without agar. Matured cotyledonary embryos were used for plantlets formation in free MS solid medium. All SEs cultures were maintained at 25 ± 2 °C under a photoperiod (16 h/8 h) with light intensity $35 \mu \text{m}^{-2} \text{s}^{-1}$ and 55-60 % relative humidity. The germinated plantlets were transplanted to plastic pots containing vermiculite supplemented with the nutrient solution (NPK 17:17:17) at weekly intervals.

Statistical analysis

Only data which showed advantageous effect were included in the tables and presented in mean of explants per treatment and repeated three times. Thirty replicates were used and repeated thrice. Experimental design was completely random and factorial with callus initiation, globular, heart, torpedo and cotyledonary stages of the callus. The data were subjected to analysis of variance and mean separation was carried out using Duncan's Multiple Range Test (DMRT) at 5% level significance (Gomez and Gomez, 1976).

RESULTS AND DISCUSSION

Callus initiation

In all media, the callus initiation didn't occur without the growth regulators (control) in leaf and stem explants (data not shown). The suitable embryogenic callus induction was observed in 2,4-D and NAA with ascorbic acid in MS, SH, B5 and WPM (Table 1). However in B5, SH and WPM media, the embryogenic potential significantly decreased (data not shown). Callus initiation and proliferation was better in 2,4-D and NAA than IAA, IBA in all media (data not shown). Between the two auxins, 2,4-D (0.6 mg/L) with ascorbic acid (10 mg/L) induced higher embryogenic callus in leaf explants (94.5 %; Fig. 1A) than stem explants (76.8 %, Fig. 1 B) after 25 days (Table 1).

Table 1 - Embryogenic callus induction from leaf and stem explants of *Phylla nodiflora* on B5, MS, SH, WPM medium supplemented with 2,4-D and NAA, after 25 days.

Medium type / Plant growth regulators / Explants		Embryogenic callus frequency (%)													
		NAA (mg/L) + Ascorbic acid (10 mg/L)							2,4-D (mg/L) + Ascorbic acid (10 mg/L)						
		0.2	0.4	0.6	0.8	1.0	1.2	1.4	0.2	0.4	0.6	0.8	1.0	1.2	1.4
B5	Leaf	32.5 ^d	37.6 ^d	40.5 ^{cd}	52.8 ^c	44.6 ^{cd}	28.5 ^d	22.4 ^d	21.4 ^d	28.4 ^d	30.8 ^d	39.2 ^d	28.5 ^d	22.6 ^d	18.4 ^d
MS	Leaf	49.5 ^a	63.2 ^a	71.6 ^a	78.5 ^a	63.0 ^a	52.4 ^a	40.6 ^a	50.4 ^a	65.6 ^a	94.5 ^a	71.3 ^a	62.6 ^a	56.0 ^a	39.2 ^a
SH	Leaf	38.5 ^{bc}	45.6 ^b	54.2 ^b	65.4 ^b	54.1 ^b	39.5 ^b	32.5 ^b	35.6 ^b	38.2 ^b	42.4 ^{bc}	45.5 ^{bc}	42.5 ^{bc}	40.0 ^b	34.6 ^{ab}
WPM	Leaf	39.2 ^b	40.0 ^{bc}	43.5 ^c	47.3 ^{cd}	45.2 ^c	32.2 ^{bc}	26.5 ^{bc}	34.2 ^{bc}	36.6 ^{bc}	45.5 ^b	51.6 ^b	48.5 ^b	35.4 ^{bc}	28.0 ^c
B5	Stem	24.6 ^{bc}	29.0 ^{bc}	34.5 ^b	40.2 ^b	36.8 ^b	30.0 ^b	26.5 ^b	30.6 ^b	36.2 ^b	52.2 ^b	41.8 ^b	39.6 ^{ab}	35.0 ^{ab}	26.8 ^b
MS	Stem	30.5 ^a	40.6 ^a	55.8 ^a	59.5 ^a	48.0 ^a	42.6 ^a	39.0 ^a	35.8 ^a	42.5 ^a	76.8 ^a	51.8 ^a	42.6 ^a	39.4 ^a	31.0 ^a
SH	Stem	26.0 ^b	29.3 ^b	32.2 ^{bc}	34.6 ^{cd}	33.2 ^{bc}	26.8 ^{bc}	20.2 ^{bc}	21.0 ^d	24.3 ^d	28.2 ^{cd}	31.6 ^d	26.2 ^{cd}	22.8 ^{cd}	18.2 ^{cd}
WPM	Stem	20.6 ^d	28.4 ^d	30.2 ^d	35.8 ^c	28.5 ^d	23.6 ^d	19.2 ^d	30.0 ^{bc}	34.2 ^{bc}	36.8 ^c	38.2 ^{bc}	30.2 ^c	26.8 ^c	22.2 ^c

Values are mean of 30 replicates per treatment and repeated thrice. Values with the same superscript are not significantly different at 5% probability level according to DMRT.

Somatic embryos maturation in suspension culture

Embryogenic tissue was maintained and bulked up through secondary somatic embryogenesis. In order to stimulate SE maturation, the pieces of embryogenic tissue (15 to 20 mg), which considered of immature embryos, were transferred into each conical flask containing suspension maturation medium supplemented with picloram and additives. The suspension culture was superior in embryogenic callus maturation than semisolid culture of leaf explants. In order to determine the effect of PGRs on somatic embryogenesis in *P. nodiflora*, the highest frequency embryos at the globular stage (89.8 %), heart stage (74.9 %) were observed onto media supplemented with 2,4-D (0.6 mg/L) + BA (1.0 mg/L) + picloram (0.1 mg/L) + coconut milk (10 ml/L) with citric acid (100 mg/L) in 40 and 60 days. During the embryos maturation, the suspension culture was continuously subculture every week to prevent the re-callus and phenolic excretion in the medium.

Torpedo and cotyledonary stages in suspension culture

The advantage of suspension culture was that the large number of free single cells were aggregated and it could be easily identified from the undifferentiated and differentiated cells (developmental stages) of SEs (Fig. 1C-I). However, the abnormal embryos were trumpet-shaped and didn't show any further development. Torpedo and cotyledonary stages embryos development were observed in MS medium with

2,4-D (0.6 mg/L) + BA (1.0 mg/L) + picloram (0.1 mg/L) + coconut milk (10 ml/L), citric acid (100 mg/L) to form torpedo (62.6 %) and cotyledonary stage embryos (55.2 %) in suspension culture (Table 2).

Germination and field survival

The mature cotyledonary embryos were transferred to half-strength MS medium without growth regulators which showed the increase frequency of plantlets than full strength medium (Fig. 2). In germination experiment, a total of 100 mature embryos (20 mg) were selected on placed on the germination media. These embryos were responded differently on half-strength and full strength germination media. 52 % of these embryos were showed root emergence in half strength MS medium, which included root and shoot in 140 days. While 26 % (100 days) and 40 % (120 days) of them demonstrated the shoot and root initiation. About 160 days (44 %) of all developed embryos were significantly reduced the germination nature without hardened. Significantly reduced abnormal embryos were observed in full-strength MS medium (data not shown). The SEs germination in the present study could be due to the persistence of auxins, which might also be the cause of dedifferentiation of embryos. However, in half-strength medium the plantlets were recovered (45 plantlets / 20 mg cotyledonary embryo callus) in 120 days. The germinated plantlets were individually transferred to the pots containing soil, sand and farmyard manure (1:1:1) and were reared in the green house (Fig. 1K).




Figure 1 - Plant regeneration from leaf and stem explants of *Phyla nodiflora* (L.) Greene via somatic embryogenesis

A – Embryogenic callus (leaf explant); B – Embryogenic callus (Stem explant); C – Different stages embryos (leaf explants); D – Globular shaped embryo; E – Heart shaped embryo; F and G – Torpedo stage embryo; H – Late torpedo stage embryo; I – Early cotyledonary stage embryo; J – Embryo germination; K – Germinated plantlets maintained in the field.

Table 2 - Somatic embryogenesis from callus induced from leaf and stem explants of *Phyla nodiflora* on MS medium supplemented with 2,4-D, picloram and BA in suspension culture at different days

Plant growth regulators (mg/L)			Globular (%) 4 th subculture	Heart (%) 6 th subculture	Torpedo (%) 8 th subculture	Cotyledonary (%) 10 th subculture
Coconut milk (10 ml/L)	+ Citric acid (100 mg/L)	BA	40 days	60 days	80 days	90 days
2,4-D	Picloram	BA				
0.6	0.01	0.5	72.6 ± 1.7 ^c	61.3 ± 1.5 ^{bc}	49.5 ± 1.0 ^c	36.2 ± 1.4 ^{bc}
0.6	0.1	0.5	65.2 ± 1.3 ^{ef}	56.4 ± 2.1 ^{de}	38.8 ± 1.5 ^{ef}	29.6 ± 2.0 ^f
0.6	0.2	0.5	59.1 ± 0.9 ^g	52.8 ± 2.5 ^f	32.6 ± 3.2 ^h	25.8 ± 1.0 ^h
0.6	0.01	1.0	78.5 ± 2.1 ^b	59.6 ± 1.8 ^d	46.1 ± 2.1 ^{cd}	38.4 ± 1.7 ^b
0.6	0.1	1.0	89.8 ± 2.8 ^a	74.9 ± 0.6 ^a	62.6 ± 1.8 ^a	55.2 ± 2.2 ^a
0.6	0.2	1.0	71.4 ± 1.5 ^{cd}	64.6 ± 2.2 ^b	56.4 ± 1.3 ^b	32.5 ± 1.9 ^d
0.6	0.01	1.5	69.7 ± 1.4 ^e	52.3 ± 1.3 ^{fg}	41.2 ± 1.4 ^e	31.0 ± 0.9 ^{de}
0.6	0.1	1.5	53.2 ± 1.7 ^h	47.6 ± 1.9 ^h	37.9 ± 1.6 ^g	27.8 ± 1.3 ^{fg}
0.6	0.2	1.5	46.0 ± 2.1 ⁱ	34.1 ± 2.0 ⁱ	29.5 ± 2.1 ^{hi}	18.2 ± 2.2 ⁱ

Values are mean of 30 replicates per treatment and repeated thrice. Values with the same superscript are not significantly different at 5% probability level according to DMRT.




Figure 2 - Somatic embryos germination from cotyledonry stage embryos of *Phyla nodiflora* in different days.

DISCUSSION

Influence of medium and plant growth regulators

When the leaf and stem explants were immersed into B5, SH, WPM media with plant growth regulators and ascorbic acid, the somatic embryos callus initiation was significantly reduced, even if they were maintained for longer period in the culture. Zimmerman, (1993) reported that the pro-embryogenic callus were containing auxins to synthesize all the necessary genes to complete the globular stage. However, the auxins were removed from the culture to make inactive genes or synthesize new gene products for the completion of embryo development. Kawahara and Komamine (1995) reported that the exogenous auxins were involved in gene expression of early stages of somatic embryogenesis. However, Wang *et al.* (2006) reported that the NAA (0.2 mg/L) and 2,4-D (0.2 mg/L) induced the embryogenic callus in *Chorispora bungeana*; similar results were observed in *Phoenix dactylifera* (Fki *et al.*, 2003, Lin *et al.*, 2004). In other plant species, the 2,4-D influenced the embryo induction and participation at initial stages of development (Gray *et al.*, 1993; Mujib and Samaj, 2006; Junaid *et al.*, 2007; Sharma *et al.*, 2007). In the present study work in *P. nodiflora* NAA was less effective compared to 2,4-D.

Influence of additives and coconut milk in somatic embryos maturation

Successful somatic embryogenesis was after obtained in the optimum concentration of auxins combined with cytokinins, picloram, additives (citric acid) and coconut milk. The activated charcoal, polyvinylpyrrolidone and malt extracts, however significantly reduced the somatic embryos quality (data not shown). Guo and Zhang (2005) reported more frequency of SEs maturation in MS medium supplemented with 2,4-D (0.2 mg/L) + BA (5.0 mg/L) in *Zingiber officinale*. Hence, tests were conducted without picloram and additives in addition to plant growth regulators, which showed that the SEs maturation was significantly reduced (data not shown). Firoozabady and Moy (2004) reported picloram as one of embryogenic potentials agent to increase the growth regulators in *Ananas comosus*. However, picloram regulated the embryogenic stages and produced maximum frequency of SEs and plant germination (Little *et al.*, 2000; Groll *et al.*, 2001). Somatic embryos maturation was stimulated by auxins combine with cytokinins in *Leptadenia reticulata* (Martin, 2004).

Influence of strength media in plant germination

The suspension culture derived from the cotyledonary embryos were transferred to full and

half strength medium with auxins. Somatic embryos were exposed to too much auxin during the development, but failed to accumulate the storage protein and germinate at a lower frequency (Stuart *et al.*, 1984). The present results showed that somatic embryos germinated without auxins. However, SEs maturation and germination were critical steps for the recovery of healthy plants (Ramanjini and Prakash, 1998). Hwang (2006) reported that 87 % SEs successfully developed into plantlets on ½ MS medium without growth regulators after six weeks culture in *Abelmoschus manihot*.

In conclusion, somatic embryos were induced from the immature zygotic embryos by picloram in suspension culture. The best embryo callus induction in MS medium with 0.1 mg/L picloram in the leaf explants were optimum for torpedo and cotyledonary stage embryos. The somatic embryo development total process was completed in 160 days. This efficient somatic embryos protocol could be useful for conservation and agronomy and in the improvement of *P. nodiflora* using gene transfer biotechnologies.

REFERENCES

- Abbasi, A.M., Khan, M.A., Ahmed, M., Zafar, M., Jahan, S., Sultana, S (2010), Ethnopharmacological application of medicinal plants to cure skin diseases and in folk cosmetics among the tribal communities of North-West Frontier Province, Pakistan. *J. Ethnopharmacol.*, **128**, 322-335.
- Ahmed, A.B.A., Gouthaman, T., Rao, A.S., Rao, M.V. (2005), Micropropagation of *Phyla nodiflora* (L.) Greene: An important medicinal plant. *Iranian J. Biotech.* **3**, 186-190.
- Ashokkumar, D., Thamilselvan, V., Senthilkumar, G.P., Mazumder, U.K., Gupta, M. (2008), Antioxidant and free radical scavenging effects of *lippia nodiflora*. *Pharmaceutical Biol.*, **46**, 762-771.
- Bhatt, T., Jain, V., Jayathiratha, M.G., Banerjee, G., Mishra, S.H. (2002), *In vitro* regeneration of roots of *Phyla nodiflora* and *Leptadenia reticulata* and comparison of roots from cultured and natural plants for secondary metabolites. *J. Exp. Biol.*, **40**, 1382-1386.
- Chopra, R.N., Nayer, S.L., Chopra, I.C. (1956), Glossary of Indian Medicinal Plants, CSIR, V edition, New Delhi, 12, p. 157.
- Costa, M., Di Stasi, L.C., Kirizawa, M., Mendacolli, S.L., Gomes, C., Trolin, G. (1989), Screening in mice of some medicinal plants used for analgesic purposes in the state of Sao Paulo. Part II. *J. Ethnopharmacol.* **27**, 25-39.
- Dodeman, V.L., Ducreux, G., Kresis, M. (1997), Zygotic embryogenesis versus somatic embryogenesis. *J. Exp. Bot.*, **48**, 1493-1509.
- Firoozabady, E., Moy, Y. (2004), Regeneration of pineapple plants via somatic embryogenesis and organogenesis. *In Vitro Cell. Dev. Biol.*, **40**, 67-74.
- Fki, L., Masmoudi, R., Drira, N., Rival, A. (2003), An optimized protocol for plant regeneration from embryogenic suspension cultures of date palm, *Phoenix dactylifera* L., cv. Deglet Nour. *Plant Cell Rep.*, **21**, 517-524.
- Forestieri, A.M., Monforte, M.T., Ragusa, S., Trovato, A., Lauk, L. (1996), Anti-inflammatory, analgesic and pyretic activity in rodents of plant extracts used in Africa medicine. *Phytothera. Resear.*, **10**, 100-106.
- Gambrog, O.L., Miller, R.A., Ojima, K. (1968), Nutrients requirements of suspension culture of soybean root cells. *Exp. Cell. Res.*, **50**, 151-158.
- Gomez, K.A., Gomez, A.A. (1976), Statistical procedures for agricultural research with emphasis on rice. International Rice Research Institute. Los Banos, Philippines, p. 264.
- Gray, D.J., McColley, D.W., Compton, M.E. (1993), High frequency embryogenesis from quiescent seed cotyledons of *Cucumis melo* cultivars. *J. Amer. Soc. Hort. Sci.*, **118**, 425-432.
- Groll, J., Mycock, D.J., Gray, V.M., Lamenski, S. (2001), Secondary somatic embryogenesis of *Cassava* on picloram supplemented media. *Plant Cell Tiss. Org. Cult.*, **65**, 201-210.
- Guo, Y., Zhang, Z. (2005), Establishment of plant regeneration of somatic embryogenic cell suspension cultures of the *Zingiber officinale* Rosc. *Scientia Hort.*, **107**, 90-96.
- Gupta, S.K., Khanuja, S.P.S., Kumar, S. (2001). *In vitro* propagation of *Lippia alba*. *Current Sci.*, **81**, 206-210.
- Hwang, S.J. (2006), Somatic embryogenesis and plant regeneration in *Abelmoschus manihot* (L.) medicus. *Prop. Orn. Plants.*, **6**, 34-38.
- Juliani, H.R., Koroch, A.R., Giuliani, H.R., Trippi, V.S. (1999), Micropropagation of *Lippia junelliana* (Mold.) Tronc. *Plant Cell Tiss. Org. Cult.*, **59**, 175-179.
- Junaid, A., Mujib, A., Bhat, M.A., Sharma, M.P., Samaj, J. (2007), Somatic embryogenesis and plant regeneration in *Catharanthus roseus*. *Biologia plant.*, **51**, 641-646.

- Kawahara, R., Komamine, A. (1995), Molecular basis of somatic embryogenesis. In: Biotechnology in Agriculture and Forestry, Somatic Embryogenesis and Synthetic Seed, Vol. 30. In: Bajaj, Y. P. S. ed. Springer-Verlag, Berlin, Heidelberg, New York, pp. 30-40.
- Kirtikar, K.R., Basu, B.D. (1975), In: Indian medicinal plants, 2nd ed. Jayeed press, New Delhi, Vol-III, 1916-1917.
- Lin, C.S., Lin, C.C., Chang, W.C. (2004), Effect of thidiazuron on vegetative tissue-derived somatic embryogenesis and flowering of bamboo *Bambusa edulis*. *Plant Cell Tiss. Org. Cult.* **76**, 75-82.
- Little, E.L., Magbana, Z.V., Parrott, W.A. (2000), A protocol for repetitive somatic embryogenesis from mature peanut epicotyl. *Plant Cell Rep.*, **19**, 351-357.
- Lloyd, G., McCown, B. (1980), Commercially feasible micropropagation of mountain laurel, *Kalmia latifolia*, by use of shoot tip culture. *Intl. Plant Prop. Soc. Proc.*, **30**, 421-427.
- Martin, K.P. (2004), Benzyl adenine induced somatic embryogenesis and plant regeneration of *Leptadenia reticulata*. *Biol. Plant.*, **48**, 285-288.
- Mujib, A., Samaj, J. (ed.) (2006), Somatic embryogenesis. Springer Verlag, Berlin-Heidelberg-New York.
- Murashige, T., Skoog, F.A. (1962), Revised media for rapid growth and bioassay with tobacco tissue cultures. *Physiol. Plant.*, **15**, 473-497.
- Nadkarni, A.K. (1954), Indian Materia Medica, Bombay: Popular Book Depot, p. 280-281.
- Nishino, C., Kobayashi, K., Fukushima, M. (1988), Halleridone, a cytotoxic constituent from *Cornus controversa*. *J. Nat Prod.*, **51**, 1281-1282.
- Ramanjini, G.P.H., Prakash, C.S. (1998), Herbicide glyphosate at sublethal concentrations enhances somatic embryo development in sweet potato (*Ipomea batatus* L.). *Current Sci.*, **75**, 508-510.
- Ravikanth, V., Ramesh, P., Diwan, P.V., Venkateswarlu, Y. (2000), Halleridone and Hallerone from *Phylla nodiflora* (L.) Greene. *Biochemical Systematic Ecology*, **28**, 905-906.
- Schenk, R.V., Hildebrandt, A.C. (1972), Medium and techniques for induction of growth of monocotyledonous and dicotyledonous plant cell cultures. *Can J. Bot.*, **50**, 166-204.
- Sharma, S.K., Bryan, G.J., Millam, S. (2007), Auxin pulse treatment holds the potential to enhance efficiency and practicability of somatic embryogenesis in potato. *Plant Cell Rep.*, **26**, 945-950.
- Shukla, S., Saluja, A.K., Pandya, S.S. (2009), *In vitro* antioxidant activity of aerial parts of *Lippia nodiflora* Rich. *Pharmacology Online*, **2**, 450-459.
- Stuart, D.A., Strickland, S.G. (1984), Somatic embryogenesis from cell cultures of *Medicago sativa* L. I. Effect of amino acid additions to the regeneration medium. *Plant Sci. Lett.*, **34**, 165-174.
- Terblanche, F.C., Kornelius, G. (1996), Essential oil constituents of the genus *lippia* (Verbinaceae) - A literature review. *Journal of Essential Oil Research*, **8**, 471-485.
- Wang J, An L, Wang R, Yang D, Si J, Fu X, Chang J, Xu S (2006), Plant regeneration of *Chorispora bungeana* via somatic embryogenesis. *In Vitro Cell. Develop. Biol.-Plant.*, **42**, 148-151.

Received: May 25, 2009;
 Revised: October 08, 2009;
 Accepted: April 22, 2010.