

Anatomical Variation in the Mandibular Foramen in Non-Atrophic and Atrophic Mandibles

Marcelo Betti Mascaro¹, Lara Cristina Picoli¹, Stefany Torres Reis Matos¹, Silvana Carla Sipos Lotائف¹, Mônica Rodrigues De Souza¹, Marcelo Ferreira Calderon¹

¹Discipline of Morphology, School of Dentistry, Nove de Julho University (UNINOVE), São Paulo, SP, Brazil.

Corresponding Author:

Marcelo Betti Mascaro

Discipline of Morphology, School of Dentistry

Nove de Julho University (UNINOVE)

Rua Vergueiro 239-245, CEP:01504-001, Liberdade, São Paulo, SP

Brazil

Phone: (55) (11) 33859164

E-mail: marcelobmascaro@gmail.com

ABSTRACT

Objectives: Previous studies of variation in mandibular foramen characteristics with age have involved comparison in different populations, but few data, between non-atrophic and atrophic mandibles are available. The aim of this original article was to compare the position, shape and area of the mandibular foramen between non-atrophic and atrophic mandibles.

Material and Methods: Morphometric methods were used to study the mandibular foramen variation. Fifty adult dry mandibles from the laboratory of anatomy were selected. Mandibles were considered non-atrophic if the distance between the base and alveolar ridge was homogeneous and greater than 25 mm in the anterior region and 20 mm in the posterior region. Conversely, mandibles were considered atrophic if that distances were lower than those described to a minimum of 11 mm in all areas. All measurements were performed with a digital caliper. For statistical analysis, the admitted level of significance was 5%.

Results: When non-atrophic mandibles were compared to atrophic ones, the mandibular foramen shifted significantly to an anterior position (mean difference [MD]: 4.81 mm; $P < 0.0001$) and to an inferior position (MD: 3.04 mm; $P < 0.0001$) and changed from an elliptical shape to round one, with a significant decrease in its area (MD: 3.66 mm²; $P < 0.05$).

Conclusions: The results indicate that there are significant differences in the position, shape and area of the mandibular foramen between non-atrophic and atrophic mandibles. These data should be considered in anaesthetic techniques and surgical procedures to prevent vascular and nervous lesions.

Keywords: anatomy; atrophy; humans; mandible; mental foramen.

Accepted for publication: 28 December 2020

To cite this article:

Mascaro MB, Picoli LC, Reis Matos ST, Sipos Lotائف SC, De Souza MR, Ferreira Calderon M.

Anatomical Variation in the Mandibular Foramen in Non-Atrophic and Atrophic Mandibles

J Oral Maxillofac Res 2020;11(4):e4

URL: <http://www.ejomr.org/JOMR/archives/2020/4/e4/v11n4e4.pdf>

doi: [10.5037/jomr.2020.11404](https://doi.org/10.5037/jomr.2020.11404)

INTRODUCTION

The mandible arises from the ossification of the embryonic tissue called ectomesenchyme [1,2]. The mandibular foramen (MF) is an opening in the medial part of the mandibular ramus [3]. It marks the entrance of the mandibular canal, which connects the mandibular and the mental foramen [4]. In the MF penetrates the artery and the inferior alveolar vein, which constitute together with the inferior alveolar nerve a neurovascular bundle [5,6]. The alveolar inferior nerve originates the mental nerve that crosses the mental foramen [6]. Mandibular atrophy that occurs predominantly after dental loss may modify the positions of the bundle [7]. The knowledge of the position of the MF and the structures that cross through that are extremely important for dental surgeons [8,9] specially when they used anaesthetic techniques like the pterygomandibular technique. It aims to block, among other nerves, the inferior alveolar nerve, before it penetrates in the MF [10]. Nervous or vascular lesions can occur during this procedure [11]. The correct position of the MF and the mandibular canal can be evaluated by panoramic radiography and mainly by mandible computed tomography. Using three-dimensional reconstruction associated with three-dimensional printing, surgical guides can be prepared. These guides will assist the surgeon in intraoperative procedures [12]. Implants must be positioned carefully in molar and premolar regions mainly in those cases with limited vertical alveolar bone what is verified in atrophic mandibles [11,12]. Moreover, these surgical guides can be used to perform an osteotomy in orthognathic surgery [12]. In these osteotomies, the inferior alveolar nerve is the structure that most suffers lesions [13-15]. Therefore, preoperative planning, early detection of sensory disturbance, and adequate treatment are fundamentals

in these surgical procedures [11,15]. This article aimed to evaluate the position, shape and area of the mandibular foramen in non-atrophic mandibles and investigate the alterations that can occur in atrophic ones.

MATERIAL AND METHODS

Ethical considerations

This project was carried out in the Human Anatomy Laboratory of the Nove de Julho University Center in Sao Paulo, Brazil, from August, 2017 to August, 2018. A letter of science and authorization was requested to the managers of the dentistry course for the use of cadaveric specimens. The present study was approved for the Ethical Committee of the University (Ethical Protocol number: 1.962.285).

Selection of the mandibles

Fifty adult dry mandibles were selected without distinction of age, sex, or ethnicity. The non-atrophic mandibles (NAT) were fully or partially dentate and the atrophic mandibles (AT) were edentulous. Mandibles were considered non-atrophic if the distance between the base and alveolar ridge was homogeneous and greater than 25 mm in the anterior region (point A, Figure 1) and 20 mm in the posterior region (points B and C, Figure 1). Conversely, mandibles were considered atrophic if the distances were lower than those described to a minimum of 11 mm in all areas. Mandibles measuring less than 11 mm were considered extremely atrophic and were not included in the analysis of this study. These measures are in agreement with the classification of mandibular atrophy degree proposed by Cawood and Howell [16] in 1988 and by Luhr et al. [17] in 1996. The measurements were performed




Figure 1. Comparison between non-atrophic and atrophic mandibles. A = points for measurement represented in a non-atrophic mandible; B = average decrease of the mandibular body represented in an atrophic mandible.

in the symphysis region (point A, Figure 1), bilaterally in the foramen mental region (point B, Figure 1) and bilaterally at a specific point in the posterior region of the mandible body (point C, Figure 1). This posterior point was determined by the mean distance between the mental foramen and the MF [16]. All linear measurements were performed with a digital caliper (150 mm capacity, 0.01 mm resolution, accuracy +/-0.03 mm - Digimess precision instruments Ltda; São Paulo, SP, Brazil) by two independent researchers (MBM, LCP). The average between the two measurements was considered [18].

Experimental procedures and data analysis

To evaluate the position of the MF in relation to ramus, straight lines from the geometric centre of the foramen were traced and measured to specific points.

The first point was the most concave point of the oblique line, what was considered the anterior margin of the ramus (AM-MF) (Figure 2). The second point was the posterior margin of the ramus (PM-MF). The third point was the most concave point of the mandibular notch, this point was considered the superior margin of the ramus (SM-MF). The fourth and last point was the inferior margin of the ramus (IM-MF). These measures are in accordance with the methodology proposed by Lima et al. [9].

The antero-posterior and latero-lateral diameter of the MF was measured to evaluate their shape and determine if it was circular or elliptical. Afterwards, a formula was applied to determine the area:

$$A = \pi \times R \times r$$

Area is equal to 3.14 times the largest radius times the smallest radius [9].




Figure 2. Mandibular foramen: position, shape and area. A = Position of the mandibular foramen to specific points of the ramus in a non-atrophic mandible; B = Average decrease of the measurements in an atrophic mandible; C = Representation of the shape of the foramen in both groups and average decrease of its area.

AM = anterior margin of the ramus; IM = inferior margin of the ramus; MF = mandibular foramen; PM = posterior margin of the ramus; SM = superior margin of the ramus.

Measurement of the gonial angle

Variations of the gonial angle will contributed to changes in MF position. The measurement of the gonial angle was performed with a goniometer at the junction of the inferior and posterior margins of the ramus of the mandibles according to the methodology proposed by Shenoy et al. [18].

Statistical analysis

The data were copied to Microsoft Office Excel program (Microsoft Corporation; Washington, USA) where they were tabulated and therefore, transferred to the GraphPad InStat v3.1 software (GraphPad Software Inc.; San Diego, California, USA). For comparison of the data between the sides of the same group or between the groups of NAT and AT mandibles, the Kolmogorov-Smirnov normality test was used, and later, when $P > 0.05$, the t-test with or without the Welch’s correction was applied. If $P < 0.05$, when the Mann-Whitney test was performed. Parametric data were expressed as mean and standard deviation (M [SD]).

RESULTS

Measurement of atrophy

There were no statistically significant differences when comparing measures obtained from the right and left sides of the mandibles in the same group. There was a highly significant difference when was compared the height of the alveolar ridge at points A, B and C between the NAT and AT mandibles (Figure 1, Table 1).

Position of the mandibular foramen

No statistical differences in the position of the MF were observed when the right and left sides of the same group were compared.

The MF in NAT mandibles was closer to the points considered as the posterior margin and superior margin of the ramus when these distances were compared to the points considered the anterior margin and inferior margin of the ramus respectively. The AT mandible demonstrated a similar pattern, but the foramen dislocated its position significantly to an anterior and inferior point. The displacement was observed because the mandible presented a great resorption of the anterior, posterior and inferior margin of the ramus, but not of the superior margin. Therefore, the foramen is more centralized mainly in the superior-inferior axis (Figure 2; Table 2).

Regarding the morphology, which in the NAT mandibles was predominantly elliptical, with an antero-posterior distance greater than latero-lateral, became circular in the AT mandibles. Moreover, the area of the MF decreased significantly between the groups (Figure 2; Table 3 and 4).

Table 1. Comparison of the height of the alveolar at points A, B and C between the non-atrophic and atrophic mandibles

Point	Non-atrophic mandibles	Atrophic mandibles
	Mean (SD), mm	Mean (SD), mm
Point A ^a	31.39 (3.21)	18.96 (2.87)
Point B right ^a	31.58 (2.47)	17.41 (2.03)
Point B left ^a	30.81 (2.44)	17.04 (2.07)
Point C right ^a	26.99 (2.55)	15.29 (2.29)
Point C left ^a	26.17 (2.32)	15.62 (2.12)

^aStatistically significant at level $P < 0.0001$ (t-test). SD = standard deviation.

Table 2. Comparison of the position of the mandibular foramen between the non-atrophic and atrophic mandibles

Distance	Non-atrophic mandibles	Atrophic mandibles
	Mean (SD), mm	Mean (SD), mm
AM-MF right ^a	19.18 (1.84)	14.54 (2.9)
PM-MF right ^a	14.69 (2.61)	11.83 (1.47)
SM-MF right ^a	23.26 (3)	21.2 (3.78)
IM-MF right ^a	25.89 (3.94)	23.6 (3.39)
AM-MF left ^a	19.48 (1.82)	14.51 (2.7)
PM-MF left ^a	14.18 (2.44)	11.73 (1.46)
SM-MF left	22.24 (3.3)	21.65 (3.58)
IM-MF left ^a	25.95 (3.44)	22.2 (3.47)

^aStatistically significant at level $P < 0.0001$ (t-test). AM = anterior margin of the ramus; IM = inferior margin of the ramus; MF = mandibular foramen; PM = posterior margin of the ramus; SM = superior margin of the ramus; SD = standard deviation.

Table 3. Comparison of the shape of the mandibular foramen between the non-atrophic and atrophic mandibles

Diameter	Non-atrophic mandibles	Atrophic mandibles
	Mean (SD), mm	Mean (SD), mm
AP right ^a	4.4 (0.92)	3.16 (0.72)
AP left ^a	4.23 (1.13)	3.21 (0.81)
LL right	3.46 (0.73)	3.41 (0.81)
LL left	3.5 (0.61)	3.18 (0.91)

^aStatistically significant at level $P < 0.0001$ (t-test). AP = antero-posterior distance; LL = latero-lateral distance; SD = standard deviation.

Table 4. Comparison of the area of the mandibular foramen between the non-atrophic and atrophic mandibles

Area	Non-atrophic mandibles	Atrophic mandibles
	Mean (SD), mm	Mean (SD), mm
MF right ^a	12.1 (4.27)	8.44 (2.81)
MF left ^a	11.9 (5.03)	8.25 (3.89)

^aStatistically significant at level P < 0.05, t-test, Welch’s correction. MF = mandibular foramen; SD = standard deviation.

Values of the gonial angle

No statistical differences were observed in the gonial angle when the right and left side of the same group were compared, but the angle increased significantly when NAT mandibles were compared to AT mandibles (Table 5).

DISCUSSION

In this study, morphometric and morphological data about the MF were obtained and compared between NAT and AT mandibles. There are two major findings in the present study. First, the knowledge of the position and shape of the MF in both NAT and AT mandibles are important for performing anaesthetic and surgical techniques in these regions. This is the first demonstration of these changes in morphology and area of the MF when NAT and AT mandibles are compared. Second, the mandibular atrophy is not equally distributed and occurs mainly because of dental loss, which leads to loss of masticatory function and atrophy of the chewing muscles. This is important in clinical practice, because these alterations can be partially reversed with oral rehabilitation [19]. Muscle strength directly influences the shape of the mandible [20]. In this article, resorption zones were detected and compared. Therefore, it contributes to the understanding of this process in mandibles.

The knowledge of the position and shape of the MF in both NAT and AT mandibles are part of the morphometric and morphological studies that adopt specific reference points to position this foramen. MF is located on the medial side of the mandibular ramus. In most studies the anterior reference point is the most concave region of the anterior margin of the ramus, and the superior one is the most concave region of the mandibular notch [9,21-23]. But the points adopted as posterior and inferior references vary widely. In this study, the points assumed as references to the

Table 5. Comparison of the gonial angle between non-atrophic and atrophic mandibles

Angle	Non-atrophic mandibles	Atrophic mandibles
	Mean (SD)	Mean (SD)
Right ^a	121.4° (8.17°)	128.9° (7.07°)
Left ^a	121.9° (7.29°)	128.96° (6.61°)

^aStatistically significant at level P < 0.0001 (t-test). SD = standard deviation.

measurements are according to Lima et al. [9].

The present study did not find a significant difference between the right and left sides when the position and shape of the MF were compared in each group NAT and AT mandibles. It is in accordance with the literature [16,21-27].

In addition, it was found that MF in the NAT mandibles was displaced to superior and posterior. These data are in agreement with other studies in the literature [7,9,16,21,26,28].

In the present study, the mandibular atrophy process shows significant resorption in the anterior, posterior, and inferior regions of the mandibular ramus and the MF moved to anterior and inferior. The unique distance that did not change significantly was that between the foramen and the lower region of the mandibular notch, which is in agreement with Prado et al. [21]. However, Matveeva et al. [27] did not find resorption in the anterior part of the ramus.

A few studies in the literature reviewed, evaluated the MF area [9], and we did not find works comparing the shape and area of MF between NAT and AT mandibles. Through the MF runs a neurovascular bundle composed by the artery, vein and inferior alveolar nerve, these structures are separated and involved by connective tissue [4]. In the present study, the MF shape was elliptical, with greater antero-posterior diameter in NAT mandibles. This shape changed to round in AT mandibles, moreover the foramen area decreased significantly. Dental losses can explain these changes. The NAT mandibles evaluated were mostly partially dentate. The dental losses bring to the atrophy process, so there is a decrease in the necessity of blood supply and, therefore, in the diameter of the inferior alveolar artery [29,30]. The same happens in relation to the necessity of innervation thus, a decrease in the number of axons of the inferior alveolar nerve occurs too [31]. In this way, a reduction of the area of the neurovascular bundle and consequently a bone remodelling can be achieved, altering the shape, and decreasing the MF area.

Regarding clinical practice, it is recommended for inferior alveolar nerve block the use of long needles, which penetrate soft tissues up to 2/3 of their length [10]. In this study, the distance between the anterior border of the ramus and the MF was approximately 19.3 mm, so the penetration of the needle should be sufficient to reach the vicinity of the MF in the NAT mandibles and caution is needed in the AT. If the needle exceeds MF and the anaesthetic was deposited near the most posterior margin of the ramus, diffusion to the parotid gland may occur. Therefore, the facial nerve can be reached, and a facial mimic muscles palsy will be observed [32,33]. Moreover, damage in the inferior alveolar neurovascular bundle can happen during the anaesthetic procedure [11]. In AT mandibles the morphological alterations observed in the position of the MF and the mandibular canal must be considered to place implants. The superficialization of the neurovascular bundle due to the resorption of the alveolar bone increases the technical difficulty. Early diagnostic of the injury and correct treatment are essential points to reach full recovery of the sensibility [11,34]

In addition to anaesthetic techniques, the mandible ramus is an area where osteotomies are performed to reposition the mandible due to aesthetic and functional problems [13,35,36]. In our study, the mean distance from the foramen to the posterior margin of the NAT mandibles was 14.3 mm and to the upper and anterior margin of the ramus were 22.7 and 19.3 mm. These distances were important in vertical and sagittal osteotomy techniques respectively and are in accordance with the results described in other studies that considered the same reference points [9,21-24].

In the clinical practice, both for anaesthetic techniques and osteotomies, the surgeon should also consider an alteration in the gonial angle when NAT mandibles

are compared with AT. In this study, the mean of this angle was 121.6° in the NAT mandibles and 128.9° in AT. The increase of the gonial angle seems to be related to the degree of mandibular atrophy [37] what happens predominantly after dental losses [38,39]. Moreover, the increase in gonial angle can occur by the aging [40]. In addition, the angle appears to be lower in men than in women [41-43]. Changes in the shape of the mandible are related to the force exerted by the masseter and temporal muscles. Therefore, the smaller muscular force leads to an increase in the angle [20].

CONCLUSIONS

The results demonstrate a change in the position, shape and area of the mandibular foramen when non-atrophic mandibles were compared to atrophic ones. The mandibular foramen shifted significantly to an anterior and inferior position and changed from an elliptical shape to a round one with a significant decrease in its area. There was a significant increase in the gonial angle. These changes are related to atrophy that occurs in the mandible due mainly to dental losses. Anaesthetic techniques and surgical procedures should consider these changes to prevent vascular and nervous lesions.

ACKNOWLEDGMENT AND DISCLOSURE STATEMENTS

We would like to thank Prof. Dr. Romeu Rodrigues de Souza for his valuable suggestions.

The authors confirm no conflicts of interest related to this original article.

REFERENCES

1. Mina M. Regulation of mandibular growth and morphogenesis. *Crit Rev Oral Biol Med*. 2001;12(4):276-300. [Medline: [11603502](#)] [doi: [10.1177/10454411010120040101](#)]
2. Kjaer I. Correlated appearance of ossification and nerve tissue in human fetal jaws. *J Craniofac Genet Dev Biol*. 1990;10(3):329-36. [Medline: [1702095](#)]
3. Radlanski RJ, Renz H, Klarkowski MC. Prenatal development of the human mandible. 3D reconstructions, morphometry and bone remodelling pattern, sizes 12-117 mm CRL. *Anat Embryol (Berl)*. 2003 Oct;207(3):221-32. [Medline: [13680218](#)] [doi: [10.1007/s00429-003-0343-4](#)]
4. Pogrel MA, Dorfman D, Fallah H. The anatomic structure of the inferior alveolar neurovascular bundle in the third molar region. *J Oral Maxillofac Surg*. 2009 Nov;67(11):2452-4. [Medline: [19837316](#)] [doi: [10.1016/j.joms.2009.06.013](#)]
5. Ikeda K, Ho KC, Nowicki BH, Haughton VM. Multiplanar MR and anatomic study of the mandibular canal. *AJNR Am J Neuroradiol*. 1996 Mar;17(3):579-84. [Medline: [8881258](#)]
6. Juodzbalys G, Wang HL, Sabalys G. Anatomy of mandibular vital structures. Part I: mandibular canal and inferior alveolar neurovascular bundle in relation with dental implantology. *J Oral Maxillofac Res*. 2010 Apr 1;1(1):e2. [Medline: [24421958](#)] [PMC free article: [PMC3886040](#)] [doi: [10.5037/jomr.2010.1102](#)]

7. Ashkenazi M, Taubman L, Gavish A. Age-associated changes of the mandibular foramen position in anteroposterior dimension and of the mandibular angle in dry human mandibles. *Anat Rec (Hoboken)*. 2011 Aug;294(8):1319-25. [Medline: [21714109](#)] [doi: [10.1002/ar.21429](#)]
8. Laishram D, Shastri D. Morphometric Analysis of Mandibular and Mental Foramen. *IOSR-JDMS* 2015 Dec;14(2)Ver. V:82-6. [doi: [10.9790/0853-141258286](#)]
9. Lima FJ, Oliveira Neto OB, Barbosa FT, Sousa-Rodrigues CF. Location, shape and anatomic relations of the mandibular foramen and the mandibular lingula: a contribution to surgical procedures in the ramus of the mandible. *Oral Maxillofac Surg*. 2016 Jun;20(2):177-82. [Medline: [26781719](#)] [doi: [10.1007/s10006-016-0547-4](#)]
10. Thangavelu K, Kannan R, Kumar NS. Inferior alveolar nerve block: Alternative technique. *Anesth Essays Res*. 2012 Jan-Jun;6(1):53-7. [Medline: [25885503](#)] [PMC free article: [4173425](#)] [doi: [10.4103/0259-1162.103375](#)]
11. Juodzbalsys G, Wang HL, Sabalys G. Injury of the Inferior Alveolar Nerve during Implant Placement: a Literature Review. *J Oral Maxillofac Res*. 2011 Apr 1;2(1):e1. [Medline: [24421983](#)] [PMC free article: [3886063](#)] [doi: [10.5037/jomr.2011.2101](#)]
12. Ghai S, Sharma Y, Jain N, Satpathy M, Pillai AK. Use of 3-D printing technologies in craniomaxillofacial surgery: a review. *Oral Maxillofac Surg*. 2018 Sep;22(3):249-259. [Medline: [29797107](#)] [doi: [10.1007/s10006-018-0704-z](#)]
13. McKenna SJ, King EE. Intraoral Vertical Ramus Osteotomy Procedure and Technique. *Atlas Oral Maxillofac Surg Clin North Am*. 2016 Mar;24(1):37-43. [Medline: [26847511](#)] [doi: [10.1016/j.cxom.2015.10.002](#)]
14. Rokutanda S, Yamada S, Yanamoto S, Omori K, Fujimura Y, Morita Y, Rokutanda H, Kohara H, Fujishita A, Nakamura T, Yoshimi T, Yoshida N, Umeda M. Comparison of osseous healing after sagittal split ramus osteotomy and intraoral vertical ramus osteotomy. *Int J Oral Maxillofac Surg*. 2018 Oct;47(10):1316-1321. [Medline: [29843949](#)] [doi: [10.1016/j.ijom.2018.05.013](#)]
15. McLeod NM, Bowe DC. Nerve injury associated with orthognathic surgery. Part 2: inferior alveolar nerve. *Br J Oral Maxillofac Surg*. 2016 May;54(4):366-71. [Medline: [26922403](#)] [doi: [10.1016/j.bjoms.2016.01.027](#)]
16. Cawood JI, Howell RA. A classification of the edentulous jaws. *Int J Oral Maxillofac Surg*. 1988 Aug;17(4):232-6. [Medline: [3139793](#)] [doi: [10.1016/S0901-5027\(88\)80047-X](#)]
17. Luhr HG, Reidick T, Merten HA. Results of treatment of fractures of the atrophic edentulous mandible by compression plating: a retrospective evaluation of 84 consecutive cases. *J Oral Maxillofac Surg*. 1996 Mar;54(3):250-4; discussion 254-5. [Medline: [8600229](#)] [doi: [10.1016/S0278-2391\(96\)90733-8](#)]
18. Shenoy V, Vijayalakshmi S, Saraswathi P. Osteometric Analysis of the Mandibular Foramen in Dry Human Mandibles. *J Clin Diagn Res*. 2012 May;6(4):557-60. doi: JCDR/2012/3873:0002.
19. Boven GC, Raghoobar GM, Vissink A, Meijer HJ. Improving masticatory performance, bite force, nutritional state and patient's satisfaction with implant overdentures: a systematic review of the literature. *J Oral Rehabil*. 2015 Mar;42(3):220-33. [Medline: [25307515](#)] [doi: [10.1111/joor.12241](#)]
20. Sella-Tunis T, Pokhojaev A, Sarig R, O'Higgins P, May H. Human mandibular shape is associated with masticatory muscle force. *Sci Rep*. 2018 Apr 16;8(1):6042. [Medline: [29662127](#)] [PMC free article: [5902585](#)] [doi: [10.1038/s41598-018-24293-3](#)]
21. Prado FB, Groppo FC, Volpato MC, Caria PH. Morphological changes in the position of the mandibular foramen in dentate and edentate Brazilian subjects. *Clin Anat*. 2010 May;23(4):394-8. [Medline: [20235169](#)] [doi: [10.1002/ca.20973](#)]
22. Valente VB, Arita WM, Gonçalves PCG, Campos JADB, Capote TSO. Location of the mandibular foramen according to the amount of dental alveoli. *Int J Morphol*. 2012 Mar;30:77-81. [doi: [10.4067/S0717-95022012000100013](#)]
23. Alves N, Deana NF. Morphometric study of mandibular foramen in macerated skulls to contribute to the development of sagittal split ramus osteotomy (SSRO) technique. *Surg Radiol Anat*. 2014 Nov;36(9):839-45. [Medline: [24671335](#)] [doi: [10.1007/s00276-014-1279-5](#)]
24. Shalini R, RaviVarman C, Manoranjitham R, Veeramuthu M. Morphometric study on mandibular foramen and incidence of accessory mandibular foramen in mandibles of south Indian population and its clinical implications in inferior alveolar nerve block. *Anat Cell Biol*. 2016 Dec;49(4):241-248. [Medline: [28127498](#)] [PMC free article: [5266106](#)] [doi: [10.5115/acb.2016.49.4.241](#)]
25. Ennes JP, Medeiros RM. Localization of mandibular foramen and clinical implications. *Int J Morphol*. 2009 Dec;27(4):1305-1311. [doi: [10.4067/S0717-95022009000400053](#)]
26. Apinhasmit W, Chompoopong S, Jansisyanont P. Alternative Landmarks of the Mandibular Foramen to Prevent Nerve Injury during Ramus Surgery. *J Med Assoc Thai*. 2015 Jun;98(6):574-81. [Medline: [26219162](#)]
27. Matveeva N, Popovska L, Evrosimovska B, Chadikovska E, Nikolovska J. Morphological alterations in the position of the mandibular foramen in dentate and edentate mandibles. *Anat Sci Int*. 2018 Jun;93(3):340-350. [Medline: [29222672](#)] [doi: [10.1007/s12565-017-0423-9](#)]
28. Oguz O, Bozkir MG. Evaluation of location of mandibular and mental foramina in dry, young, adult human male, dentulous mandibles. *West Indian Med J*. 2002 Mar;51(1):14-6. [Medline: [12089867](#)]
29. Wadu SG, Penhall B, Townsend GC. Morphological variability of the human inferior alveolar nerve. *Clin Anat*. 1997;10(2):82-7. [Medline: [9058013](#)] [doi: [10.1002/\(SICI\)1098-2353\(1997\)10:23.0.CO;2-V](#)]
30. Poirot G, Delattre JF, Palot C, Flament JB. The inferior alveolar artery in its bony course. *Surg Radiol Anat*. 1986;8(4):237-44. [Medline: [3107147](#)] [doi: [10.1007/BF02425073](#)]

31. Heasman PA. The myelinated fibre content of human inferior alveolar nerves from dentate and edentulous subjects. *J Dent.* 1984 Dec;12(4):283-6. [Medline: [6596307](#)] [doi: [10.1016/0300-5712\(84\)90083-6](#)]
32. Blanton PL, Jeske AH; ADA Council on Scientific Affairs; ADA Division of Science. Avoiding complications in local anesthesia induction: anatomical considerations. *J Am Dent Assoc.* 2003 Jul;134(7):888-93. [Medline: [12892447](#)] [doi: [10.14219/jada.archive.2003.0288](#)]
33. Ilea A, Cristea A, Tărmure V, Trombitaş VE, Câmpian RS, Albu S. Management of patients with facial paralysis in the dental office: A brief review of the literature and case report. *Quintessence Int.* 2014 Jan;45(1):75-86. [Medline: [24392498](#)] [doi: [10.3290/j.qi.a30770](#)]
34. Shavit I, Juodzbalys G. Inferior alveolar nerve injuries following implant placement - importance of early diagnosis and treatment: a systematic review. *J Oral Maxillofac Res.* 2014 Dec 29;5(4):e2. [Medline: [25635209](#)] [PMC free article: [4306320](#)] [doi: [10.5037/jomr.2014.5402](#)]
35. Agbaje JO, Salem AS, Lambrechts I, Jacobs R, Politis C. Systematic review of the incidence of inferior alveolar nerve injury in bilateral sagittal split osteotomy and the assessment of neurosensory disturbances. *Int J Oral Maxillofac Surg.* 2015 Apr;44(4):447-51. [Medline: [25496848](#)] [doi: [10.1016/j.ijom.2014.11.010](#)]
36. Reyneke JP, Ferretti C. The Bilateral Sagittal Split Mandibular Ramus Osteotomy. *Atlas Oral Maxillofac Surg Clin North Am.* 2016 Mar;24(1):27-36. [Medline: [26847510](#)] [doi: [10.1016/j.cxom.2015.10.005](#)]
37. Xie QF, Ainamo A. Correlation of gonial angle size with cortical thickness, height of the mandibular residual body, and duration of edentulism. *J Prosthet Dent.* 2004 May;91(5):477-82. [Medline: [15153856](#)] [doi: [10.1016/j.prosdent.2004.02.020](#)]
38. Huuonen S, Sipilä K, Haikola B, Tapio M, Söderholm AL, Remes-Lyly T, Oikarinen K, Raustia AM. Influence of edentulousness on gonial angle, ramus and condylar height. *J Oral Rehabil.* 2010 Jan;37(1):34-8. [Medline: [19912483](#)] [doi: [10.1111/j.1365-2842.2009.02022.x](#)]
39. Tozoğlu U, Cakur B. Evaluation of the morphological changes in the mandible for dentate and totally edentate elderly population using cone-beam computed tomography. *Surg Radiol Anat.* 2014 Sep;36(7):643-9. [Medline: [24292424](#)] [doi: [10.1007/s00276-013-1241-y](#)]
40. Leversha J, McKeough G, Myrteza A, Skjellrup-Wakefiled H, Welsh J, Sholapurkar A. Age and gender correlation of gonial angle, ramus height and bigonial width in dentate subjects in a dental school in Far North Queensland. *J Clin Exp Dent.* 2016 Feb 1;8(1):e49-54. [Medline: [26855706](#)] [PMC free article: [4739368](#)] [doi: [10.4317/jced.52683](#)]
41. Casey DM, Emrich LJ. Changes in the mandibular angle in the edentulous state. *J Prosthet Dent.* 1988 Mar;59(3):373-80. [Medline: [3162279](#)] [doi: [10.1016/0022-3913\(88\)90193-X](#)]
42. Aragão JA, Souto ML, Mateus CR, Menezes Ldos S, Reis FP. Edentulousness in relation to remodeling of the gonial angles and incisures in dentate and edentate mandibles: morphometric study using the Image J software. *Surg Radiol Anat.* 2014 Nov;36(9):889-94. [Medline: [25015783](#)] [doi: [10.1007/s00276-014-1335-1](#)]
43. Direk F, Uysal II, Kivrak AS, Unver Dogan N, Fazliogullari Z, Karabulut AK. Reevaluation of Mandibular Morphometry According to Age, Gender, and Side. *J Craniofac Surg.* 2018 Jun;29(4):1054-1059. [Medline: [29461369](#)] [doi: [10.1097/SCS.0000000000004293](#)]

To cite this article:

Mascaro MB, Picoli LC, Reis Matos ST, Sipos Lotaif SC, De Souza MR, Ferreira Calderon M.

Anatomical Variation in the Mandibular Foramen in Non-Atrophic and Atrophic Mandibles

J Oral Maxillofac Res 2020;11(4):e4

URL: <http://www.ejomr.org/JOMR/archives/2020/4/e4/v11n4e4.pdf>

doi: [10.5037/jomr.2020.11404](#)

Copyright © Mascaro MB, Picoli LC, Reis Matos ST, Sipos Lotaif SC, De Souza MR, Ferreira Calderon M. Published in the JOURNAL OF ORAL & MAXILLOFACIAL RESEARCH (<http://www.ejomr.org>), 31 December 2020.

This is an open-access article, first published in the JOURNAL OF ORAL & MAXILLOFACIAL RESEARCH, distributed under the terms of the [Creative Commons Attribution-Noncommercial-No Derivative Works 3.0 Unported License](#), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work and is properly cited. The copyright, license information and link to the original publication on (<http://www.ejomr.org>) must be included.