



Intestinal Parasites and Theta Brainwave Changes in Children

Massomeh Davoudi¹, Raheleh Rafiei Sefiddashti^{1, **}, Ahmadreza Meamar¹, Somayeh Toreyhi² and Ramtin Hadighi^{1, *}

¹Department of Parasitology, Medical School, Iran University of Medical Sciences, Tehran, Iran

²Payame Noor University, Alborz, Iran

*Corresponding author: Department of Parasitology, Medical School, Iran University of Medical Sciences, Tehran, Iran. Email: hadighi.r@iums.ac.ir

**Corresponding author: Department of Parasitology, Medical School, Iran University of Medical Sciences, Tehran, Iran. Email: rafiei.r@iums.ac.ir

Received 2020 March 29; Revised 2020 July 13; Accepted 2020 July 19.

Abstract

Background: Parasitic diseases are one of the health problems in all societies, especially in developing countries. Infection with parasites, especially in children, can be associated with learning disorders, malnutrition, anaemia, and physical disorders. On the other hand, an abnormal increase in the theta brainwave amplitude may also be one of the causes of mental slowness and disability in children's learning, decentralization, or attention deficit hyperactivity disorder (ADHD).

Objectives: In this study, we assessed the possible effects of intestinal parasites on brainwave changes in children in Karaj Health centers.

Methods: A stool examination was done to separate healthy and infected children. Then, using a diagnostic interview and electroencephalography, the theta-wave was recorded in both groups.

Results: From 69 children who participated in this study, 8.7% were infected with one or more intestinal parasites. Around 14.49% of all children had a high theta wave, but the interesting point was that 100% of infected children had a high theta brainwave (P value = 0.001). In contrast, only 6.34% of healthy people had high theta waves.

Conclusions: In this study, there was a significant relationship between parasitic infection and brain wave changes. Parasitic diseases may alter brainwaves with unknown mechanisms and affect the ability and activities of the brain.

Keywords: Parasitic Intestinal Diseases, Electroencephalography, Attention Deficit Hyperactivity Disorder

1. Background

Parasitic infections are one of the most common health problems in developing countries. Various types of intestinal parasites can cause human infections and can produce a wide range of symptoms (1, 2). The most common pathogens of the intestine are *Giardia lamblia*, *Entamoeba histolytic*, *Cryptosporidium* sp., and *Blastocystis* (1, 3). The prevalence of intestinal parasites in primary school children was reported 47% in the city of Varamin of Iran (4). Gastrointestinal complications such as diarrhoea, weight loss, anaemia, abdominal distension, and abdominal pain, as well as some additional disorders such as teeth grinding, night sweats, sleep deprivation, restlessness, anger, and anal itching have been reported as symptoms of these parasites (1).

In Iran, many efforts have been made to control parasitic infections for many years. As a result of this endeavor, the prevalence of different helminths such as *Ascaris lumbricoides* and *Trichuris trichiura* have greatly decreased in recent years (5), but other worms such as *Enterobius vermicularis*, as well as parasitic protozoa like *Giardia lamblia* are still observed among different classes of the society, especially young people because of the social, economic, cultural, geographical, and demographic conditions (2). Some studies have been done on the effects of blood-parasitic infections and their relationship with personality and behavior changes, but few surveys have been done on the association between intestinal parasites and childhood behavioral disorders (6-9). In a study of 194 malnourished children with several parasitic agents, parasites that disturb a person's nutrition seem to have the potential to affect neuropsychological functions (10).

Neurofeedback is a method that assists persons to control brain waves consciously. Neurologic information is given back to the central nervous system (CNS), and it is attempted to teach the person how to correct their brain function. In this method, brain activity is described by using electroencephalography (EEG) (11). The neurofeedback EEG has brain sensors to measure brain waves, and it can change the predetermined range of brain waves, degrees

of oscillation, or phase differences (12). The brain waves are divided from low to high frequency into delta, theta, alpha, beta, and gamma waves, respectively. Theta waves (4 - 8 Hz) play an important role in the internal, memory focus, creativity, intuitive and sensory understanding (13). The most important wave in children is theta wave, the abnormal increase in theta wave amplitude leads to an increase of reaction time and disability in learning and concentration, and hyperactivity (14). As the age increases from childhood to adulthood, the theta wave amplitude decreases gradually, and the alpha wave amplitude increases, but at the end of middle ages and aging, the theta amplitude increases, and the alpha wave decreases again (13). One of the most common neurobehavioral disorders is Attention-deficit/hyperactivity disorder (ADHD) of childhood, which causes an abnormal increase in theta wave and affects the social interactions of children (15).

2. Objectives

Because of the limited studies on the probable efficacy of intestinal parasites on the brain waves of children, the present study examines the changes in the level of theta waves recorded by the neurofeedback system in infected and non-infected children with parasitic agents aged 7 - 11 years in the city of Karaj.

3. Methods

This cross-sectional descriptive study was carried out during the years 2017 - 2018 on 69 girls and boys aged 7 - 11 years in the Karaj City Center who were convinced to participate in this project. All children who were in the above range and the kindergarten officials had given their consent to participate in the project were included in this study. The sample size was estimated by the Cochran formula using the Morgan table in an unlimited society and based on retrospective studies with 95% confidence and a margin of error of 2.5%.

The data gathering tool was a demographic information question form with a query of age, gender, and general health conditions. Participants were assured that personal information was effectively protected against improper usage. The usual procedure for intestinal parasite laboratory diagnosis is a collection of 3 small stool samples that were performed by direct or wet-smear preparation and various concentration methods. Therefore after collection of stool specimens and the completion of the information form, all specimens were examined by macroscopic and microscopic methods, including direct test and formalin ether

concentration. Scotch tape or Graham test was used to isolate and identify the eggs and adult forms of *Enterobius vermicularis* (16).

In the next stage, brain wave assessments were performed with the help of the neurofeedback procedure. The instrument of neurofeedback measurement consists of a diagnostic interview by a psychologist (lasted about an hour) and quantitative electroencephalography. The psychologist's interview included taking history and finding mental health problems. This report represented the current situation, background information about the child, how he/she works in the school, medical history, details on medicine consumption, sleep, exercise, diet, and family (parents, siblings, and psychiatric signs in the whole family). Quantitative electroencephalography (qEEG) was saved as a tool for recording brain activity with electrodes attached to the head. The brain waves were recorded from the 19th Eb Neuro EEG channel 40 using a special hat on the head. The qEEG was done for one-hour with a 128-Hz sampling rate. In an article in 1994, helminth infections were associated with poor memory and IQ abnormalities (17). Lenartowicz et al. (13) in 2014 found that electroencephalography (EEG), with only the use of electrodes at the forehead, plays an important role in assessing the performance of ADHD children due to the cheapness, availability, and non-use of harmful electromagnetic radiation. Any changes of EEG rhythm in the forehead sensors were associated with behavioural problems and as a sign of abnormal brain function. Abnormalities are shown by graphs on the tape in children with attention deficit hyperactivity disorder (13). Brain waves were written in three situations of closed eyes, open eyes, and cognitive tasks (like reading) for at least six minutes. SPSS X16 was used as project software. The relationship between intestinal parasitic infections, demographic factors, and brainwave results was analyzed by the chi-square test, and $P < 0.050$ is considered as statistically significant.

3.1. Ethical Approval

The study was approved by the Institutional Research Ethics Committee School of Medicine of Iran with code number: IR.IUMS.FMD.REC.1397.008.

3.2. Patient Consent

Written informed consent was obtained from all study participants' parents.

4. Results

The statistical population included children aged 7 - 11 years old in Karaj City of Welfare centers in Iran. The age and symptoms of participants were shown in Figure 1.




Figure 1. The left-pie chart shows the percentage of different reported symptoms, and the right pie chart is about the scattering of the project population

The results of epidemiologic studies showed that 6 out of 69 persons (8.7%) had one or more intestinal parasites, from which 4 cases (5.8%) were infected with *E. coli*, 2 cases (2.9%) were infected with *Blastocystis*, and 1 case (1.44%) were infected with *G. lamblia*.

Theta brainwave results display 14.49% (10 persons) of children had an increase in their diagram, but the interesting point was that 100% of infected children had a high brainwave, and a significant relationship was observed between theta wave and intestinal parasites in this survey (P value = 0.001). On the other hand, only 6.34% of healthy people had high theta waves. Theta wave test results according to infected and non-infected individuals are shown in Table 1.

5. Discussion

Parasitic infections are reported globally with high prevalence in poor health and economy societies and can affect the quality of life and learning in infected children (18). In this study, all 69 children were tested with scotch tape and three-time stool examination to find different intestinal helminths like *A. lumbricoides*, *T. trichiura*, and *E. vermicularis* and different protozoa like *G. lamblia*, *E. histolytic*, *Cryptosporidium* sp. and *Blastocystis*. In Iran, the Khash children's study reported the prevalence of 24.1% for *Enterobius* (19), which is probably due to the severe deprivation of health status. Fortunately, the present study showed differing results. In another study in Iran in Rey City, the prevalence of Oxyuridae was 15.49%; again this discrepancy with recent surveys emphasizes the improvement of health levels in the past of 19 years (20). In a relatively new study on mental disabilities in Urmia in Iran (21), the prevalence of *E. vermicularis* was 1.3%, which highlighted, like this study, the promotion of health and welfare in the cities.

Here, the prevalence of intestinal parasitic infections was 8.7%. In similar studies with the same method, in

southern Tehran (2005) (22) and Karaj (2008) of Iran, the prevalence was 10.7% and 7.7%, respectively. Also, the prevalence of parasites was 21.2% in Tehran in 2008 (5), 74.6% in Tonekabon in 1992 (23), 25% in Mazandaran in 2005 (24), and 19.66% in Kirkuk in 2015, by formalin-ether concentration and Ziehl-Neelsen staining (25) that can assume the proportions of contamination were somewhat similar considering the year and cities. Although the spread of parasites was very high in the past based on the overall prevalence of intestinal parasites in wastewaters (26) but today it has declined due to the rising levels of health. Accordingly, the data gathered from this study shows no helminthic infection, which is probably due to the upgrade of urban lifestyles and the improvement of public health and environmental sanitation (27). Another important point in this study was the prevalence of different protozoa. *E. coli* had a rate of 5.8%, similar to the Soleimani et al. study in Ghaemshahr, Iran, in 2014 with an incidence of 2.1% (28) and 9.7% in Tappeh (21), Iran in 2008 between disabled children. In both studies like this, *E. coli* was the most common type of protozoan found in the stool.

In a 2017 review article, cognitive-behavioural studies were done on children infected with soil-transmitted helminths, but the result was very conflicting and contradictory. It was probably because of various methods for the study of children's cognitive activities (29). So in this project, EEG was used as one of the best instruments for evaluating abnormality in brain function both in infected and non-infected children with parasitic agents. It was said that multiple etiologies' conditions can be early markers for ADHD, such as infectious diseases (30). On the other hand, the findings regarding parasite prevalence and cognitive ability have shown a negative relationship between the intensity of infectious diseases and average national IQ worldwide (31). Intestinal parasitic infections are more common in developing countries, which can affect the

Table 1. Theta Wave Test Results According to Infected and Non-Infected Individuals with Different Parasites

Theta Wave	Non-Infected Individuals	Infected Individuals	Total	P Value
High	4	6	10	0.001
Low	59	0	59	
Total	63	6	69	

quality of life of people infected with these infections and in children, the effects of which are reported to be more severe. On the other hand, with increasing age from childhood to adulthood, the range of theta wave gradually decreases, and the amplitude of the alpha wave increases. However, in late middle age and old age, the amplitude of the theta wave, and the decrease of the alpha wave amplitude are observed again. It is seen in people with a variety of neurological disorders, as well as in children with ADHD and learning disabilities, an increase in theta wave amplitude and a decrease in the alpha amplitude similar to what happens when a child is infected with parasites.

5.1. Conclusions

The data of this survey express that intestinal parasitic infection rates have relatively decreased, probably be due to the rise of hygiene standards. In addition, 100% of children infected with parasites had high theta wavelengths that could lead to long reaction time, learning disability, and lack of concentration observed in these children. In other words, it might be said that abnormalities of the theta wavelength are associated with parasitic diseases. It seems, parasites probably could be capable of altering the brain. It is suggested that this study be conducted in a larger society and on a larger number of people with parasitic agents before and after treatment.

Footnotes

Authors' Contribution: Study concept and design: Ramtin Hadighi and Somayeh Toreyhi. Analysis and interpretation of data: Massomeh Davoudi and Somayeh Toreyhi. Drafting of the manuscript: Massomeh Davoudi. Co-corresponding and critical revision of the manuscript for important intellectual content: Raheleh Rafiei Sefidashti.

Conflict of Interests: The authors declare that they have no conflict of interest.

Ethical Approval: The study was approved by the Institutional Research Ethics Committee School of Medicine of Iran with code number: IR.IUMS.FMD.REC.1397.008.

Funding/Support: This research was supported by grant no.: 31707 from Iran University of Medical Sciences.

Informed Consent: Written informed consent was obtained from all study participants' parents.

References

1. Haque R. Human intestinal parasites. *J Health Popul Nutr.* 2007;**25**(4):387-91. [PubMed: [18402180](#)]. [PubMed Central: [PMC2754014](#)].
2. Sayyari AA, Imanzadeh F, Bagheri Yazdi SA, Karami H, Yaghoobi M. Prevalence of intestinal parasitic infections in the Islamic Republic of Iran. *EMHJ-East Mediterr Health J.* 2005;**11**(3):377-83.
3. Horton J. Human gastrointestinal helminth infections: are they now neglected diseases? *Trends Parasitol.* 2003;**19**(11):527-31. doi: [10.1016/j.pt.2003.09.007](#). [PubMed: [14580965](#)].
4. Aminzadeh Z, Tarami M, Gachkar L. Prevalence of intestinal parasites and related factors in primary school children in Varamin. *Iran J Pediatr Soc.* 2007;**1**:55-8.
5. Akhlaghi L, Shamseddin J, Meamar AR, Razmjou E, Oormazdi H. Frequency of intestinal parasites in Tehran. *Iranian Journal of Parasitology.* 2009;**44**-7.
6. Arpino C, Gattinara GC, Piergili D, Curatolo P. Toxocara infection and epilepsy in children: a case-control study. *Epilepsia.* 1990;**31**(1):33-6. doi: [10.1111/j.1528-1157.1990.tb05356.x](#). [PubMed: [2303010](#)].
7. Brooks JM, Carrillo GL, Su J, Lindsay DS, Fox MA, Blader IJ. Toxoplasma gondii Infections Alter GABAergic Synapses and Signaling in the Central Nervous System. *mBio.* 2015;**6**(6):e01428-15. doi: [10.1128/mBio.01428-15](#). [PubMed: [26507232](#)]. [PubMed Central: [PMC4626855](#)].
8. Stommel EW, Seguin R, Thadani VM, Schwartzman JD, Gilbert K, Ryan KA, et al. Cryptogenic epilepsy: an infectious etiology? *Epilepsia.* 2001;**42**(3):436-8. doi: [10.1046/j.1528-1157.2001.25500.x](#). [PubMed: [11442166](#)].
9. Oberhelman RA, Guerrero ES, Fernandez ML, Silio M, Mercado D, Comiskey N, et al. Correlations between intestinal parasitosis, physical growth, and psychomotor development among infants and children from rural Nicaragua. *Am J Trop Med Hyg.* 1998;**58**(4):470-5. doi: [10.4269/ajtmh.1998.58.470](#). [PubMed: [9574794](#)].
10. Levav M, Mirsky AF, Schantz PM, Castro S, Cruz ME. Parasitic infection in malnourished school children: effects on behaviour and EEG. *Parasitology.* 1995;**110** (Pt 1):103-11. doi: [10.1017/s0031182000081105](#). [PubMed: [7845707](#)].
11. Nachvak SM, Haghighat HR, Rezaei M. Prevalence and monitoring of epilepsy in mentally retarded students, 2002. *Behav Sci Q.* 2004;**8**:34-42.
12. Arns M, de Ridder S, Strehl U, Breteler M, Coenen A. Efficacy of neuro-feedback treatment in ADHD: the effects on inattention, impulsivity and hyperactivity: a meta-analysis. *Clin EEG Neurosci.* 2009;**40**(3):180-9. doi: [10.1177/155005940904000311](#). [PubMed: [19715181](#)].
13. Lenartowicz A, Loo SK. Use of EEG to diagnose ADHD. *Curr Psychiatry Rep.* 2014;**16**(11):498. doi: [10.1007/s11920-014-0498-0](#). [PubMed: [25234074](#)]. [PubMed Central: [PMC4633088](#)].

14. Persinger MA, St-Pierre LS, Saroka KS. LORETA predicts electromagnetic sensitivity and "hearing voices" in a predictable, increasingly prevalent subpopulation: possible QEEG-based differential diagnosis. *Neuropsychiatr Electrophysiol*. 2015;**1**(1). doi: [10.1186/s40810-015-0007-7](https://doi.org/10.1186/s40810-015-0007-7).
15. Subcommittee on Attention-Deficit/Hyperactivity D, Steering Committee on Quality I, Wolraich M, Brown L, Brown RT; Management, et al. ADHD: clinical practice guideline for the diagnosis, evaluation, and treatment of attention-deficit/hyperactivity disorder in children and adolescents. *Pediatrics*. 2011;**128**(5):1007-22. doi: [10.1542/peds.2011-2654](https://doi.org/10.1542/peds.2011-2654). [PubMed: [22003063](https://pubmed.ncbi.nlm.nih.gov/22003063/)]. [PubMed Central: [PMC4500647](https://pubmed.ncbi.nlm.nih.gov/PMC4500647/)].
16. Garcia LS. Laboratory identification of the microsporidia. *J Clin Microbiol*. 2002;**40**(6):1892-901. doi: [10.1128/jcm.40.6.1892-1901.2002](https://doi.org/10.1128/jcm.40.6.1892-1901.2002). [PubMed: [12037040](https://pubmed.ncbi.nlm.nih.gov/12037040/)]. [PubMed Central: [PMC130667](https://pubmed.ncbi.nlm.nih.gov/PMC130667/)].
17. Nokes C, Bundy DA. Does helminth infection affect mental processing and educational achievement? *Parasitol Today*. 1994;**10**(1):14-8. doi: [10.1016/0169-4758\(94\)90348-4](https://doi.org/10.1016/0169-4758(94)90348-4). [PubMed: [15275558](https://pubmed.ncbi.nlm.nih.gov/15275558/)].
18. World Health Organization. *Deworming for health and development: report of the Third Global Meeting of the Partners for Parasite Control*. World Health Organization; 2005.
19. Ebrahimzadeh A, Saryazdipoor KH, Gharaei A, Mohammadi S, Jamshidi A. Prevalence of Enterobius vermicularis infection among preschool children of Khash city kindergartens, Iran in 2012. *J North Khorasan Univ Med Sci*. 2014;**6**(3):477-81.
20. Rafiei M, Torkaman M, Sharbatdar AMR. Asymptomatic Giardiasis in school children in Rey city. *Tehran Univ Med J*. 2000.
21. Tappeh Kh H, Mohammadzadeh H, Rahim RN, Barazesh A, Khashaveh S, Taherkhani H. Prevalence of Intestinal Parasitic Infections among Mentally Disabled Children and Adults of Urmia, Iran. *Iran J Parasitol*. 2010;**5**(2):60-4. [PubMed: [22347245](https://pubmed.ncbi.nlm.nih.gov/22347245/)]. [PubMed Central: [PMC3279829](https://pubmed.ncbi.nlm.nih.gov/PMC3279829/)].
22. Arani AS, Alaghebandan R, Akhlaghi L, Shahi M, Lari AR. Prevalence of intestinal parasites in a population in south of Tehran, Iran. *Rev Inst Med Trop Sao Paulo*. 2008;**50**(3):145-9. doi: [10.1590/s0036-46652008000300003](https://doi.org/10.1590/s0036-46652008000300003). [PubMed: [18604414](https://pubmed.ncbi.nlm.nih.gov/18604414/)].
23. Rezaian M, Hooshyar H. The prevalence of intestinal parasitic infection in rural areas of Tonekabon, Iran. *Iran J Public Health*. 1996;47-58.
24. Kia EB, Hosseini M, Nilforoushan MR, MEMAR AR, Rezaeian M. Study of intestinal protozoan parasites in rural inhabitants of Mazandaran province, Northern Iran. *Iran J Parasitol*. 2008.
25. Salman YJ, Aziz Al-Tae AR, Abid AM. Prevalence of Giardia lamblia among Iraqi Displaced Peoples in Kirkuk Province. *Int J Curr Microbiol Appl Sci*. 2016;**5**(1):753-60. doi: [10.20546/ijcmas.2016.501.076](https://doi.org/10.20546/ijcmas.2016.501.076).
26. Hatam-Nahavandi K, Mohebbali M, Mahvi AH, Keshavarz H, Khanaliha K, Tarighi F, et al. Evaluation of Cryptosporidium oocyst and Giardia cyst removal efficiency from urban and slaughterhouse wastewater treatment plants and assessment of cyst viability in wastewater effluent samples from Tehran, Iran. *J Water Reuse Desalination*. 2015;**5**(3):372-90.
27. van Lieshout L, Verweij JJ. Newer diagnostic approaches to intestinal protozoa. *Curr Opin Infect Dis*. 2010;**23**(5):488-93. doi: [10.1097/QCO.0b013e32833de0eb](https://doi.org/10.1097/QCO.0b013e32833de0eb). [PubMed: [20683263](https://pubmed.ncbi.nlm.nih.gov/20683263/)].
28. Soleymani E, Davoodi L, Azami D. The prevalence of intestinal parasitic infections among the mentally retarded patients in Lamook Rehabilitation Center of Qaemshahr, Mazandaran Province, 2015. *Tabari Biomed Stud Res J*. 2016;**2**(1):1-5.
29. Owada K, Nielsen M, Lau CL, Clements ACA, Yakob L, Soares Magalhaes RJ. Measuring the Effect of Soil-Transmitted Helminth Infections on Cognitive Function in Children: Systematic Review and Critical Appraisal of Evidence. *Adv Parasitol*. 2017;**98**:1-37. doi: [10.1016/bs.apar.2017.05.002](https://doi.org/10.1016/bs.apar.2017.05.002). [PubMed: [28942767](https://pubmed.ncbi.nlm.nih.gov/28942767/)].
30. Silva D, Colvin L, Hagemann E, Stanley F, Bower C. Children diagnosed with attention deficit disorder and their hospitalisations: population data linkage study. *Eur Child Adolesc Psychiatry*. 2014;**23**(11):1043-50. doi: [10.1007/s00787-014-0545-8](https://doi.org/10.1007/s00787-014-0545-8). [PubMed: [24770488](https://pubmed.ncbi.nlm.nih.gov/24770488/)].
31. Eppig C, Fincher CL, Thornhill R. Parasite prevalence and the worldwide distribution of cognitive ability. *Proc Biol Sci*. 2010;**277**(1701):3801-8. doi: [10.1098/rspb.2010.0973](https://doi.org/10.1098/rspb.2010.0973). [PubMed: [20591860](https://pubmed.ncbi.nlm.nih.gov/20591860/)]. [PubMed Central: [PMC2992705](https://pubmed.ncbi.nlm.nih.gov/PMC2992705/)].