

Epigenetic regulation of the pathological process in endometriosis

Kuei-Yang Hsiao¹  | Meng-Hsing Wu² | Shaw-Jenq Tsai¹ 

¹Department of Physiology, College of Medicine, National Cheng Kung University, Tainan, Taiwan

²Department of Obstetrics and Gynecology, College of Medicine, National Cheng Kung University, Tainan, Taiwan

Correspondence

Shaw-Jenq Tsai, Department of Physiology, College of Medicine, National Cheng Kung University, Tainan, Taiwan.
Email: seantsai@mail.ncku.edu.tw

Funding information

Ministry of Science and Technology, Taiwan, Grant/Award Number: MOST 104-2320-B-006 -036 -MY3

Abstract

Background: Endometriosis is one of the most common gynecological diseases that greatly compromises the quality of life in affected individuals. A growing body of evidence shows that the remodeling of retrograde endometrial tissues to the ectopic endometriotic lesions involves multiple epigenetic alterations, such as DNA methylation, histone modification, and microRNA expression.

Methods: This article retrospectively reviewed the studies that were related to the epigenetic regulatory factors that contribute to the development and maintenance of endometriosis. A literature search was performed in order to collect scientific articles that were written in English by using the key words of “endometriosis,” “epigenetics,” “DNA methylation,” “histone modification,” and “microRNA.”

Results: Epigenetic modifications, including DNA methylation, histone modification, and microRNA expression, are involved in the pathogenesis of endometriosis. These epigenetic players are regulated or tuned by microenvironmental cues, such as locally produced estradiol, proinflammatory cytokines, and hypoxic stress, and reciprocally regulate the process or response to those stimuli.

Conclusion: Understanding the molecular mechanisms that underlie these epigenetic regulatory processes would shed light on the etiology and/or progression of endometriosis and facilitate the development of novel therapeutic strategies.

KEYWORDS

DNA methylation, endometriosis, epigenetics, histone modification, microRNA

1 | INTRODUCTION

Endometriosis, defined as the presence of endometrial glandular and stromal tissues, is one of the most common gynecological diseases, with a 10%-15% prevalence rate in women of reproductive age. The combination of retrograded menses and the immunosuppression hypothesis is the most accepted theory of the pathogenesis of endometriosis. Although the ectopic lesions were established from eutopic tissues, mounting evidence indicated that the characteristics of each are very distinct, suggesting that epigenetic regulation could be involved in the alteration of these phenotypes.

Epigenetic mechanisms have been recognized as important players in the development of a broad range of human diseases, including cancerous, neurologic, endocrine, and immune diseases. The alteration of chromatin conformation constitutes the basis of epigenetic regulation because the pattern of gene expression changes without changing the genomic sequence. Chromatin conformation can be altered by DNA methylation and post-translational modifications of histones. The change of chromatin conformation alters the accessibility of DNA to its modulator, which can be either a transcription activator or repressor, and thus the subsequent outcome of an open chromatin would be highly context-dependent. In this mini-review, the roles or implications

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2017 The Authors. Reproductive Medicine and Biology published by John Wiley & Sons Australia, Ltd on behalf of Japan Society for Reproductive Medicine.

of different epigenetic elements, including DNA methylation, histone modification, and microRNA (miRNA) expression, will be discussed.

2 | ALTERATION OF DNA METHYLATION IN ENDOMETRIOSIS

It has been known that the alteration of DNA methylation plays an important role and has great impacts on the chromatin remodeling and transcription networks. The major form of DNA methylation in mammalian cells is 5-methyl cytosine, which is catalyzed by a group of DNA methyltransferases (DNMTs), consisting of DNMT1, DNMT3a, and DNMT3b. DNA methyltransferase1 takes hemi-methylated DNA as a substrate, which is responsible for cell cycle-coupled DNA methylation that transmits the epigenetic marks from passage to passage. In contrast, DNMT3a and 3b are de novo methyltransferases that use unmethylated DNA as a substrate. Although both hypomethylated and hypermethylated DNA for specific genes have been reported in endometriotic epithelial and stromal cells (see below for details), the authors' recent study revealed that the genome of endometriotic stromal cells is globally hypomethylated due to the downregulation of DNMT1.¹

It has been comprehensively discussed that DNA methylation plays an important role during the pathogenesis of endometriosis.² Thus, in this review, the focus is on how it might be regulated and interact with other epigenetic elements. Hypoxia and inflammation, two critical driving forces for the development of endometriosis,^{3,4} modulate the expression of DNMTs distinctly and can act together to cause aberrant DNA methylation patterns^{1,5,6} (Figure 1). Two recent reports




FIGURE 1 Hypoxia and inflammation regulate DNA methylation through microRNAs (miRs). These two major microenvironmental cues promote epigenetic remodeling. Hypoxia causes global hypomethylation through miR-148a-mediated DNA methyltransferase (DNMT)1 downregulation (left). Inflammatory stimulation induces DNMT3a expression and loci-specific hypermethylation (right). Hypoxia-induced miR-20a and inflammatory cytokine-induced miR-302a suppress dual specificity phosphatase (DUSP)2 and chicken ovalbumin upstream promoter (COUP)-transcription factor (TF)II, respectively, to potentiate an inflammatory response and to contribute to the modulation of DNA methylation. COX, cyclo-oxygenase; E₂, estradiol; IL, interleukin; PGE₂, prostaglandin E₂; TGF, transforming growth factor; TNF, tumor necrosis factor

found that global methylation decreases in ectopic stromal cells,^{1,6} which is mainly caused by hypoxia-mediated DNMT1 downregulation (Figure 1, left). In contrast to the suppressive effect of hypoxia on DNMT1, a blockage of the prostaglandin E₂ (PGE₂) pathway has no effect on the level of DNMT1 but suppresses DNMT3a expression⁵ (Figure 1, right), implying that the level of DNMT3a might be maintained or stimulated by an inflammation pathway.

The development and maintenance of endometriosis highly depend on the estrogen pathway. The expression of steroidogenic acute regulatory protein (StAR) and aromatase (CYP19), two proteins that control the key steps of 17β-estradiol biosynthesis, was shown to be elevated in the ectopic tissues and isolated primary stromal cells.^{7,8} During the last two decades, studies have shown that hypoxia and inflammation play a central role in the regulation of this steroidogenic pathway during the development of endometriosis (Figure 2). It has been shown that StAR and CYP19 harbor less methylated caffeine phenylethanoid glycoside islands in their promoter and/or intronic regions,⁹⁻¹¹ contributing to their aberrant expression in ectopic lesions. In addition, the promoter and/or intronic regions of several aberrantly expressed nuclear receptors that mediate the effect of steroid hormones or modulate the steroidogenic activity, such as estrogen receptor (ER)β¹² and steroidogenic factor (SF)-1,^{13,14} also were hypomethylated, highlighting the central role of epigenetic dysregulation on the pathway of steroid hormones. In contrast, the inactivation of 17β-estradiol also is regulated by DNA methylation. The gene body of 17β-hydroxysteroid dehydrogenase type II, the enzyme that converts 17β-estradiol to the less potent form, estrone, is hypermethylated and is inactivated in ectopic stromal cells (Figure 2).¹⁵ Not only so, even the promoter of progesterone receptor isoform B, a functional nuclear receptor that induces 17β-hydroxysteroid dehydrogenase type II expression, is also hypermethylated in endometriotic cells,¹⁶ a phenomenon that is probably mediated by tumor necrosis factor (TNF) α.¹⁷ As a result, the conversion of the potent 17β-estradiol to the less potent estrone is suppressed. These data indicate that DNA methylation is coordinately regulated to facilitate the production or to enhance the activity of 17β-estradiol in endometriosis.

In parallel to 17β-estradiol biosynthesis, PGE₂ production that is mediated by the elevated expression of cyclo-oxygenase (COX)-2 also plays critical roles in the development of endometriosis.¹⁸ It has been reported that the promoter of COX-2 is hypomethylated⁹ and contributes to aberrant COX-2 induction in ectopic stromal cells, which in turn enhances the positive feedback to PGE₂-mediated 17β-estradiol production¹⁹ and DNMT3a elevation.

3 | ALTERATION OF THE HISTONE CODE IN ENDOMETRIOSIS

3.1 | Histone acetylation

Histones, the key components of the nucleosome, pack the lengthy genomic DNA molecules into compact forms in the nuclei. The core histones share a conserved tripartic structure, consisting of the amino-terminal tail, a globular domain, and a carboxyl-terminal tail.




FIGURE 2 DNA methylation contributes to inflammation and the acquisition of steroidogenic capacity and enhances estradiol (E_2) signaling. Global DNA hypomethylation contributes to the upregulation of the genes that are involved in prostaglandin E_2 (PGE_2) biosynthesis and E_2 biosynthesis and signaling, while loci-specific hypermethylation suppresses the genes that are related to E_2 inactivation. COX, cyclo-oxygenase; CYP, cytochrome P450; ER, estrogen receptor; HSD, hydroxysteroid dehydrogenase; PRB, progesterone receptor isoform B; SF, steroidogenic factor; StAR, steroidogenic acute regulatory protein

The majority of post-translational modification takes place on the amino-terminal tail of histones extruding from the nucleosome, which regulates various molecular processes, such as chromatin remodeling, transcription, splicing, and DNA damage.

Histone acetylation, one of the earliest discovered modifications, promotes transcriptional activation through the disruption of electric charges between the DNA and histone tail and/or acetyl-lysine reading proteins. Two categories of proteins, histone acetyltransferases and histone deacetylases (HDACs), counteract each other to modulate the levels of histone acetylation. It has been reported that the levels of HDAC1 and/or HDAC2, two of the most abundant HDACs in human cells, were deregulated in endometriotic stromal cells. While one study reported that both HDAC1 and HDAC2 were upregulated in endometriotic stromal cells,²⁰ two other studies reported the upregulation of HDAC1 and HDAC2.^{21,22} Nevertheless, the expression of HDAC1 and HDAC2 were induced by the steroid hormones, 17 β -estradiol and progesterone,²⁰ a notion that is consistent with the central role of steroid hormones in the development of endometriosis (Figure 3). Accompanying the aberrant expression of HDACs, the global levels of histone H3 and H4 acetylation decreased in the endometriotic stromal cells.^{21,23} Of special note, while both of these studies found decreased levels of histone H3 acetylation, one reported no difference in histone H4 acetylation. This discrepancy could partly result from the distinct antibodies that are used for detecting acetylation sites (not specified in all studies). In addition, it has been known that the application of antibodies to explore the combinational modifications of histone is constrained by the nature of multiple adjacent modifications,^{24,25} which can vary the results from laboratory to laboratory. Intriguingly, although inactive genes (e.g. *ER α* , *CCAAT-enhancer-binding protein* [*C/EBP*] α , *CDH1*, *p21*, and *homeobox A10*) in ectopic endometriotic lesions have lower levels of histone H3 and/or H4 acetylation in their promoters, aberrantly expressed genes (eg, *SF-1*) have a high level of histone acetylation in their promoters,^{23,26} suggesting that the level or distribution of acetylation is not solely regulated by HDACs, but also by other acetyltransferases in a gene-specific manner. For example, the histone acetyltransferases, such as steroid receptor coactivator-1, p300, and cyclic adenosine monophosphate response element binding




FIGURE 3 Histone acetylation contributes to an acquired steroidogenic capacity. Estradiol (E_2) and progesterone (P_4) together induce the expression of histone deacetylase (HDAC)1 and HDAC2, which further cause the downregulation of estrogen receptor (ER) α . Histone acetyltransferases, such as steroid receptor coactivator (SRC)-1, p300, and cyclic adenosine monophosphate response element binding protein (CBP), are required for the actively expressed genes, steroidogenic factor (SF)-1 and steroidogenic acute regulatory protein (StAR) in endometriotic cells. PGE_2 , prostaglandin E_2

protein, have been reported to regulate the function of the estrogen receptor, PGE_2 -induced 17 β -estradiol synthesis, and the development of endometriotic lesions (Figure 3).^{19,27,28}

3.2 | Histone deacetylase inhibitor for endometriosis treatment

The critical role of histone acetylation in the transactivation of key genes in endometriosis makes it an attractive therapeutic target.

Treatment with HDAC inhibitors in vitro (immortalized human endometrial stromal and epithelial cells) and in vivo (rat model of endometriosis) caused cell cycle arrest, apoptosis, and thus reduced the lesion size in vivo.^{29,30} Other works that focused on the reactivation of the genes that suppress the development of endometriosis, such as *C/EBPα* and *death receptor 6*, also are reported.^{26,31} Of particular note, although modulating histone acetylation might ameliorate endometriosis, one should bear in mind that other molecular events also are controlled by the means of acetylation. For example, the application of HDAC inhibitors to cause histone hyperacetylation inhibits the mitosis and DNA damage responses.³²⁻³⁵ Thus, how to maximize the therapeutic impact and minimize cytotoxicity should be thoroughly investigated before HDAC inhibitors can be used to treat endometriosis.

4 | DYSREGULATION OF MICRORNA EXPRESSION IN ENDOMETRIOSIS

MicroRNAs belong to a group of single-stranded, non-coding RNAs with an average size of 22 nucleotides. They play important regulatory roles in gene expression through pairing with messengerRNA (mRNA) to modulate RNA splicing, degradation, and translation.³⁶ The genome-wide analysis of the miRNA expression profile demonstrated that dysregulated miRNAs play critical roles during the development of endometriosis through modulating the cell cycle progression, apoptosis, steroidogenic pathway, hormone signaling, inflammation, and response to hypoxia.^{37,38} MicroRNAs that target the mediators of inflammation (COX-2, interleukin [IL]-6, and IL-6 receptors), inducer of apoptosis (B-cell lymphoma-2), cycle regulator (cyclin D1), and angiogenic factors (vascular endothelial growth factor, IL-8) are typically downregulated in the endometrium and/or endometriotic tissues of women with endometriosis, supporting the multifaceted role of miRNAs during the pathogenesis of endometriosis.

4.1 | MicroRNAs regulate the steroidogenic pathway


It has been reported that miRNAs modulate the signaling pathway of steroid hormones, which plays a central role in the pathogenesis

of endometriosis. The expression levels of miR-23a and miR-23b are aberrantly suppressed in the endometrium of women with endometriosis, compared to those of women without endometriosis. Furthermore, the levels of miR-23a and b are much lower in the ectopic endometriotic lesions.³⁹ Typically, a high level of miR-23a and b in the normal endometrium inhibits the expression of SF-1 and thus keeps the normal endometrium in low steroidogenic activity. However, in the endometrium of women with endometriosis, the suppressed level of miR-23a and b allows for the elevated expression of SF-1 and further promotes the expression of StAR and CYP19, suggesting that miR-23a and b play an important role in the acquisition of the steroidogenic capacity of the endometriotic tissues (Figure 4). Although a hypomethylated promoter was reported for SF-1 in endometriosis, a hypermethylated promoter of miR-23a and b was one of the mechanisms responsible for its downregulation in gynecological cancers.⁴⁰

4.2 | MicroRNA-mediated hyperactivated inflammatory responses

In addition to the steroidogenic pathway, miRNAs also modulate inflammation. The effect of PGE₂ was prolonged by hypoxia-induced miR-20a that targets dual specificity phosphatase (DUSP)2, which is a repressor of extracellular signal-regulated kinase (ERK)-1 and -2 signaling downstream of PGE₂. The miR-20a-mediated hypoxia-inhibited DUSP2 prolongs the phosphorylation of ERK-1 and -2, which further augments the PGE₂ signaling and thus potentiates PGE₂-induced gene expression⁴¹ (Figures 1 and 4). In contrast, the biosynthesis of PGE₂ also is stimulated through a miRNA-mediated positive feedback loop.⁴² Chicken ovalbumin upstream promoter (COUP)-transcription factor (TF)II, which binds the COX-2 promoter region and elevates the expression of COX-2 in the eutopic endometrium, is suppressed by inflammatory stimuli from the peritoneal fluid. Treatment with IL-1β, TNFα, or transforming growth factor β suppresses the expression of COUP-TFII through miR-302a binding to its 3'-untranslated region (UTR). In a similar manner to DUSP2, the downregulated repressive transcription factor, COUP-TFII, augments the inflammatory cytokine-induced COX-2 expression, which forms a positive feedback loop for the inflammatory stimulation (Figures 1 and 4).

FIGURE 4 MicroRNAs (miRs) that are involved in the development of endometriosis. DNA hypermethylation-suppressed miRNAs and hypoxia- and inflammation-induced miRNAs work coordinately to regulate multiple cellular processes and to promote the development of endometriosis. Bcl, B-cell lymphoma; COUP, chicken ovalbumin upstream promoter; DNMT, DNA methyltransferase; DUSP, dual specificity phosphatase; SF, steroidogenic factor; TF, transcription factor; VEGFA, vascular endothelial growth factor A



4.3 | MicroRNA-mediated hypoxic response

As mentioned previously, the global DNA hypomethylation in the ectopic tissues is triggered by microenvironmental hypoxic stress.¹ MicroRNA-148a, the key player mediating this global passive demethylation, is aberrantly elevated in the ectopic stromal cells. The administration of exogenous oligo-mimicking miR-148a suppresses the expression of DNMT1, while treatment with the miRNA inhibitor to miR-148a rescues the hypoxia-inhibited DNMT1 expression. Under hypoxia, miR-148a and argonaute 2, an essential component of the RNA-induced silencing complex, coordinate with adenylate-uridylate-rich element RNA-binding protein 1 to bind to 3'-UTR of DNMT1 to cause the degradation of its mRNA. The decreased DNMT1 level in the endometriotic stromal cells causes the passive demethylation of numerous genes through multiple cell cycle progression. The downregulated DNMT1 and global hypomethylation might further trigger the DNMT3a- and 3b-mediated locus-specific hypermethylation. Two recent studies reported that the locus-specific hypermethylation-mediated miR-196b and miR-503 suppression derepresses the genes that are related to the inhibition of cell proliferation, induction of apoptosis, and angiogenesis, thus promoting the pathogenesis of endometriosis.^{38,43} These observations suggest the regulatory complexity and feasibility between epigenetics and small non-coding RNAs (Figure 4).

5 | CONCLUSION

It has been demonstrated that estrogen signaling, hypoxia, and inflammation are three interlinked driving forces in the development of endometriosis, while epigenetic components play a central role in coordinating these three factors. Through the understanding of epigenetic modulations in retrograde endometrial tissues, a more comprehensive picture about how ectopic lesions are transformed and remodeled to a distinct steroidogenic tissue could be shown. By dissecting the interactions among the microenvironmental factors (ie, hypoxia and inflammation) and epigenetic regulation (ie, DNA methylation, histone modification, and miRNA expression), it might be possible to develop novel remedies that target the epigenetic effectors for the better treatment of endometriosis.

DISCLOSURES

Conflict of interest: The authors declare no conflict of interest.
Human and Animal Rights: This article does not contain any study with human or animal participants that has been performed by any of the authors.

ORCID

Kuei-Yang Hsiao  <http://orcid.org/0000-0002-1648-974X>

Shaw-Jenq Tsai  <http://orcid.org/0000-0002-3569-5813>

REFERENCES

1. Hsiao KY, Wu MH, Chang N, et al. Coordination of AUF1 and miR-148a destabilizes DNA methyltransferase 1 mRNA under hypoxia in endometriosis. *Mol Hum Reprod*. 2015;21:894-904.
2. Nasu K, Kawano Y, Tsukamoto Y, et al. Aberrant DNA methylation status of endometriosis: epigenetics as the pathogenesis, biomarker and therapeutic target. *J Obstet Gynaecol Res*. 2011;37:683-695.
3. Hsiao KY, Lin SC, Wu MH, Tsai SJ. Pathological functions of hypoxia in endometriosis. *Front Biosci (Elite Ed)*. 2015;7:309-321.
4. Wu MH, Hsiao KY, Tsai SJ. Endometriosis and possible inflammation markers. *GMIT*. 2015;4:61-67.
5. Arosh JA, Lee J, Starzinski-Powitz A, Banu SK. Selective inhibition of prostaglandin E2 receptors EP2 and EP4 modulates DNA methylation and histone modification machinery proteins in human endometriotic cells. *Mol Cell Endocrinol*. 2015;409:51-58.
6. Senthong A, Kitkumthorn N, Rattanatanyong P, Khemapech N, Triratanachart S, Mutirangura A. Differences in LINE-1 methylation between endometriotic ovarian cyst and endometriosis-associated ovarian cancer. *Int J Gynecol Cancer*. 2014;24:36-42.
7. Tsai SJ, Wu MH, Lin CC, Sun HS, Chen HM. Regulation of steroidogenic acute regulatory protein expression and progesterone production in endometriotic stromal cells. *J Clin Endocrinol Metab*. 2001;86:5765-5773.
8. Noble LS, Simpson ER, Johns A, Bulun SE. Aromatase expression in endometriosis. *J Clin Endocrinol Metab*. 1996;81:174-179.
9. Wang D, Chen Q, Zhang C, Ren F, Li T. DNA hypomethylation of the COX-2 gene promoter is associated with up-regulation of its mRNA expression in eutopic endometrium of endometriosis. *Eur J Med Res*. 2012;17:12.
10. Izawa M, Harada T, Taniguchi F, Ohama Y, Takenaka Y, Terakawa N. An epigenetic disorder may cause aberrant expression of aromatase gene in endometriotic stromal cells. *Fertil Steril*. 2008;89:1390-1396.
11. Izawa M, Taniguchi F, Uegaki T, et al. Demethylation of a nonpromoter cytosine-phosphate-guanine island in the aromatase gene may cause the aberrant up-regulation in endometriotic tissues. *Fertil Steril*. 2011;95:33-39.
12. Meyer JL, Zimbardi D, Podgaec S, Amorim RL, Abrao MS, Rainho CA. DNA methylation patterns of steroid receptor genes ESR1, ESR2 and PGR in deep endometriosis compromising the rectum. *Int J Mol Med*. 2014;33:897-904.
13. Xue Q, Lin Z, Yin P, et al. Transcriptional activation of steroidogenic factor-1 by hypomethylation of the 5' CpG island in endometriosis. *J Clin Endocrinol Metab*. 2007;92:3261-3267.
14. Xue Q, Zhou YF, Zhu SN, Bulun SE. Hypermethylation of the CpG island spanning from exon II to intron III is associated with steroidogenic factor 1 expression in stromal cells of endometriosis. *Reprod Sci*. 2011;18:1080-1084.
15. Yamagata Y, Nishino K, Takaki E, et al. Genome-wide DNA methylation profiling in cultured eutopic and ectopic endometrial stromal cells. *PLoS ONE*. 2014;9:e83612.
16. Wu Y, Strawn E, Basir Z, Halverson G, Guo SW. Promoter hypermethylation of progesterone receptor isoform B (PR-B) in endometriosis. *Epigenetics*. 2006;1:106-111.
17. Wu Y, Starzinski-Powitz A, Guo SW. Prolonged stimulation with tumor necrosis factor-alpha induced partial methylation at PR-B promoter in immortalized epithelial-like endometriotic cells. *Fertil Steril*. 2008;90:234-237.
18. Wu M, Lu C, Chuang P, Tsai S. Prostaglandin E2: the master of endometriosis? *Exp Biol Med*. 2010;235:668-677.
19. Sun HS, Hsiao KY, Hsu CC, Wu MH, Tsai SJ. Transactivation of steroidogenic acute regulatory protein in human endometriotic stromal cells is mediated by the prostaglandin EP2 receptor. *Endocrinology*. 2003;144:3934-3942.

20. Colon-Diaz M, Baez-Vega P, Garcia M, et al. HDAC1 and HDAC2 are differentially expressed in endometriosis. *Reprod Sci*. 2012;19:483-492.
21. Xia M, Zhao M, Ma J, Fang X. Aberrant histone acetylation and methylation levels in woman with endometriosis. *Arch Gynecol Obstet*. 2013;287:487-494.
22. Samartzis EP, Noske A, Samartzis N, Fink D, Imesch P. The expression of histone deacetylase 1, but not other class I histone deacetylases, is significantly increased in endometriosis. *Reprod Sci*. 2013;20:1416-1422.
23. Monteiro JB, Colon-Diaz M, Garcia M, et al. Endometriosis is characterized by a distinct pattern of histone 3 and histone 4 lysine modifications. *Reprod Sci*. 2014;21:305-318.
24. Pesavento JJ, Bullock CR, LeDuc RD, Mizzen CA, Kelleher NL. Combinatorial modification of human histone H4 quantitated by two-dimensional liquid chromatography coupled with top down mass spectrometry. *J Biol Chem*. 2008;283:14927-14937.
25. Thomas CE, Kelleher NL, Mizzen CA. Mass spectrometric characterization of human histone H3: a bird's eye view. *J Proteome Res*. 2006;5:240-247.
26. Kawano Y, Nasu K, Hijiya N, et al. CCAAT/enhancer-binding protein alpha is epigenetically silenced by histone deacetylation in endometriosis and promotes the pathogenesis of endometriosis. *J Clin Endocrinol Metab*. 2013;98:E1474-E1482.
27. Han SJ, Hawkins SM, Begum K, et al. A new isoform of steroid receptor coactivator-1 is crucial for pathogenic progression of endometriosis. *Nat Med*. 2012;18:1102-1111.
28. Hanstein B, Eckner R, DiRenzo J, et al. p300 is a component of an estrogen receptor coactivator complex. *Proc Natl Acad Sci USA*. 1996;93:11540-11545.
29. Wu Y, Guo SW. Histone deacetylase inhibitors trichostatin A and valproic acid induce cell cycle arrest and p21 expression in immortalized human endometrial stromal cells. *Eur J Obstet Gynecol Reprod Biol*. 2008;137:198-203.
30. Imesch P, Fink D, Fedier A. Romidepsin reduces histone deacetylase activity, induces acetylation of histones, inhibits proliferation, and activates apoptosis in immortalized epithelial endometriotic cells. *Fertil Steril*. 2010;94:2838-2842.
31. Kai K, Nasu K, Kawano Y, et al. Death receptor 6 is epigenetically silenced by histone deacetylation in endometriosis and promotes the pathogenesis of endometriosis. *Am J Reprod Immunol*. 2013;70:485-496.
32. Hsiao KY, Mizzen CA. Histone H4 deacetylation facilitates 53BP1 DNA damage signaling and double-strand break repair. *J Mol Cell Biol*. 2013;5:157-165.
33. Tang J, Cho NW, Cui G, et al. Acetylation limits 53BP1 association with damaged chromatin to promote homologous recombination. *Nat Struct Mol Biol*. 2013;20:317-325.
34. Kruhlak MJ, Hendzel MJ, Fischle W, et al. Regulation of global acetylation in mitosis through loss of histone acetyltransferases and deacetylases from chromatin. *J Biol Chem*. 2001;276:38307-38319.
35. Cimini D, Mattiuzzo M, Torosantucci L, Degraffi F. Histone hyperacetylation in mitosis prevents sister chromatid separation and produces chromosome segregation defects. *Mol Biol Cell*. 2003;14:3821-3833.
36. Catalanotto C, Cogoni C, Zardo G. MicroRNA in control of gene expression: an overview of nuclear functions. *Int J Mol Sci*. 2016;17:E1712.
37. Braza-Boils A, Mari-Alexandre J, Gilabert J, et al. MicroRNA expression profile in endometriosis: its relation to angiogenesis and fibrinolytic factors. *Hum Reprod*. 2014;29:978-988.
38. Hirakawa T, Nasu K, Abe W, et al. miR-503, a microRNA epigenetically repressed in endometriosis, induces apoptosis and cell-cycle arrest and inhibits cell proliferation, angiogenesis, and contractility of human ovarian endometriotic stromal cells. *Hum Reprod*. 2016;31:2587-2597.
39. Shen L, Yang S, Huang W, et al. MicroRNA23a and microRNA23b deregulation derepresses SF-1 and upregulates estrogen signaling in ovarian endometriosis. *J Clin Endocrinol Metab*. 2013;98:1575-1582.
40. Campos-Viguri GE, Jimenez-Wences H, Peralta-Zaragoza O, et al. miR-23b as a potential tumor suppressor and its regulation by DNA methylation in cervical cancer. *Infect Agent Cancer*. 2015;10:42.
41. Lin SC, Wang CC, Wu MH, Yang SH, Li YH, Tsai SJ. Hypoxia-induced microRNA-20a expression increases ERK phosphorylation and angiogenic gene expression in endometriotic stromal cells. *J Clin Endocrinol Metab*. 2012;97:E1515-E1523.
42. Lin SC, Li YH, Wu MH, et al. Suppression of COUP-TFII by proinflammatory cytokines contributes to the pathogenesis of endometriosis. *J Clin Endocrinol Metab*. 2014;99:E427-E437.
43. Abe W, Nasu K, Nakada C, Kawano Y, Moriyama M, Narahara H. miR-196b targets c-myc and Bcl-2 expression, inhibits proliferation and induces apoptosis in endometriotic stromal cells. *Hum Reprod*. 2013;28:750-761.

How to cite this article: Hsiao K-Y, Wu M-H, Tsai S-J.

Epigenetic regulation of the pathological process in endometriosis. *Reprod Med Biol*. 2017;16:314-319.

<https://doi.org/10.1002/rmb2.12047>