

A group theoretical approach to graviton two-point function

S. Rahbardehghan¹, H. Pejhan^{2,a}, M. Elmizadeh²

¹ Department of Physics, Islamic Azad University, Central Branch, Tehran, Iran

² Department of Physics, Science and Research Branch, Islamic Azad University, Tehran, Iran

Received: 23 January 2015 / Accepted: 27 February 2015 / Published online: 13 March 2015

© The Author(s) 2015. This article is published with open access at Springerlink.com

Abstract Respecting the group theoretical approach, it is debated that the theory of linear conformal gravity should be formulated through a tensor field of rank-3 and mixed symmetry (Binegar et al., *Phys Rev D* 27: 2249, 1983). Pursuing this path, such a field equation was obtained in de Sitter space (Takook et al., *J Math Phys* 51:032503, 2010). In the present work, considering the de Sitter ambient space notation, a proper solution to the physical part of this field equation is obtained. We have also calculated the related two-point function, which is interestingly de Sitter invariant and free of an infrared divergence.

1 Introduction

Many people believe that conformal invariance may be the key to a future theory of quantum gravity. In this paper, we consider linear theories of gravitation, in which not only the field equations but also the free field commutation relations are conformal invariant. The main input into this construction of linear gravity is to insist that the propagating modes must be a pair of massless particles with helicity ± 2 . It was supposed that a natural choice for such a field is a symmetric tensor field of rank-2. However, as proved in Ref. [3], for the physical representation of conformal group, the value of conformal Casimir operator is 9. However, by considering a rank-2 tensor field, the related value will become 8 [1]. Hence, such a tensor field does not correspond to any unitary irreducible representation (UIR) of the conformal group. Indeed, the mentioned physical requirement implies that the theory of linear conformal quantum gravity must be formulated in terms of a tensor field of rank-3 and mixed symmetry with conformal degree zero [1]. The mixed symmetry means that

$$\Psi_{abc} = -\Psi_{bac}, \quad \sum_{\text{cycl}} \Psi_{abc} = 0,$$

while a field of conformal degree zero satisfies $u^d \partial_d \Psi_{abc} = 0$.

On the other hand, according to the Wigner theorem, a linear gravitational field should transform under the UIR of its space-time symmetry group. In this regard, it seems that the theory should also be invariant under the de Sitter (dS) group as the space-time symmetry group. Our choice of dS space-time is due to the recent cosmological observations. These observational data are strongly in favor of a positive acceleration of the present universe [4–7], which means, in the first approximation, our universe might currently be in a dS phase. Accordingly, a mixed symmetry tensor field of rank-3 with conformal degree zero, which transforms according to both UIRs of the conformal and de Sitter groups, was obtained in Ref. [2]. In the present work, a proper solution for the physical part of this conformal field equation is calculated. Then the related conformally invariant (CI) two-point function is obtained. It is, interestingly, de Sitter invariant and free of any pathological large-distance behavior. Our method to calculate the two-point function is based on a rigorous group theoretical approach combined with a suitable adaption of Krein space quantization.

This Krein quantization method is a canonical quantization of Gupta–Bleuler type in which the Fock space is constructed over the total space $\mathcal{H}_+ \oplus \mathcal{H}_-$, where \mathcal{H}_+ (\mathcal{H}_-) stands for the Hilbert (anti-Hilbert) space [8–10]. Through this construction, recently, a covariant quantization of the massless minimally coupled scalar field on de Sitter space has been carried out [11, 12]; according to Allen’s theorem [13, 14], no invariant vacuum exists, therefore no covariant Hilbert space quantization is possible. It is reputed that the graviton propagator in the linear approximation on dS background suffers from the same problem. Actually, for largely separated points, it has a pathological behavior (infrared divergence) and also breaks the de Sitter invariance [15–17].¹

^a e-mails: h.pejhan@yahoo.com; h.pejhan@piaou.ac.ir

¹ On this basis, it has been proposed that the infrared divergence might lead to instability of de Sitter space [18, 19]. So, some authors, by

Respecting the Krein quantization method, however, these difficulties are solved. The singularity of the Wightman two-point function, which appears due to the zero mode problem of the Laplace–Beltrami operator on dS space [13,14], is removed, and interestingly the de Sitter invariance survives.

The layout of the paper is as follows. In Sect. 2, we briefly introduce the notations, and in particular, study the CI massless spin-2 wave equations in dS space. In Sect. 3, by focusing on the physical part of the field equations, the corresponding solution is calculated. It is actually constructed over the massless minimally coupled scalar field. In Sect. 4, we calculate the two-point function $\mathcal{W}_{\alpha\beta\gamma\alpha'\beta'\gamma'}(x, x')$ in the ambient space notations. Especially, it is shown that, through Krein space quantization, we are capable of calculating the physical graviton two-point function that is dS-invariant and free of any divergences. Finally, in Sect. 5, the results of the paper are discussed. Some mathematical relations are given in the appendices.

2 De Sitter space and Dirac’s six-cone formalism

2.1 De Sitter space

The de Sitter solution to the cosmological Einstein field equation (with positive cosmological constant Λ) can be viewed as a one-sheeted hyperboloid embedded in a 5-dimensional Minkowski space M^5

$$X_H = \left\{ x \in R^5; x^2 = \eta_{\alpha\beta}x^\alpha x^\beta = -H^{-2} = -\frac{3}{\Lambda} \right\},$$

$$\alpha, \beta = 0, 1, 2, 3, 4, \tag{2.1}$$

where $\eta_{\alpha\beta} = \text{diag}(1, -1, -1, -1, -1)$ and H is the Hubble parameter. The dS metric is

$$ds^2 = \eta_{\alpha\beta}dx^\alpha dx^\beta = g_{\mu\nu}^{\text{dS}}dX^\mu dX^\nu, \quad \mu, \nu = 0, 1, 2, 3.$$

We use x^α for ambient space formalism (five global coordinates) whereas X^μ stand for de Sitter intrinsic coordinates (four local coordinates). In what follows, the ambient space notation is used, because working in the embedding space has two advantages, first it is close to the group theoretical language and second the equations are obtained in an easier way than they might be found in de Sitter intrinsic space.

The dS kinematical group is the 10-parameter group $SO_0(1, 4)$ (connected component of the identity in $O(1, 4)$),

Footnote 1 continued

considering a dS field operator for linear gravity in terms of flat coordinates (it covers only one-half of the de Sitter hyperboloid), have investigated the possibility of quantum instability and have found a quantum field, which violates the de Sitter invariance [20,21]. However, recently it was shown that the infrared divergence of the graviton propagator in the one-loop approximation is gauge dependent, therefore, it should not appear in an effective way as a physical quantity [22–25].

for which there are two Casimir operators,

$$Q^{(1)} = -\frac{1}{2}L^{\alpha\beta}L_{\alpha\beta}, \quad Q^{(2)} = -W_\alpha W^\alpha, \tag{2.2}$$

where $W_\alpha = -\frac{1}{8}\epsilon_{\alpha\beta\gamma\sigma\eta}L^{\beta\gamma}L^{\sigma\eta}$ and $\epsilon_{\alpha\beta\gamma\sigma\eta}$ is the antisymmetric tensor in the ambient space notation with $\epsilon_{01234} = 1$. The generator of the de Sitter group is $L_{\alpha\beta} = M_{\alpha\beta} + \sum_{\alpha\beta}$, in which the action of the orbital, $M_{\alpha\beta}$, and the spinorial, $\sum_{\alpha\beta}$, parts are, respectively, defined by [26,27]

$$M_{\alpha\beta} \equiv -i(x_\alpha\partial_\beta - x_\beta\partial_\alpha) = -i(x_\alpha\bar{\partial}_\beta - x_\beta\bar{\partial}_\alpha),$$

$$\sum_{\alpha\beta} \mathcal{K}_{\gamma\delta\dots} \equiv -i(\eta_{\alpha\gamma}\mathcal{K}_{\beta\delta\dots} - \eta_{\beta\gamma}\mathcal{K}_{\alpha\delta\dots} + \eta_{\alpha\delta}\mathcal{K}_{\gamma\beta\dots} - \eta_{\beta\delta}\mathcal{K}_{\gamma\alpha\dots} + \dots). \tag{2.3}$$

$\bar{\partial}_\alpha$ is the tangential (or transverse) derivative on dS space, defined by

$$\bar{\partial}_\alpha = \theta_{\alpha\beta}\partial^\beta = \partial_\alpha + H^2x_\alpha x \cdot \partial, \quad \text{with } x \cdot \bar{\partial} = 0, \tag{2.4}$$

and $\theta_{\alpha\beta}$ is the transverse projector ($\theta_{\alpha\beta} = \eta_{\alpha\beta} + H^2x_\alpha x_\beta$).

The operator $Q^{(1)}$ commutes with the action of the group generators; thus, it is constant in each UIR. The eigenvalues of $Q^{(1)}$ can be used to classify the UIRs, i.e.

$$(Q^{(1)} - \langle Q^{(1)} \rangle)\mathcal{K}(x) = 0. \tag{2.5}$$

Following Dixmier [28], one can get a classification scheme considering a pair (p, q) of parameters involved in the following possible spectral values of the Casimir operators:

$$Q^{(1)} = (-p(p+1) - (q+1)(q-2))I_d,$$

$$Q^{(2)} = (-p(p+1)q(q-1))I_d. \tag{2.6}$$

According to the range of values of the parameters p and q , there exist three distinct types of UIRs for $SO(1, 4)$ [28,29], namely: principal, complementary and discrete series. In the case of the principal and complementary series, the flat limit compels the value of p to bear the meaning of spin. For the discrete series case, the only representation which has a physically meaningful Minkowskian counterpart is $p = q$ case. For more mathematical details of the group contraction and the physical principles underlying the relationship between dS and Poincaré groups, one can refer to Refs. [30,31].

The spin-2 tensor representations relevant to the present work are as follows:

- (I) The UIRs $U^{2,\nu}$ in the principal series, $p = s = 2$ and $q = \frac{1}{2} + i\nu$, correspond to the Casimir spectral values

$$\langle Q^{(1)} \rangle = \nu^2 - \frac{15}{4}, \quad \nu \in R, \tag{2.7}$$

in which $U^{2,\nu}$ and $U^{2,-\nu}$ are equivalent.

(II) The UIRs $V^{2,q}$ in the complementary series, $p = s = 2$ and $q - q^2 = \mu$, correspond to

$$\langle Q^\mu \rangle = q - q^2 - 4 \equiv \mu - 4, \quad 0 < \mu < \frac{1}{4}. \quad (2.8)$$

(III) The UIRs $\Pi_{2,q}^\pm$ in the discrete series, $p = s = 2$, correspond to

$$\begin{aligned} \langle Q^{(1)} \rangle &= -4, \quad q = 1 \ (\Pi_{2,1}^\pm); \\ \langle Q^{(2)} \rangle &= -6, \quad q = 2 \ (\Pi_{2,2}^\pm). \end{aligned} \quad (2.9)$$

Regarding the de Sitter group, the ‘‘massless’’² spin-2 field is symbolized by $\Pi_{2,2}^\pm$ and $\Pi_{2,1}^\pm$ (the signs \pm correspond to the two types of helicity for the massless tensor field). In these cases, the two representations $\Pi_{2,2}^\pm$, in the discrete series with $p = q = 2$, have a Minkowskian interpretation. It is worth to mention that p and q do not bear the meaning of mass and spin. For discrete series in the limit $H \rightarrow 0$, $p = q = s$ are veritably none other than spin.

The compact subgroup of the conformal group $SO(2, 4)$ is $SO(2) \otimes SO(4)$, in which, by considering E as the eigenvalues of the conformal energy generator of $SO(2)$ and (j_1, j_2) as the $(2j_1 + 1)(2j_2 + 1)$ -dimensional representation of $SO(4) = SU(2) \otimes SU(2)$, the mathematical symbols $C(E; j_1, j_2)$ can be used to denote the irreducible projective representation of the conformal group. The representation $\Pi_{2,2}^+$ has a unique extension to a direct sum of two UIRs $C(3; 2, 0)$ and $C(-3; 2, 0)$ of the conformal group, with positive and negative energies respectively [3, 30]. The latter is restricted to the massless Poincaré UIRs $P^>(0, 2)$ and $P^<(0, 2)$ with positive and negative energies, respectively. $\mathcal{P}^>(0, 2)$ (resp. $\mathcal{P}^<(0, -2)$) are the massless Poincaré UIRs with positive and negative energies and positive (resp. negative) helicity. The following diagrams elucidate these connections:

$$\begin{array}{ccc} \Pi_{2,2}^+ \hookrightarrow & \begin{array}{ccc} C(3, 2, 0) & C(3, 2, 0) & \leftrightarrow \mathcal{P}^>(0, 2) \\ \oplus & \xrightarrow{H=0} & \oplus & \oplus \\ C(-3, 2, 0) & C(-3, 2, 0) & \leftrightarrow \mathcal{P}^<(0, 2), \end{array} \end{array} \quad (2.10)$$

$$\begin{array}{ccc} \Pi_{2,2}^- \hookrightarrow & \begin{array}{ccc} C(3, 0, 2) & C(3, 0, 2) & \leftrightarrow \mathcal{P}^>(0, -2) \\ \oplus & \xrightarrow{H=0} & \oplus & \oplus \\ C(-3, 0, 2) & C(-3, 0, 2) & \leftrightarrow \mathcal{P}^<(0, -2), \end{array} \end{array} \quad (2.11)$$

² It should be noted that in de Sitter space, the mass concept does not exist by itself as a conserved quantity. It is actually referred to the conformal invariance (propagating on the dS light cone). The term ‘‘massive’’, however, is used in reference to fields that in the flat limit would be reduced to massive Minkowskian fields [3].

the arrows \hookrightarrow indicate a unique extension. It is worth to mention that the representations $\Pi_{2,1}^\pm$ do not have a corresponding zero curvature limit [30, 31].

2.2 Dirac’s six-cone formalism and conformal-invariant field equations

The concept of conformal space and six-cone formalism was firstly used by Dirac to obtain the field equations for spinor and vector fields in 1 + 3-dimensional space-time in the conformally covariant form [32]. He suggested a manifestly CI formulation in which the Minkowski coordinates are embedded as the hypersurface $\eta_{ab}u^a u^b = 0$, ($a, b = 0, 1, 2, 3, 4, 5$), $\eta_{ab} = \text{diag}(1, -1, -1, -1, -1, 1)$ in R^6 . Then the fields are extended by homogeneity requirements to the whole of the space of homogeneous coordinates, namely R^6 . The reduction to four dimensions (physical space-time) is carried out by projection, that is, by fixing the degrees of homogeneity of all fields. Wave equations, subsidiary conditions, etc., must be expressed in terms of well-defined operators which are determined intrinsically on the cone (they actually map tensor fields to tensor fields with the same rank on the cone $u^2 = 0$). Thus, the equations obtained through this method are conformally invariant. This approach to the conformal symmetry was then developed by Mack and Salam [33] and many others [34, 35].

Considering this method in de Sitter space provides us with the opportunity to acquire the CI field equations for massless scalar, vector and tensor fields [2, 36, 37]. It has been shown that these CI equations in the zero curvature limit ($H \rightarrow 0$) would be reduced exactly to their Minkowskian counterparts, e.g., the Maxwell equations are obtained from the vector field case [36, 37].

As discussed in Sect. 1, we are interested in the conformal invariance properties of massless spin-2 field in dS space, i.e. the dS linear gravity. Generalizing the group theoretical approach, based on proposals by Binegar et al. [1], to de Sitter space and using a mixed symmetry tensor field of rank-3 with conformal degree zero, the related CI wave equation in dS space is best obtained as follows [2]³:

$$\begin{aligned} 2Q_0^{(1)}(Q_0^{(1)} - 2)(F_{\alpha\beta\gamma} - \frac{1}{4}x_\gamma \mathcal{A}_{\alpha\beta}) + (\bar{\partial}_\alpha + 3x_\alpha)(Q_0^{(1)} - 2) \\ \times (4\bar{\partial} \cdot F_{\beta\gamma} - \mathcal{A}_{\gamma\beta} - x_\gamma \bar{\partial} \cdot \mathcal{A}_\beta) \\ + (\bar{\partial}_\beta + 3x_\beta)(Q_0^{(1)} - 2) \\ \times (4\bar{\partial} \cdot F_{\alpha\gamma} - \mathcal{A}_{\alpha\gamma} - x_\gamma \bar{\partial} \cdot \mathcal{A}_\alpha) = 0, \end{aligned} \quad (2.12)$$

in which $Q_0^{(1)} = -\frac{1}{2}M^{\alpha\beta}M_{\alpha\beta}$, $F_{\alpha\beta\gamma}$ is the projected tensor field to dS space and $\mathcal{A}_{\alpha\beta} \equiv \bar{\partial}^\gamma F_{\alpha\beta\gamma} - x_\alpha F_{\gamma\beta}^\gamma + x_\beta F_{\gamma\alpha}^\gamma$. Now,

³ Note: for the sake of simplicity, from now on we take $H = 1$ and use the notation $\bar{\partial}^\alpha F_{\alpha\beta\gamma} \equiv \bar{\partial} \cdot F_{\beta\gamma}$.

by imposing the mixed symmetry, transversality, divergenceless, and traceless conditions on the tensor field $F_{\alpha\beta\gamma}$, which are necessary for UIRs of the dS and conformal groups, the CI equation (2.12) reduces to (see Appendix A)

$$\begin{aligned} Q_0^{(1)}(Q_0^{(1)} - 2)F_{\alpha\beta\gamma} &= 0, \quad \text{or equivalently,} \\ (Q^{(1)} + 6)(Q^{(1)} + 4)F_{\alpha\beta\gamma} &= 0. \end{aligned} \tag{2.13}$$

Obviously this CI field corresponds to the two representations of discrete series, $\Pi_{2,1}^\pm$ and $\Pi_{2,2}^\pm$ (the physical representation of the de Sitter group). Accordingly, the parameter p does have a physical significance. It is indeed spin. In what follows, however, we are only interested in the tensor field that corresponds to the representations of $\Pi_{2,2}^\pm$, i.e.

$$(Q^{(1)} + 6)F_{\alpha\beta\gamma} = 0. \tag{2.14}$$

As already pointed out, these are actually the only two representations in the discrete series which have a Minkowskian interpretation.

3 De Sitter field solution

In this section, we want to obtain solution of the physical part of the CI field equation. To start, we consider the most generic form of $F_{\alpha\beta\gamma}$ as follows:

$$F_{\alpha\beta\gamma} = (\bar{\partial}_\alpha + x_\alpha)K_{\beta\gamma} - (\bar{\partial}_\beta + x_\beta)K_{\alpha\gamma} + \bar{Z}_\alpha H_{\beta\gamma} - \bar{Z}_\beta H_{\alpha\gamma}, \tag{3.1}$$

where $K_{\alpha\beta}$ and $H_{\alpha\beta}$ are two rank-2 tensor fields and Z is a 5-dimensional constant vector. Bar over the vector makes it a tangential (or transverse) vector on dS space (see (2.4)). Imposing the mixed symmetry, transversality, divergenceless, and traceless conditions on $F_{\alpha\beta\gamma}$, which are needed in order to relate it to the physical representation, leads to

$$\begin{aligned} K_{\alpha\beta} &= K_{\beta\alpha}, \quad x \cdot K_{\cdot\beta} = x \cdot K_{\alpha\cdot} = 0, \\ H_{\alpha\beta} &= H_{\beta\alpha}, \quad x \cdot H_{\cdot\beta} = x \cdot H_{\alpha\cdot} = 0, \\ \bar{\partial} \cdot H_{\cdot\beta} &= \bar{\partial} \cdot H_{\alpha\cdot} = 0, \quad \mathcal{H}' = 0, \end{aligned} \tag{3.2}$$

where $\mathcal{H}' = H_\alpha^\alpha$ is the trace of $H_{\alpha\beta}$. In addition one obtains useful relations as follows:

$$\begin{aligned} (Q_0^{(1)} - 2)K_{\beta\gamma} + (\bar{\partial}_\beta + 2x_\beta)\bar{\partial} \cdot K_{\cdot\gamma} - (Z \cdot \bar{\partial} + 3x \cdot Z)H_{\beta\gamma} \\ + x_\beta Z \cdot H_{\cdot\gamma} &= 0, \quad \text{(I)} \\ (\bar{\partial}_\alpha + 2x_\alpha)\bar{\partial} \cdot K_{\cdot\beta} - (\bar{\partial}_\beta + 2x_\beta)\bar{\partial} \cdot K_{\alpha\cdot} + x_\alpha Z \cdot H_{\cdot\beta} \\ - x_\beta Z \cdot H_{\alpha\cdot} &= 0, \quad \text{(II)} \\ (\bar{\partial}_\alpha + x_\alpha)\mathcal{K}' - \bar{\partial} \cdot K_{\alpha\cdot} \\ - Z \cdot H_{\alpha\cdot} &= 0, \quad \text{(III)} \end{aligned} \tag{3.3}$$

$\mathcal{K}' = K_\alpha^\alpha$ is the trace of $K_{\alpha\beta}$.

On the other hand, substituting $F_{\alpha\beta\gamma}$ in (2.14), results in [From now on, in order to get shorthand equations, we define a symmetrizer operator, i.e. $S_{\alpha\beta}K_{\alpha\beta} \equiv K_{\alpha\beta} + K_{\beta\alpha}$, and an anti-symmetrizer operator, i.e. $\bar{S}_{\alpha\beta}K_{\alpha\beta} \equiv K_{\alpha\beta} - K_{\beta\alpha}$.]

$$\begin{cases} \bar{S}_{\alpha\beta}((\bar{\partial}_\alpha + 3x_\alpha)Q_0^{(1)} - 4x_\alpha)K_{\beta\gamma} \\ = \bar{S}_{\alpha\beta}((8x_\alpha + 2\bar{\partial}_\alpha)(x \cdot Z) + 2x_\alpha(Z \cdot \bar{\partial}))H_{\beta\gamma}, \quad \text{(I)} \\ Q_0^{(1)}H_{\beta\gamma} = 0. \quad \text{(II)} \end{cases} \tag{3.4}$$

From Eq. (3.4-I) along with the conditions given in (3.2), (3.3), and after following the procedure given in Appendix B, it is proved that $K_{\beta\gamma}$ can be written in terms of $H_{\beta\gamma}$ as

$$\begin{aligned} K_{\beta\gamma}(x) &= \left(-\frac{1}{2}(x \cdot Z) + \frac{1}{8}(Z \cdot \bar{\partial}) \right) H_{\beta\gamma} \\ &\quad - \frac{1}{8}(x_\beta Z \cdot H_{\cdot\gamma} + x_\gamma Z \cdot H_{\beta\cdot}). \end{aligned} \tag{3.5}$$

Thus we can construct the tensor field (3.1) as follows:

$$\begin{aligned} F_{\alpha\beta\gamma}(x) &= \bar{S}_{\alpha\beta} \left[(\bar{\partial}_\alpha + x_\alpha) \left(-\frac{1}{2}(x \cdot Z) + \frac{1}{8}(Z \cdot \bar{\partial}) \right) \right. \\ &\quad \left. + \bar{Z}_\alpha \right] H_{\beta\gamma} - \frac{1}{8}\bar{S}_{\alpha\beta}(\bar{\partial}_\alpha + x_\alpha) \\ &\quad \times (x_\beta Z \cdot H_{\cdot\gamma} + x_\gamma Z \cdot H_{\beta\cdot}), \end{aligned} \tag{3.6}$$

where $H_{\beta\gamma}$ must satisfy Eq. (3.4)-II. After utilizing a similar procedure, which is given in Ref. [36], it is proved that

$$\begin{aligned} H(x) &= \left[-\frac{2}{3}\theta Z_1 \cdot + S\bar{Z}_1 + \frac{1}{3}S(\bar{\partial} - x) \left(\frac{1}{9}\bar{\partial} Z_1 \cdot + x \cdot Z_1 \right) \right] \\ &\quad \times \left[\bar{Z}_2 - \frac{1}{2}\bar{\partial} (Z_2 \cdot \bar{\partial} + 2x \cdot Z_2) \right] \phi, \end{aligned} \tag{3.7}$$

Z_1 , Z_2 , and Z_3 are another 5-dimensional constant vectors and ϕ is the massless minimally coupled scalar field.

Now, let us briefly describe Krein quantization of the massless minimally coupled scalar field in de Sitter space

$$\square_H \phi(x) = 0, \tag{3.8}$$

where \square_H is the Laplace–Beltrami operator on de Sitter space. Considering the de Sitter ambient space notation, the solution of (3.8) can be written in terms of the dS plane wave as [38]

$$\phi(x) = (x \cdot \xi)^\sigma, \quad \sigma = 0, -3$$

where ξ lies on the positive null cone $C^+ = \{\xi \in \mathbb{R}^5; \xi^2 = 0, \xi^0 > 0\}$. As already mentioned, Allen proved that, for the dS minimally coupled massless scalar field, the covariant canonical quantization cannot be constructed over the Hilbert space [13, 14]. Actually, in this case, due to the zero mode problem (or constant solution $\sigma = 0$), the constructed Fock space over the Hilbert space (generated by any complete set of modes including the zero mode, $\mathcal{H}_+ = \{\sum_{k \geq 0} \alpha_k \phi_k; \sum_{k \geq 0} |\alpha_k|^2 < \infty\}$, ϕ_k is defined in [11]) is not

de Sitter invariant. More precisely, it is not closed under the action of the dS group. However, by adding all the conjugate modes to the previous ones, a fully covariant quantum field is accessible through a new construction, which is called the Krein space quantization method [11, 12]. In this quantization method, the set of states is different from the set of physical states. Indeed, the observables are defined on the total space $(\mathcal{H}_+ \oplus \mathcal{H}_-)$, while the average values of the observables are calculated on the sub-space of physical states. This provides a remarkable advantage for the theory; we no longer require that the space \mathcal{H}_+ be closed under the action of the isometry group. It is actually enough that the larger space $\mathcal{H}_+ \oplus \mathcal{H}_-$ is closed under the latter, which is clearly a weaker condition. Pursuing this path, it is proved that in the case of the dS minimally coupled massless scalar field—which is of great importance for this paper because the final answer would be written based on it—the Fock space, which is constructed over the total space, is closed under an indecomposable representation of the dS group [11]. It is worth to mention that the total space is equipped with an indefinite inner product which results in some (un-physical) states have a negative norm. [To obtain a detailed construction of the quantization method, and in particular, the unitarity condition and compatibility with (Hilbert space) QFT’s counterpart in the Minkowskian limit, one could refer to Refs. [10,39].]

Accordingly, the field operator is defined as follows:

$$\phi(x) = \frac{1}{\sqrt{2}}[\phi_p(x) + \phi_n(x)], \tag{3.9}$$

in which

$$\begin{aligned} \phi_p(x) &= \sum_{k \geq 0} a_k \phi_k(x) + H.C., \\ \phi_n(x) &= \sum_{k \geq 0} b_k \phi_k^*(x) + H.C.. \end{aligned} \tag{3.10}$$

Here, the positive mode $\phi_p(x)$ is the scalar field that was used by Allen [13, 14]. A significant difference from the standard QFT, which is based on canonical commutation relation, lies in the requirement of the following commutation relations ($|\Omega\rangle$ is the Fock vacuum state):

$$a_k|\Omega\rangle = 0, [a_k, a_{k'}^\dagger] = \delta_{kk'}, \quad b_k|\Omega\rangle = 0, [b_k, b_{k'}^\dagger] = -\delta_{kk'}, \tag{3.11}$$

the other commutation relations are zero. Note that, in spite of the presence of negative norm modes in the theory, no negative energy can be measured; $\langle \mathbf{k}|T_{00}|\mathbf{k}\rangle \geq 0$, for any physical state $|\mathbf{k}\rangle$ and this quantity vanishes if and only if $|\mathbf{k}\rangle = |\Omega\rangle$. Therefore the “normal ordering” process for eliminating the ultraviolet divergence in the vacuum energy, which appears in the usual QFT is not needed [11, 39].

In the following section, respecting the vacuum state $|\Omega\rangle$, the related two-point function is calculated.

4 Two-point function

In this section, we deal with conformally invariant two-point function of the massless spin-2 field. We write the two-point function in dS space in terms of bi-tensors which are called maximally symmetric if they respect dS invariance. Bi-tensors are functions of two points (x, x') and behave like tensors under coordinate transformations at each point [40]. Moreover, the dS axiomatic field theory is constructed over the bi-tensor Wightman two-point function [38, 41]. On this basis, the two-point function is given by

$$\mathcal{W}_{\alpha\beta\gamma\alpha'\beta'\gamma'}(x, x') = \langle \Omega|F_{\alpha\beta\gamma}(x)F_{\alpha'\beta'\gamma'}(x')|\Omega\rangle, \tag{4.1}$$

where $x, x' \in X_H$. In this respect, by considering Eqs. (3.1) and (4.1), the following form for the two-point function is proposed⁴:

$$\begin{aligned} \mathcal{W}_{\alpha\beta\gamma\alpha'\beta'\gamma'}(x, x') &= \bar{S}_{\alpha\beta}(\bar{\partial}_\alpha + x_\alpha)(\bar{S}'_{\alpha'\beta'}(\bar{\partial}'_{\alpha'} \\ &\quad + x'_{\alpha'})\mathcal{W}_{\beta\gamma\beta'\gamma'}^K(x, x') + \bar{S}_{\alpha\beta}\bar{S}'_{\alpha'\beta'} \\ &\quad \times \left((\theta_\alpha \cdot \theta'_{\alpha'})\mathcal{W}_{\beta\gamma\beta'\gamma'}^H(x, x') \right). \end{aligned} \tag{4.2}$$

$\mathcal{W}_{\beta\gamma\beta'\gamma'}^K$ and $\mathcal{W}_{\beta\gamma\beta'\gamma'}^H$ are two transverse bi-tensor two-point functions which will be determined through the similar procedure of the previous section. Actually, the two-point function (4.2) must verify Eq. (2.14) with respect to x and x' (without any difference), and also the physical requirements; mixed symmetry, transversality, divergenceless, and traceless conditions, which imply that

- $\mathcal{W}_{\alpha\beta\gamma\alpha'\beta'\gamma'} = -\mathcal{W}_{\beta\alpha\gamma\alpha'\beta'\gamma'}, \quad \mathcal{W}_{\alpha\beta\gamma\alpha'\beta'\gamma'} = -\mathcal{W}_{\alpha\beta\gamma\beta'\alpha'\gamma'}$.
- $\sum_{\text{cycl}\{\alpha,\beta,\gamma\}} \mathcal{W}_{\alpha\beta\gamma\alpha'\beta'\gamma'} = 0, \quad \sum_{\text{cycl}\{\alpha',\beta',\gamma'\}} \mathcal{W}_{\alpha\beta\gamma\alpha'\beta'\gamma'} = 0$.
- $x \cdot \mathcal{W}_{\beta\gamma\alpha'\beta'\gamma'} = \dots = 0, \quad x' \cdot \mathcal{W}_{\alpha\beta\gamma\beta'\gamma'} = \dots = 0$.
- $\bar{\partial} \cdot \mathcal{W}_{\beta\gamma\alpha'\beta'\gamma'} = \dots = 0, \quad \bar{\partial}' \cdot \mathcal{W}_{\alpha\beta\gamma\beta'\gamma'} = \dots = 0$.
- $\mathcal{W}_{\alpha\beta\alpha'\beta'\gamma'}^\beta = 0, \quad \mathcal{W}_{\alpha\beta\gamma\alpha'\beta'}^{\beta'} = 0$.

At the first step, with regard to the above considerations, we investigate the two-point function (4.2) with the choice of x . Accordingly, by imposing the mentioned requirements on the two-point function, we have

$$\begin{aligned} \mathcal{W}_{\alpha\beta\alpha'\beta'}^K &= \mathcal{W}_{\beta\alpha\alpha'\beta'}^K, \quad x \cdot \mathcal{W}_{\beta\alpha\alpha'\beta'}^K = x \cdot \mathcal{W}_{\alpha\beta\alpha'\beta'}^K = 0, \\ \mathcal{W}_{\alpha\beta\alpha'\beta'}^H &= \mathcal{W}_{\beta\alpha\alpha'\beta'}^H, \quad x \cdot \mathcal{W}_{\beta\alpha\alpha'\beta'}^H = x \cdot \mathcal{W}_{\alpha\beta\alpha'\beta'}^H = 0, \end{aligned}$$

⁴ Note that the primed operators act only on the primed coordinates and vice versa, so that $\bar{\partial}\bar{\partial}' = \bar{\partial}'\bar{\partial}$.

$$\bar{\partial} \cdot \mathcal{W}_{\beta\alpha'\beta'}^H = \bar{\partial} \cdot \mathcal{W}_{\alpha'\beta'}^H = 0, \quad \mathcal{W}_{\alpha\alpha'\beta'}^H = 0, \quad (4.3)$$

and also

$$\begin{aligned} & (\mathcal{Q}_0^{(1)} - 2) \left(\bar{S}'_{\alpha'\beta'} (\bar{\partial}'_{\alpha'} + x'_{\alpha'}) \mathcal{W}_{\beta\gamma\beta'\gamma'}^K \right) + (\bar{\partial}_\beta + 2x_\beta) \\ & \times \left(\bar{S}'_{\alpha'\beta'} (\bar{\partial}'_{\alpha'} + x'_{\alpha'}) \bar{\partial} \cdot \mathcal{W}_{\gamma\beta'\gamma'}^K \right) \\ & - \bar{S}'_{\alpha'\beta'} (\theta'_{\alpha'} \cdot \bar{\partial} + 3x \cdot \theta'_{\alpha'}) \mathcal{W}_{\beta\gamma\beta'\gamma'}^H \\ & + \bar{S}'_{\alpha'\beta'} x_\beta \theta'_{\alpha'} \cdot \mathcal{W}_{\gamma\beta'\gamma'}^H = 0, \\ & \bar{S}_{\alpha\beta} (\bar{\partial}_\alpha + 2x_\alpha) \left(\bar{S}'_{\alpha'\beta'} (\bar{\partial}'_{\alpha'} + x'_{\alpha'}) \bar{\partial} \cdot \mathcal{W}_{\beta\beta'\gamma'}^K \right) \\ & + \bar{S}_{\alpha\beta} x_\alpha \left(\bar{S}'_{\alpha'\beta'} \theta'_{\alpha'} \cdot \mathcal{W}_{\beta\beta'\gamma'}^H \right) = 0, \\ & \times (\bar{\partial}_\alpha + x_\alpha) \left(\bar{S}'_{\alpha'\beta'} (\bar{\partial}'_{\alpha'} + x'_{\alpha'}) \mathcal{W}_{\beta\beta'\gamma'}^K \right) - \bar{S}'_{\alpha'\beta'} \\ & \times (\bar{\partial}'_{\alpha'} + x'_{\alpha'}) \bar{\partial} \cdot \mathcal{W}_{\alpha\beta'\gamma'}^K - \bar{S}'_{\alpha'\beta'} \theta'_{\alpha'} \cdot \mathcal{W}_{\alpha\beta'\gamma'}^H = 0. \end{aligned} \quad (4.4)$$

On the other side, $\mathcal{W}_{\alpha\beta\gamma\alpha'\beta'\gamma'}(x, x')$ must satisfy Eq. (2.14), so one can easily show

$$\begin{aligned} & \bar{S}_{\alpha\beta} \bar{S}'_{\alpha'\beta'} (\bar{\partial}'_{\alpha'} + x'_{\alpha'}) \left((\bar{\partial}_\alpha + 3x_\alpha) \mathcal{Q}_0^{(1)} - 4x_\alpha \right) \mathcal{W}_{\beta\gamma\beta'\gamma'}^K \\ & = \bar{S}_{\alpha\beta} \bar{S}'_{\alpha'\beta'} \left((8x_\alpha + 2\bar{\partial}_\alpha) (x \cdot \theta'_{\alpha'}) + 2x_\alpha (\theta'_{\alpha'} \cdot \bar{\partial}) \right) \mathcal{W}_{\beta\gamma\beta'\gamma'}^H, \end{aligned} \quad (4.5)$$

$$\mathcal{Q}_0^{(1)} \mathcal{W}_{\beta\gamma\beta'\gamma'}^H = 0. \quad (4.6)$$

Consistently with (4.3), (4.4), (4.5), and based on the procedure presented in Sect. 3, it is a matter of a simple calculation to get

$$\begin{aligned} & \bar{S}'_{\alpha'\beta'} (\bar{\partial}'_{\alpha'} + x'_{\alpha'}) \mathcal{W}_{\beta\gamma\beta'\gamma'}^K(x, x') = \bar{S}'_{\alpha'\beta'} \left(-\frac{1}{2} (x \cdot \theta'_{\alpha'}) \right. \\ & \left. + \frac{1}{8} (\theta'_{\alpha'} \cdot \bar{\partial}) \right) \mathcal{W}_{\beta\gamma\beta'\gamma'}^H(x, x') \\ & - \frac{1}{8} \bar{S}'_{\alpha'\beta'} \left(x_\beta \theta'_{\alpha'} \cdot \mathcal{W}_{\gamma\beta'\gamma'}^H(x, x') \right. \\ & \left. + x_\gamma \theta'_{\alpha'} \cdot \mathcal{W}_{\beta\beta'\gamma'}^H(x, x') \right). \end{aligned} \quad (4.7)$$

Then, according to Eqs. (4.2) and (4.7), we have

$$\begin{aligned} \mathcal{W}_{\alpha\beta\gamma\alpha'\beta'\gamma'}(x, x') & = \bar{S}_{\alpha\beta} \bar{S}'_{\alpha'\beta'} \left[(\bar{\partial}_\alpha + x_\alpha) \left(-\frac{1}{2} (x \cdot \theta'_{\alpha'}) \right. \right. \\ & \left. \left. + \frac{1}{8} (\theta'_{\alpha'} \cdot \bar{\partial}) \right) + (\theta_\alpha \cdot \theta'_{\alpha'}) \right] \mathcal{W}_{\beta\gamma\beta'\gamma'}^{\mathcal{H}}(x, x') \\ & - \frac{1}{8} \bar{S}_{\alpha\beta} \bar{S}'_{\alpha'\beta'} (\bar{\partial}_\alpha + x_\alpha) \left(x_\beta \theta'_{\alpha'} \cdot \mathcal{W}_{\gamma\beta'\gamma'}^{\mathcal{H}} \right. \\ & \left. \times (x, x') + x_\gamma \theta'_{\alpha'} \cdot \mathcal{W}_{\beta\beta'\gamma'}^{\mathcal{H}}(x, x') \right), \end{aligned} \quad (4.8)$$

here $\mathcal{W}_{\beta\gamma\beta'\gamma'}^{\mathcal{H}}(x, x')$ applies in the Eq. (4.6). Meanwhile, such transverse function was found in Ref. [36] as

$$\begin{aligned} \mathcal{W}^{\mathcal{H}}(x, x') & = \left(-\frac{2}{3} S' \theta \theta' \cdot + S S' \theta \cdot \theta' \right. \\ & \left. + \frac{1}{3} S S' (\bar{\partial} - x) [x \cdot \theta' + \frac{1}{9} \bar{\partial} \theta' \cdot] \right) \\ & \times \left(\theta \cdot \theta' - \frac{1}{2} \bar{\partial} [\theta' \cdot \bar{\partial} + 2\theta' \cdot x] \right) \mathcal{W}_{mc}(x, x'), \end{aligned} \quad (4.9)$$

\mathcal{W}_{mc} is the two-point function for the minimally coupled massless scalar field in dS space.

Now, at the second step, we investigate the two-point function (4.2) with respect to x' . In this case, the physical requirements imply that

$$\begin{aligned} \mathcal{W}_{\alpha\beta\alpha'\beta'}^{\{K,H\}} & = \mathcal{W}_{\alpha\beta\beta'\alpha'}^{\{K,H\}}, \quad x' \cdot \mathcal{W}_{\alpha\beta\beta'}^{\{K,H\}} = x' \cdot \mathcal{W}_{\alpha\beta\alpha'}^{\{K,H\}} = 0, \\ \bar{\partial}' \cdot \mathcal{W}_{\alpha\beta\beta'}^H & = \bar{\partial}' \cdot \mathcal{W}_{\alpha\beta\alpha'}^H = 0, \quad \mathcal{W}_{\alpha\beta\alpha'}^H = 0, \end{aligned}$$

in addition

$$\begin{aligned} & (\mathcal{Q}'_0^{(1)} - 2) \left(\bar{S}_{\alpha\beta} (\bar{\partial}_\alpha + x_\alpha) \mathcal{W}_{\beta\gamma\beta'\gamma'}^K \right) + (\bar{\partial}'_{\beta'} + 2x'_{\beta'}) \\ & \times \left(\bar{S}_{\alpha\beta} (\bar{\partial}_\alpha + x_\alpha) \bar{\partial}' \cdot \mathcal{W}_{\beta\gamma\beta'}^K \right) \\ & - \bar{S}_{\alpha\beta} (\theta_\alpha \cdot \bar{\partial}' + 3x' \cdot \theta_\alpha) \mathcal{W}_{\beta\gamma\beta'\gamma'}^H \\ & + \bar{S}_{\alpha\beta} x'_{\beta'} \theta_\alpha \cdot \mathcal{W}_{\beta\gamma\beta'\gamma'}^H = 0, \\ & \times \bar{S}'_{\alpha'\beta'} (\bar{\partial}'_{\alpha'} + 2x'_{\alpha'}) \left(\bar{S}_{\alpha\beta} (\bar{\partial}_\alpha + x_\alpha) \bar{\partial}' \cdot \mathcal{W}_{\beta\gamma\beta'}^K \right) \\ & + \bar{S}'_{\alpha'\beta'} x'_{\alpha'} \left(\bar{S}_{\alpha\beta} \theta_\alpha \cdot \mathcal{W}_{\beta\gamma\beta'}^H \right) = 0, \\ & \times (\bar{\partial}'_{\alpha'} + x'_{\alpha'}) \left(\bar{S}_{\alpha\beta} (\bar{\partial}_\alpha + x_\alpha) \mathcal{W}_{\beta\gamma\beta'}^K \right) \\ & - \bar{S}_{\alpha\beta} (\bar{\partial}_\alpha + x_\alpha) \bar{\partial}' \cdot \mathcal{W}_{\beta\gamma\alpha'}^K - \bar{S}_{\alpha\beta} \theta_\alpha \cdot \mathcal{W}_{\beta\gamma\alpha'}^H = 0. \end{aligned} \quad (4.10)$$

Substituting $\mathcal{W}_{\alpha\beta\gamma\alpha'\beta'\gamma'}(x, x')$ into Eq. (2.14) leads to

$$\begin{aligned} & \bar{S}_{\alpha\beta} \bar{S}'_{\alpha'\beta'} (\bar{\partial}_\alpha + x_\alpha) \left((\bar{\partial}'_{\alpha'} + 3x'_{\alpha'}) \mathcal{Q}'_0^{(1)} - 4x'_{\alpha'} \right) \mathcal{W}_{\beta\gamma\beta'\gamma'}^K \\ & = \bar{S}_{\alpha\beta} \bar{S}'_{\alpha'\beta'} \left((8x'_{\alpha'} + 2\bar{\partial}'_{\alpha'}) (x' \cdot \theta_\alpha) + 2x'_{\alpha'} (\theta_\alpha \cdot \bar{\partial}') \right) \mathcal{W}_{\beta\gamma\beta'\gamma'}^H, \\ & \times \mathcal{Q}'_0^{(1)} \mathcal{W}_{\beta\gamma\beta'\gamma'}^H = 0. \end{aligned}$$

As stated so far, it is the work of a few lines to show that

$$\begin{aligned} \mathcal{W}_{\alpha\beta\gamma\alpha'\beta'\gamma'}(x, x') & = \bar{S}_{\alpha\beta} \bar{S}'_{\alpha'\beta'} \left[(\bar{\partial}'_{\alpha'} + x'_{\alpha'}) \left(-\frac{1}{2} (x' \cdot \theta_\alpha) \right. \right. \\ & \left. \left. + \frac{1}{8} (\theta_\alpha \cdot \bar{\partial}') \right) + (\theta'_{\alpha'} \cdot \theta_\alpha) \right] \mathcal{W}_{\beta\gamma\beta'\gamma'}^{\mathcal{H}} \\ & - \frac{1}{8} \bar{S}_{\alpha\beta} \bar{S}'_{\alpha'\beta'} (\bar{\partial}'_{\alpha'} + x'_{\alpha'}) \\ & \times \left(x'_{\beta'} \theta_\alpha \cdot \mathcal{W}_{\beta\gamma\beta'\gamma'}^{\mathcal{H}} + x'_{\gamma'} \theta_\alpha \cdot \mathcal{W}_{\beta\gamma\beta'}^{\mathcal{H}} \right), \end{aligned} \quad (4.11)$$

where $\mathcal{W}_{\beta\gamma\beta'\gamma'}^{\mathcal{H}}$ is [36]

$$\begin{aligned} \mathcal{W}^{\mathcal{H}}(x, x') = & \left(-\frac{2}{3}S\theta'\theta \cdot + S'S\theta' \cdot \theta \right. \\ & \left. + \frac{1}{3}S'S(\bar{\partial}' - x')[x' \cdot \theta + \frac{1}{9}\bar{\partial}'\theta \cdot] \right) \\ & \times \left(\theta' \cdot \theta - \frac{1}{2}\bar{\partial}'[\theta \cdot \bar{\partial}' + 2\theta \cdot x'] \right) \mathcal{W}_{mc}(x, x'). \end{aligned} \tag{4.12}$$

Meanwhile, respecting the Krein construction, the dS minimally coupled massless scalar field two-point function, \mathcal{W}_{mc} , on the vacuum state is [42]

$$\begin{aligned} \mathcal{W}_{mc}(x, x') = & \frac{1}{2}[\langle \Omega | \phi_p(x) \phi_p(x') | \Omega \rangle \\ & + \langle \Omega | \phi_n(x) \phi_n(x') | \Omega \rangle] \\ = & \frac{1}{2}[\mathcal{W}_p(x, x') + \mathcal{W}_n(x, x')], \end{aligned} \tag{4.13}$$

where $\mathcal{W}_n(x, x') = -\mathcal{W}_p^*(x, x')$ and $\mathcal{W}_p(x, x')$ is the two-point function for the positive modes [13, 14, 43],

$$\begin{aligned} \mathcal{W}_p(x, x') = & \frac{1}{8\pi^2} \left[\frac{1}{1 - \mathcal{Z}(x, x')} - \ln(1 - \mathcal{Z}(x, x')) \right. \\ & \left. + \ln 2 + f(\eta, \eta') \right]. \end{aligned} \tag{4.14}$$

It is worth to mention that \mathcal{Z} is an invariant object under the isometry group $O(1, 4)$ which is defined, for two given points on the dS hyperboloid x and x' , by

$$\mathcal{Z} \equiv -x \cdot x' = 1 + \frac{1}{2}(x - x')^2,$$

so that any function of \mathcal{Z} is dS-invariant, as well. f is a function of the conformal time η that breaks the dS invariance. In addition, the term $\ln(1 - \mathcal{Z}(x, x'))$, at largely separated points, is responsible for the advent of the infrared divergence. However, by constructing a covariant quantization of the massless minimally coupled scalar field through Krein space quantization (see (4.13)), we have [42]

$$\begin{aligned} \mathcal{W}_{mc}(x, x') = & \frac{i}{8\pi^2} \epsilon(x^0 - x'^0) \\ & \times [\delta(1 - \mathcal{Z}(x, x')) + \vartheta(\mathcal{Z}(x, x') - 1)], \end{aligned} \tag{4.15}$$

where ϑ is the Heaviside step function and

$$\epsilon(x^0 - x'^0) = \begin{cases} 1 & x^0 > x'^0, \\ 0 & x^0 = x'^0, \\ -1 & x^0 < x'^0. \end{cases} \tag{4.16}$$

Note that this two-point function has been written in terms of \mathcal{Z} , thus the de Sitter invariance is indeed preserved. It is also free of any pathological large-distance behavior.

5 Conclusion

A group theoretical approach to quantum gravity, based on Wigner’s theorem and Dirac’s six-cone formalism, led to the CI field equation for the massless spin-2 field in de Sitter space [2]. In this paper, the corresponding CI two-point function was computed. The calculations were carried out through Krein quantization method. This method has already been successfully applied to the massless minimally coupled scalar field in de Sitter space-time for which it preserves covariance [11, 12]. On this basis, it was shown that the two-point function is dS invariant and also free of any infrared divergences.

At the end, we would like to mention that, although the geometrical interpretation of this linear theory is not entirely clear, it may have an interesting property linked to the quantum approach to the modified gravitational theories, for instance metric-affine theories of gravity. The advent of a rank-3 tensor field implies that, contrary to General Relativity (GR) assumptions, the space-time geometry is not fully described by the metric only, and other geometrical objects which can be independent of metric, such as connections, must be taken into account. In general, the connection does carry dynamics, so that the theory presents more degrees of freedom than GR. Consequently, torsion⁵ does not remain non-propagating [44].

Actually, if we accept that a quantum theory of gravity should be an effective field theory, as many do [45–47], we can conclude remarkable results. It is proved that the torsion is zero in vacuum and in the presence of a scalar field or the electromagnetic field, however, in the presence of a Dirac field or other vector and tensor fields it does not necessarily vanish [44]. This shows a correspondence between torsion and the presence of fields that describe particles with spin. So, though when torsion is present, the concept of a perfect fluid has to be generalized if one wants to include particles with spin, but since many cosmological and astrophysical applications are related to either the vacuum or the environments where matter can more or less be accurately described as a perfect fluid, these contributions to torsion will be negligible in most cases [48]. Therefore, it seems that these dynamical degrees of freedom can be eliminated in low-energy regimes [44],⁶ and still, one can consider the dS space-time as the classical background with good accuracy. Nevertheless, we believe that in high-energy physics, where quantum corrections are important, these effects cannot be ignored. In this respect, the calculated two-point function

⁵ The antisymmetric part of the connection is often called the Cartan torsion tensor.

⁶ It is expected that at some intermediate or high-energy regimes, the spin of particles might interact with the geometry [49].

may have an important role in formulating the future theory of quantum gravity.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.
 Funded by SCOAP³ / License Version CC BY 4.0.

Appendix A: Mathematical relations underlying Eq. (2.13)

Regarding Eqs. (2.2) and (2.3), the action of the Casimir operator $Q^{(1)}$ on a rank-3 tensor field can be written as follows:

$$\begin{aligned}
 Q^{(1)}F_{\alpha\beta\gamma} &= (Q_0^{(1)} - 6)F_{\alpha\beta\gamma} \\
 &+ 2(\eta_{\alpha\beta}F_{\delta\delta\gamma} + \eta_{\beta\gamma}F_{\alpha\delta\delta} + \eta_{\alpha\gamma}F_{\delta\beta\delta}) \\
 &+ 2(x_\alpha\partial \cdot F_{\beta\gamma} + x_\beta\partial \cdot F_{\alpha\gamma} \\
 &+ x_\gamma\partial \cdot F_{\alpha\beta}) - 2(\partial_\alpha x \cdot F_{\beta\gamma} + \partial_\beta x \cdot F_{\alpha\gamma} \\
 &+ \partial_\gamma x \cdot F_{\alpha\beta}) - 2(F_{\beta\alpha\gamma} + F_{\gamma\beta\alpha} + F_{\alpha\gamma\beta}).
 \end{aligned}
 \tag{A.1}$$

It is important to note the following. Impose the following conditions on the tensor field:

- $F_{\alpha\beta\gamma} = -F_{\beta\alpha\gamma}$ and $\sum_{\text{cycl}} F_{\alpha\beta\gamma} = F_{\alpha\beta\gamma} + F_{\beta\gamma\alpha} + F_{\gamma\alpha\beta} = 0$; the mixed symmetry conditions. Note that these conditions are necessary for UIRs of the conformal group [1].
- $x \cdot F_{\beta\gamma} = x \cdot F_{\alpha\gamma} = x \cdot F_{\alpha\beta} = 0$; the transversality conditions.
- $\partial \cdot F_{\beta\gamma} = \partial \cdot F_{\alpha\gamma} = \partial \cdot F_{\alpha\beta} = 0$; the divergenceless conditions. Note that, for transverse tensors, like $F_{\alpha\beta\gamma}$, $\partial \cdot F_{\beta\gamma} = \bar{\partial} \cdot F_{\beta\gamma}$.
- $F_{\alpha\delta\delta} = 0$; the traceless condition.

These are necessary for UIRs of the dS and conformal groups. Then Eq. (A.1) reduces to

$$(Q^{(1)} + 6)F_{\alpha\beta\gamma} = Q_0^{(1)}F_{\alpha\beta\gamma}.$$

For more mathematical details of the action of the Casimir operators ($Q^{(1)}$ and $Q^{(2)}$), the commutation rules and algebraic identities of the various operators and fields, one can refer to [36,50].

Appendix B: Mathematical relations underlying Eq. (3.5)

Generally, the following form for $K_{\beta\gamma}$ can be considered:

$$\begin{aligned}
 K_{\beta\gamma} &= C_1(x \cdot ZH_{\beta\gamma}) + C_2(Z \cdot \bar{\partial}H_{\beta\gamma}) \\
 &+ C_3(\bar{\partial}_\beta Z \cdot H_{\gamma} + \bar{\partial}_\gamma Z \cdot H_{\beta}) + C_4(x_\beta Z \cdot H_{\gamma} \\
 &+ x_\gamma Z \cdot H_{\beta}) + C_5(\theta_{\beta\gamma} Z \cdot H \cdot Z)
 \end{aligned}$$

$$+ C_6(\bar{\partial}_\beta \bar{\partial}_\gamma - x_\gamma \bar{\partial}_\beta)Z \cdot H \cdot Z, \tag{B.1}$$

clearly $K_{\beta\gamma} = K_{\gamma\beta}$. C_1, \dots, C_6 are six arbitrary real numbers, which are determined by considering the following physical requirements:

The transversality conditions ($x \cdot K_{\gamma} = x \cdot K_{\beta} = 0$) require that

$$C_2 + C_3 + C_4 = 0. \tag{B.2}$$

Then the condition (3.3)-III leaves us with

$$C_5 = -C_6, \text{ and } C_1 + 4C_4 + 1 = 0. \tag{B.3}$$

Regarding the conditions (3.3)-I and -II, one can obtain

$$C_1 = -\frac{1}{2}, \quad C_4 = -\frac{1}{8}, \tag{B.4}$$

and also a new auxiliary equation $\bar{\partial}_\beta Z \cdot H_{\gamma} = x_\gamma Z \cdot H_{\beta}$, which states that the third and fourth terms in (B.1) are not independent, so, without any damage to the generality of the solution, one can take $C_3 = 0$. Then we have $C_2 = \frac{1}{8}$ and so one can rewrite the general solution for $K_{\beta\gamma}$ as follows:

$$\begin{aligned}
 K_{\beta\gamma} &= -\frac{1}{2}(x \cdot ZH_{\beta\gamma}) + \frac{1}{8}(Z \cdot \bar{\partial}H_{\beta\gamma}) \\
 &- \frac{1}{8}(x_\beta Z \cdot H_{\gamma} + x_\gamma Z \cdot H_{\beta}) \\
 &+ C_5(\bar{\partial}_\beta x_\gamma - \bar{\partial}_\beta \bar{\partial}_\gamma)Z \cdot H \cdot Z.
 \end{aligned}
 \tag{B.5}$$

Note that a straightforward calculation shows that Eq. (3.4) does not create new constraints to be imposed on (B.5). Therefore, since we are looking for the easiest possible answer, we choose $C_5 = 0$.

References

1. B. Binetgar, C. Fronsdal, W. Heidenreich, Phys. Rev. D. **27**, 2249 (1983)
2. M.V. Takook, M.R. Tanhayi, S. Fatemi, J. Math. Phys. **51**, 032503 (2010)
3. A.O. Barut, A. Böhm, J. Math. Phys. **11**, 2938 (1970)
4. A.G. Riess et al., Supernova Search Team Collaboration, Astro. J. **116**, 1009 (1998)
5. S. Perlmutter et al., Supernova Cosmology Project Collaboration, Astro. J. **517**, 567 (1999)
6. U. Seljak, A. Slosar, P. McDonald, JCAP **014**, 610 (2006)
7. A.G. Riess et al., Astro. J. **98**, 659 (2007)
8. M. Mintchev, J. Phys. A **13**, 1841 (1990)
9. J. Bogner, *Indefinite inner product space* (Springer, Berlin, 1974)
10. T. Garidi, E. Huguët, J. Renaud, J. Phys. A **38**, 245 (2005)
11. J.P. Gazeau, J. Renaud, M.V. Takook, Class. Quant. Grav. **17**, 1415 (2000)
12. S. De Bievre, J. Renaud, Phys. Rev. D **57**, 6230 (1998)
13. B. Allen, Phys. Rev. D **32**, 3136 (1985)
14. B. Allen, A. Folacci, Phys. Rev. D **35**, 3771 (1987)
15. B. Allen, M. Turyn, Nucl. Phys. B **292**, 813 (1987)
16. E.G. Floratos, J. Iliopoulos, T.N. Tomaras, Phys. Lett. B **197**, 373 (1987)
17. I. Antoniadis, E. Mottola, J. Math. Phys. **32**, 1037 (1991)

18. H.L. Ford, Phys. Rev. D **31**, 710 (1985)
19. I. Antoniadis, J. Iliopoulos, T.N. Tomaras, Phys. Rev. Lett. **56**, 1319 (1986)
20. N.C. Tsamis, R.P. Woodard, Phys. Lett. B **292**, 269 (1992)
21. N.C. Tsamis, R.P. Woodard, Commun. Math. Phys. **162**, 217 (1994)
22. I. Antoniadis, J. Iliopoulos, T.N. Tomaras, Nucl. Phys. B **462**, 437 (1996)
23. A. Higuchi, S.S. Kouris, Class. Quant. Grav. **17**, 3077 (2000)
24. A. Higuchi, S.S. Kouris, Class. Quant. Grav. **20**, 3005 (2003)
25. H.J. de Vega, J. Ramirez, N. Sanchez, Phys. Rev. D **60**, 044007 (1999)
26. J.P. Gazeau, Lett. Math. Phys. **8**, 507 (1984)
27. J.P. Gazeau, M. Hans, J. Math. Phys. **29**, 2533 (1988)
28. J. Dixmier, Bull. Soc. Math. France **89**, 9 (1961)
29. B. Takahashi, Bull. Soc. Math. France **91**, 289 (1963)
30. M. Levy-Nahas, J. Math. Phys. **8**, 1211 (1967)
31. H. Bacry, J.M. Levy-Leblond, J. Math. Phys. **9**, 1605 (1968)
32. P.A.M. Dirac, Ann. Math. **36**, 657 (1935)
33. G. Mack, A. Salam, Ann. Phys. **53**, 174 (1969)
34. H.A. Kastrup, Phys. Rev. **150**, 1189 (1964)
35. C.R. Preitschop, M.A. Vosiliev, Nucl. Phys. B. **549**, 450 (1999)
36. M. Dehghani, S. Rouhani, M.V. Takook, M.R. Tanhayi, Phys. Rev. D. **77**, 064028 (2008)
37. S. Behroozi, S. Rouhani, M.V. Takook, M.R. Tanhayi, Phys. Rev. D. **74**, 124014 (2006)
38. J. Bros, U. Moschella, Rev. Math. Phys. **8**, 327 (1996)
39. H. Pejhan, S. Rahbardehghan, Krein quantization approach to Hawking radiation (2015). [arXiv:1408.4531](https://arxiv.org/abs/1408.4531)
40. B. Allen, T. Jacobson, Commun. Math. Phys. **103**, 669 (1986)
41. T. Garidi, J.P. Gazeau, S. Rouhani, M.V. Takook, J. Math. Phys. **49**, 032501 (2008)
42. M.V. Takook, Mod. Phys. Lett. A **16**, 1691 (2001)
43. A. Folacci, J. Math. Phys. **32**, 2828 (1991)
44. V. Vitagliano, T.P. Sotiriou, S. Liberati, Ann. Phys. **326**, 1259–1273 (2011)
45. J.F. Donoghue, Phys. Rev. D. **50**, 3874 (1994)
46. J.F. Donoghue, [arXiv:gr-qc/9512024](https://arxiv.org/abs/gr-qc/9512024) (1995)
47. C.P. Burgess, Living Rev. Relativ. **7**, 5 (2004)
48. T.P. Sotiriou, S. Liberati, Ann. Phys. **322**, 935–966 (2007)
49. T.P. Sotiriou, V. Faraoni, Rev. Mod. Phys. **82**, 451–497 (2010)
50. M.V. Takook, H. Pejhan, M.R. Tanhayi, Eur. Phys. J. C **72**, 2052 (2012)