

Is there any connection between choroidal thickness and obesity?

Farshad Askarizadeh, Mohsen Heirani^{ID}, Masoud Khorrami-Nejad^{ID}, Foroozan Narooie-Noori, Mehdi Khabazkhoob and Alireza Ostadrahimi

Ther Adv Ophthalmol

2022, Vol. 14: 1–10

DOI: 10.1177/
25158414221100649

© The Author(s), 2022.
Article reuse guidelines:
sagepub.com/journals-
permissions

Abstract: Obesity is a health-threatening and epidemic medical condition that can affect individuals of different ages and is potentially associated with an increased risk of systemic and ocular disorders. Despite the well-documented adverse effects of obesity on different parts of the body vasculature, less published data are available concerning obesity-related consequences on the ocular vasculature. As the human choroid is a highly vascularized tissue, its morphology and function might be altered in obese individuals. The micro-structural changes within the choroid could also trigger development of subsequent functional abnormalities of the eye. Previous population-based studies have asserted an association between obesity and choroidal thickness; however, they reported conflicting patterns of association between obesity and changes in choroidal thickness. Therefore, to enhance our understanding of the changes in choroidal morphology secondary to obesity, we reviewed studies describing the micro-structural consequences of obesity on the choroidal thickness profile and its underlying physiological and anatomical basis. This review includes all original publications related to the association between choroidal thickness and obesity published until mid-2021 that were indexed in PubMed, Google Scholar, ScienceDirect, or Scopus.

Keywords: body mass index, choroidal thickness, enhanced depth imaging, obesity, optical coherence tomography

Received: 29 November 2021; revised manuscript accepted: 27 April 2022.

Introduction

Obesity is typically described as excessive fatty tissue accumulation in the body, and is one of the most prominent health concerns worldwide, reaching pandemic levels.^{1,2} Obesity can develop due to the interaction among different etiologies such as genetic, lifestyle, socioeconomic, and environmental factors.^{3–6} Obesity is a serious risk factor for developing various comorbid diseases, including metabolic syndromes (dyslipidemia and insulin resistance), cardiovascular diseases, diabetes mellitus type 2, hypertension, stroke, and sleep apnea.^{7–10} Another negative impact of obesity is the risk of developing depressive disorder, which can contribute to social isolation from peers.¹¹ Therefore, the psychosocial and economic burden of obesity and its comorbidities and coexisting conditions are quite substantial.^{1,3–5,7,8,11,12}

Obesity may affect different structural and functional aspects of the visual system. Previous studies have found serious consequences of obesity for some visual functions, including visual acuity and visual field.¹³ The effects of obesity on ocular anatomy have also been extensively studied. The majority of previously published studies have shown that obese people are at increased risk of glaucomatous manifestations, including higher intraocular pressure, increased cup-to-disk ratio, and thinner retinal nerve fiber layer.¹⁴ However, contradictory results have also been reported.^{15,16} The link between obesity and the risk of cataract formation,^{17,18} age-related macular degeneration,^{19,20} and diabetic retinopathy^{21,22} is also a controversial issue. The incompatible findings of studies on obese individuals may be due to the variations in diet and lifestyle of the participants.

Correspondence to:

Mohsen Heirani
Translational
Ophthalmology Research
Center, Farabi Eye
Hospital, Tehran University
of Medical Sciences,
Qazvin Square, Tehran
1336616351, Iran.
Dr.heirani@hotmail.com

Farshad Askarizadeh
Department of Optometry,
Faculty of Rehabilitation
Sciences, Tabriz University
of Medical Sciences,
Tabriz, Iran

Masoud Khorrami-Nejad
Translational
Ophthalmology Research
Center, Farabi Eye
Hospital, Tehran University
of Medical Sciences,
Tehran, Iran

Foroozan Narooie-Noori
Department of Optometry,
Faculty of Rehabilitation
Sciences, Iran University of
Medical Sciences, Tehran,
Iran

Mehdi Khabazkhoob
Department of Medical
Surgical Nursing, School
of Nursing and Midwifery,
Shahid Beheshti University
of Medical Sciences,
Tehran, Iran

Noor Research Center for
Ophthalmic Epidemiology,
Noor Eye Hospital, Tehran,
Iran

Alireza Ostadrahimi
Nutrition Research Center,
Tabriz University of
Medical Sciences, Tabriz,
Iran

The body hormones of leptin and insulin regulate the appetite and metabolism and distribute the adipose tissue throughout the body.^{23,24} People with obesity have abnormally raised levels of leptin that may stimulate abnormal metabolism and insulin resistance. As a result, it can lead to fatty tissue build-up in the body. Insulin activates glucose uptake and inhibits the release of fatty acids from the adipose tissues; thereby, its abnormal secretion by virtue of the increased body weight may substantially impact ocular vasculature.²⁵ Leptin is produced by fat cells and regulates appetite by sending signals to the central nervous system.²⁶ Previous studies reported the link between leptin level and different posterior segment diseases such as choroidal neovascularization, diabetic retinopathy, and retinal detachment.^{27,28}

It is evident that obesity can be associated with different macrovascular and microvascular dysfunctions.^{8,21,29–33} Notably, the choroid predominantly comprises a dense microvasculature system and is the major blood supplier of the retina.³⁴ Understanding the proposed impacts of obesity on the choroidal structure and profile could be a predictor in identification of obesity-induced ophthalmic complications. The main objective of this study is to review the potential effects of obesity and overweight on choroidal morphology.

Method for literature search

For this literature review, we conducted an extensive search on PubMed, Google Scholar, ScienceDirect, and Scopus search engines for original scientific articles with the following key words in different combinations: obesity and choroid, overweight and choroid, body mass index (BMI) and choroid, obesity and choroidal thickness, BMI, and CT. We retrieved a total of 115 articles published from 1 January 1995, through 1 May 2021, including observational studies, randomized clinical trials (RCTs), systematic reviews, and meta-analyses. A thorough review of all publications in English and the abstracts of non-English papers was undertaken. The reference lists of the selected studies were also checked for potentially relevant papers. Selected sources were also mutually agreed upon by the authors. Irrelevant articles to the main purposes of the study were excluded.

Classification criteria for weight status

BMI is a typical index of weight status and calculates as body weight divided by body height squared (kg/m^2). According to the World Health Organization (WHO) Expert Committee on Physical Status, underweight, normal weight, overweight (pre-obese), obese class I, obese class II, and obese class III (extreme obesity) are defined as BMI values of ≤ 18.50 , 18.50–24.99, 25.00–29.99, 30.00–34.99, 35–39.99, and ≥ 40 , respectively. Morbid obesity is defined as BMI value $> 35 \text{ kg/m}^2$ when the patient experiences obesity-related diseases or BMI values $> 40 \text{ kg/m}^2$ without any coexisting disease. The main limitation of the current WHO classification is that it does not consider the body adipose distribution. In practice, excessive accumulation of fatty tissue in the central or abdominal body part can pose a real risk to the body's metabolic process and circulatory system.^{8,35} Therefore, other indices have been developed to represent the level of central adipose tissue, such as waist circumference and the waist-hip ratio.^{12,36,37} However, most studies applied BMI as a clinically approved criterion for weight-related health problems.^{36,38} This criterion is clear-cut and non-invasive for clinical and study settings.

Effects of obesity on systemic and ocular vasculature

Obesity-related cardiovascular disorder is due to comorbid diseases such as diabetes,³⁹ hypertension,⁴⁰ and stroke.⁴¹ The progression of atherosclerosis in the carotid^{42,43} and femoral⁴³ arteries adversely affects these diseases. Studies on obese individuals determined that higher BMI values might reduce capillary density and tissue perfusion.³⁰ This sign is one of the earliest detectable microvasculature changes in the course of weight gain.^{44,45} Furthermore, obesity causes endothelial defects following hyperglycemia-induced oxidative stress.^{46–48} There are two proposed histopathological theories for microvascular damage. First, vascular endothelial cells demonstrate abnormal synthesis of nitric oxide (NO) molecules following exposure to glycation end-products and stimulate vascular endothelial growth factor (VEGF).⁴⁹ The second theory is attributed to the oxidative stress phenomenon, in which hyperglycemia accelerates the production of reactive oxygen molecules. In consequence, it interacts with proteins and macromolecules like DNA and finally causes damage to the vascular endothelial cells.⁴⁹ In addition to

hyperglycemia, dyslipidemia and inflammatory chemicals may also trigger the development of microvascular disorders.⁵⁰

The normal microcirculation of the two vascular layers of the eye, retina and choroid, could be indicative of the whole-body microvasculature status. Some optical coherence tomography-angiography (OCT-A) studies on obese individuals demonstrated changes in choroidal and retinal microvasculature of these population.^{51,52} For example, Kurtul *et al.*,⁵¹ in a prospective study on obese children, found significantly higher flow area for choriocapillaris and foveal capillary densities than normal healthy subjects. The retinal and choroidal vascular beds are observable locations to directly explore the microvascular system. Notably, any retinal and choroidal vasculature changes could predict the pre-clinical phase of systemic vascular disorders.^{33,44} Previous investigations have revealed direct associations between the retinal venular diameter and BMI measurements.^{32,53} Therefore, widening the retinal venular diameter could also be a clinical sign of systemic venular problems linked to obesity.⁵³

Choroidal thickness changes secondary to obesity

The importance of choroidal thickness evaluation

The human choroid is a highly vascularized structure located behind the globe and extended from the borders of the optic nerve head to the ora serrata. In addition to providing nutrients and oxygen to the sensory and pigmentary retina, the choroid regulates ocular temperature and intraocular pressure.³⁴ The structure and function of the retinal photoreceptors essentially depend on normal choroidal blood flow.³⁴ Because of its hyper-vascularized nature, the choroid is potentially subject to impairment from systemic and local diseases of the body circulatory system.⁵⁴ Therefore, abnormal morphology and malfunction of the choroid can be seen in various ocular and systemic disorders.^{55,56} Assessment of the choroid could thereby manifest obesity-induced microvascular impairments.^{51,54}

Numerous studies on choroidal thickness have addressed its association with age,^{57,58} axial length,⁵⁹ refractive error,^{60,61} and ethnicity.⁶² Thickness changes in choroidal tissue have also been noted in ocular diseases like diabetic retinopathy, age-related macular degeneration, central

serous chorioretinopathy, and retinitis pigmentosa.^{63–66} Moreover, choroidal thickness measurement has diagnostic value in evaluating localized abnormalities of the posterior segment secondary to systemic diseases such as cardiovascular disorders and metabolic syndromes.^{55,56,67} In addition, changes in the choroidal microvasculature could indicate systemic vascular involvement due to obesity.^{32,54} Therefore, it has been hypothesized that choroidal thickness could be considered an indirect predictor of choroidal microvasculature.^{65,68,69} Some studies found the increasing effect of sildenafil citrate on choroidal thickness due to the vasodilatory effect of the drug and consequent rise in choroidal blood flow.^{68,69} Conflicting results have been reported by Sogawa *et al.*,⁷⁰ and they found no significant correlation between subfoveal choroidal thickness (SFCT) measurement and choroidal circulation in healthy young subjects. However, their study had a small sample size and was performed on normal individuals. In addition, the potential effect of associated local and systemic diseases was not investigated.

Measurement of choroidal thickness using the enhanced depth imaging technique

Conventional imaging techniques such as indocyanine green angiography and B-scan ultrasonography represent low image resolution in differentiating posterior layers of the eye.^{71,72} Following the advent of OCT technology and providing *in vivo* investigation of the ocular structures, several studies have analyzed changes in choroidal thickness secondary to systemic conditions.^{56,63,67} The novel OCT techniques (spectral domain) represent higher resolution and less acquisition time because of faster scanning speed compared to the conventional time-domain method. In recent years, cross-sectional and quantitative assessments of choroidal structures have been carried out by adding a modality to spectral-domain OCT known as enhanced depth imaging (EDI).⁷³ In this technique, choroidal thickness is manually outlined as the vertical space between the external border of the hyper-reflective row of the retinal pigment epithelium layer and the internal border of the hypo-reflective choroidoscleral interface layer (Figure 1).⁶² It should be noted that previous studies have indicated that choroidal thickness data recorded by different OCT devices are highly repeatable and have minimum variability.^{73,74} In addition, various OCT instruments are strongly correlated and can be used interchangeably.^{73,74}




Figure 1. Enhanced depth imaging modality of optical coherence tomography representing the anterior (green line) and posterior (red line) borders of the choroid.

The effects of obesity on choroidal thickness

The effect of obesity on choroidal thickness has received limited attention in the literature. Table 1 summarizes the EDI-based studies on the association between SFCT and BMI value.

A number of studies have consistently reported a negative association between SFCT and BMI measurements.^{14,25,54,76–78} In a prospective study on obese children aged 5–15 years, Erşan *et al.*⁵⁴ found a significant reduction in subfoveal and parafoveal choroidal thickness. They recorded baseline blood pressure and performed blood tests (serum insulin, plasma glucose, and lipids) after overnight fasting to control parameters that may affect choroidal thickness. Their findings showed that mean SFCT values in normal weight and age-matched obese children were $301.95 \pm 56.72 \mu\text{m}$, and $270.20 \pm 56.13 \mu\text{m}$, respectively. They ascribed the choroidal thinning to the microvascular alterations in obese participants. Boillot *et al.*²⁹ also supported this hypothesis. They proposed that ocular microvascular changes in obese subjects might be secondary to the arteriole restriction and venular diameter widening. In a cross-sectional study, Yilmaz *et al.*⁷⁶ investigated the association between BMI value and choroidal thickness in the adult population. Their results revealed a negative correlation between SFCT and weight status. They hypothesized that this finding might be due to the impaired dilation of the choroidal vessels following reduced endothelium-derived NO as a vasodilator agent in the obese population.^{32,33} The autonomic system controls choroidal blood flow, and when NO level decreases, parasympathetic control of choroidal circulation is prevented.⁸³ In addition, they found a positive

correlation between obesity and the amount of two main vasoconstrictor molecules, that is, endothelin-1 (ET-1) and angiotensin II (Ang-II).⁸⁴ As a result, the imbalance between obesity-induced vasodilator/vasoconstrictor agents (declining NO level and rising ET-1 and Ang-II) may be associated with choroidal thinning.¹⁴

Hypoxia is another potential underlying factor for obesity-induced choroidal thinning. This condition has been detected in obese individuals and fundamentally has a key role in cell death. Periodic episodes of hypoxia induce oxidative stress and inflammatory reaction. It causes damage to the endothelial cells of the vessel wall, increasing inflammatory cytokines, reducing cell reactivity to vasodilator factors, and stimulating the sympathetic nervous system.^{47,48} These consequences can contribute to dysregulation of the vessels and affect the choroidal circulation, resulting in a decrease in choroidal thickness measurements.⁸⁵ Although some progress has been made, further investigations are needed to prove this hypothesis.

In contrast, several studies reported a positive association between choroidal thickness and obesity.^{75,80,81} Yumusak *et al.*⁸⁰ demonstrated that SFCT was positively correlated with BMI in obese women which might be due to the changes in ophthalmic blood flow. Microvascular damage in obese women might be a complication of elevated blood insulin levels and high blood pressure.³¹ In a cross-sectional study, Bulus *et al.*⁸¹ examined obese children and illustrated a positive correlation between BMI and SFCT. They measured the blood pressure of participants to control the confounding effect of hypertension on CT. Their findings revealed that the mean SFCT values in normal and obese children were $348.43 \pm 73.21 \mu\text{m}$ and $385.77 \pm 6.09 \mu\text{m}$, respectively. The elevated SFCT was attributed to the raised leptin level and overproduction of inflammatory cells in obese children. As discussed earlier, leptin plays a key role in controlling appetite and body weight. Higher leptin production in obese patients than in normal-weight individuals leads to further adipose tissue accumulation.⁸⁶ Available evidence shows that leptin also acts as an angiogenic factor contributing to oxidative stress on endothelial cells.⁸⁷ Conflicting results have been found by Panon *et al.*,⁸² and they found no correlation between SFCT and obesity. These inconsistent outcomes could be explained by various sources such as

Table 1. Studies investigating the association between body mass index and subfoveal choroidal thickness using enhanced depth imaging.

The first author (year of publication)	Number of eyes	Mean age (years)	OCT modality	Association of BMI and SFCT
Nagasawa <i>et al.</i> ⁷⁵	100 pediatrics, 83 adults	7.9 (pediatrics), 54.5 (adults)	Swept source (DRI OCT-1; Topcon)	Negative in both groups
Yilmaz <i>et al.</i> ⁷⁶	160 eyes equally distributed in four groups as group 1 (BMI < 18.50), group 2 (BMI between 18.50 and 24.99), group 3 (BMI between 25 and 29.99), and group 4 (BMI between 30 and 34.99)	26.5 (group 1) 27.2 (group 2) 28.5 (group 3) 29.25 (group 4)	Spectral domain (Spectralis; Heidelberg Engineering)	Negative
Dogan <i>et al.</i> ⁷⁷	67 morbid obese (BMI ≥ 40), 29 healthy subjects	37.92 (morbidly obese) 35.31 (healthy subjects)	Spectral domain (Cirrus HD OCT; Carl Zeiss Meditec)	Negative
Erşan <i>et al.</i> ⁵⁴	40 obese children, 40 non-obese children	11.20 (obese) 11.29 (non-obese)	Spectral domain (Cirrus HD OCT; Carl Zeiss Meditec)	Negative
Özen <i>et al.</i> ⁷⁸	76 obese, 40 healthy subjects	12.8 (obese) 12.9 (healthy subjects)	Spectral domain (Spectralis; Heidelberg Engineering)	Negative
Öner and Karadağ ²⁵	32 obese (BMI > 30), 45 non-obese subjects	41.9 (obese) 36.04 (non-obese)	Spectral domain (Spectralis; Heidelberg Engineering)	Negative
Teberik <i>et al.</i> ¹⁴	101 morbid obese (BMI ≥ 40), 95 healthy subjects	35.9 (morbid obese) 36.6 (healthy subjects)	Spectral domain (Spectralis; Heidelberg Engineering)	Negative
Öncül <i>et al.</i> ⁷⁹	197 eyes of 197 women, in five groups of normal weight (<i>n</i> =44), overweight (<i>n</i> =40), class 1 obese (<i>n</i> =40), class 2 obese (<i>n</i> =38), morbidly obese (<i>n</i> =35)	37.5 (normal weight), 39.4 (overweight), 36 (class 1 obese), 38.7 (class 2 obese), 41.2 (morbidly obese)	Spectral domain (Spectralis; Heidelberg Engineering)	Negative
Yumusak <i>et al.</i> ⁸⁰	72 obese female (BMI > 30), 68 non-obese female (BMI < 25)	37.27 (obese female) 37.85 (non-obese female)	Spectral domain (OCT Advance Nidek RS-3000; Nidek)	Positive
Bulus <i>et al.</i> ⁸¹	44 obese children (BMI > 30), 42 non-obese children	13.3 (obese children) 13.4 (non-obese children)	Spectral domain (Spectralis; Heidelberg Engineering)	Positive
Panon <i>et al.</i> ⁸²	67 overweight (BMI 23.0–29.9), 53 non-obese subjects	47 (overweight) 45.6 (healthy subjects)	Spectral domain (Cirrus HD OCT; Carl Zeiss Meditec)	Not associated
BMI, body mass index (kg/m ²); OCT, optical coherence tomography; SFCT, subfoveal choroidal thickness.				

differences in age and gender of the study population, different choroidal imaging modalities, associated ocular and systemic diseases, and BMI status of the subjects.

Few works have been conducted on choroidal thickness changes in morbidly obese patients.^{14,88} In a study on morbidly obese patients who undertook laparoscopic surgery, Dogan *et al.*⁸⁸ found a

continuous increase in post-operative SFCT at 3- and 6-month follow-ups. In a recent study, Gonul *et al.*⁸⁹ evaluated the choroidal thickness changes in 40 morbidly obese patients following bariatric surgery. They showed a positive association between choroidal thickness and BMI. Terra *et al.*⁹⁰ also found that choroidal thickness decreased gradually along with progressive reduction of BMI following bariatric surgery, which may be due to the mechanical reduction of fatty tissue in orbit as well as the reduction of leptin level after bariatric surgery.

In normal populations, choroidal thickness differs topographically across the choroidal tissue and decreases from the fovea to the outer locations.^{61,78} Furthermore, it has been shown that the superior and temporal areas of the choroid are the thickest quadrants, and the nasal area is significantly the thinnest quadrant across the horizontal choroidal scans.^{91,92} In line with the findings of normal subjects, previous studies on the obese population indicated that choroidal thickness significantly reduced toward nasal regions. These findings show that topographical variations across the choroid of obese people are similar to normal-weight people.^{76,78}

The association between obesity and choroidal thickness might be attributed to the development of obesity comorbidities such as diabetes mellitus, hypertension, dyslipidemia, cardiovascular diseases, stroke, and ischemia.^{7,8,21,22,84} It is hypothesized that microvascular changes secondary to obesity in these associate diseases may contribute to choroidal thickness changes with the same mechanisms in obese patients.^{54,61,65} Several studies have speculated that identification of choroidal thickness changes in obese patients could be an early and noninvasive biomarker of the following vascular changes in the body.^{29,76}

Conclusion

CT variations in obese people can be explained by biological changes in the circulatory system; however, precise mechanism has not been well established. Besides, available data are limited, and therefore many of these findings are inconclusive. In essence, the available literature appears inadequate to explicitly establish the association between choroidal thickness and obesity. It is suggested that future cohort studies examine choroidal thickness in people with a wide age range

from childhood to the elderly. Furthermore, prospective follow-ups of subjects with different stages of obesity, from abnormally underweight to morbid obesity, could help understanding the trend of choroidal thickness changes in different weight conditions. Another challenging aspect of choroidal thickness investigation is the method of measurement. State-of-the-art technologies that incorporate high-resolution OCT modalities having automatic choroidal delineation are recommended to enhance the accuracy of the choroidal image recordings.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Author contribution(s)

Farshad Askarizadeh: Conceptualization; Investigation; Methodology; Project administration; Supervision; Visualization; Writing – original draft; Writing – review & editing.

Mohsen Heirani: Conceptualization; Investigation; Methodology; Project administration; Software; Supervision; Validation; Visualization; Writing – original draft; Writing – review & editing.

Masoud Khorrami-Nejad: Conceptualization; Investigation; Methodology; Supervision; Visualization; Writing – review & editing.


Foroozan Narooie-Noori: Investigation; Visualization; Writing – review & editing.

Mehdi Khabazkhoob: Conceptualization; Supervision; Writing – review & editing.

Alireza Ostadrahimi: Conceptualization; Supervision; Writing – review & editing.

ORCID iDs

Mohsen Heirani  <https://orcid.org/0000-0002-8426-5924>

Masoud Khorrami-Nejad  <https://orcid.org/0000-0002-8270-9704>

Acknowledgements

The authors would like to thank Dr. Asieh Ehsaei and Dr. Javad Heravian for their guidance before starting the research project and Dr. Hosein Hoseini-Yazdi for his valuable advice regarding the related articles in the field of the choroid.

Funding

The authors received no financial support for the research, authorship, and/or publication of this article.

Conflict of interest statement

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Availability of data and materials

Not applicable.

References

- Blüher M. Obesity: global epidemiology and pathogenesis. *Nat Rev Endocrinol* 2019; 15: 288–298.
- Tutunchi H, Ostadrahimi A, Hosseinzadeh-Attar MJ, *et al.* A systematic review of the association of neuregulin 4, a brown fat-enriched secreted factor, with obesity and related metabolic disturbances. *Obes Rev* 2020; 21: e12952.
- Srivastava A, Srivastava N and Mittal B. Genetics of obesity. *Indian J Clin Biochem* 2016; 31: 361–371.
- Chung A, Backholer K, Wong E, *et al.* Trends in child and adolescent obesity prevalence in economically advanced countries according to socioeconomic position: a systematic review. *Obes Rev* 2016; 17: 276–295.
- Cobb LK, Appel LJ, Franco M, *et al.* The relationship of the local food environment with obesity: a systematic review of methods, study quality, and results. *Obesity* 2015; 23: 1331–1344.
- Tutunchi H, Jafarabadi MA, Hoojehani S, *et al.* General and abdominal obesity is related to socioeconomic status and food choices: a cross-sectional study. *Nutr Food Sci* 2019; 50: 61–73.
- Jiang SZ, Lu W, Zong XF, *et al.* Obesity and hypertension. *Exp Ther Med* 2016; 12: 2395–2399.
- Ortega FB, Lavie CJ and Blair SN. Obesity and cardiovascular disease. *Cir Res* 2016; 118: 1752–1770.
- Tutunchi H, Naeini F, Ebrahimi-Mameghani M, *et al.* The association of the steatosis severity, NAFLD fibrosis score and FIB-4 index with atherogenic dyslipidaemia in adult patients with NAFLD: a cross-sectional study. *Int J Clin Pract* 2021; 75: e14131.
- Tutunchi H, Saghafi-Asl M, Asghari-Jafarabadi M, *et al.* Association between dietary patterns and non-alcoholic fatty liver disease: results from a case-control study. *Arch Iran Med* 2021; 24: 35–42.
- Di Pasquale R and Celsi L. Stigmatization of overweight and obese peers among children. *Front Psychol* 2017; 8: 524.
- James PT, Leach R, Kalamara E, *et al.* The worldwide obesity epidemic. *Obes Res* 2001; 9(Suppl. 11): 228S–233S.
- Szewka AJ, Bruce BB, Newman NJ, *et al.* Idiopathic intracranial hypertension: relation between obesity and visual outcomes. *J Neuroophthalmol* 2013; 33: 4–8.
- Teberik K, Eski MT, Doğan S, *et al.* Ocular abnormalities in morbid obesity. *Arq Bras Oftalmol* 2019; 82: 6–11.
- Pasquale LR, Willett WC, Rosner BA, *et al.* Anthropometric measures and their relation to incident primary open-angle glaucoma. *Ophthalmology* 2010; 117: 1521–1529.
- Leske MC, Connell A, Wu S-Y, *et al.* Risk factors for open-angle glaucoma: the Barbados Eye Study. *Arch Ophthalmol* 1995; 113: 918–924.
- Mares JA, Volland R, Adler R, *et al.* Healthy diets and the subsequent prevalence of nuclear cataract in women. *Arch Ophthalmol* 2010; 128: 738–749.
- Ye J, Lou L-X, He J-J, *et al.* Body mass index and risk of age-related cataract: a meta-analysis of prospective cohort studies. *PLoS ONE* 2014; 9: e89923.
- Mares JA, Volland RP, Sondel SA, *et al.* Healthy lifestyles related to subsequent prevalence of age-related macular degeneration. *Arch Ophthalmol* 2011; 129: 470–480.
- Adams MK, Simpson JA, Aung KZ, *et al.* Abdominal obesity and age-related macular degeneration. *Am J Epidemiol* 2011; 173: 1246–1255.
- Klein R, Klein BE and Moss SE. Is obesity related to microvascular and macrovascular complications in diabetes? The Wisconsin Epidemiologic Study of Diabetic Retinopathy. *Arch Intern Med* 1997; 157: 650–656.
- Raman R, Rani PK, Gnanamoorthy P, *et al.* Association of obesity with diabetic retinopathy: Sankara Nethralaya Diabetic Retinopathy Epidemiology and Molecular Genetics Study (SN-DREAMS Report no. 8). *Acta Diabetol* 2010; 47: 209–215.
- Mantzoros CS. The role of leptin in human obesity and disease: a review of current evidence. *Ann Intern Med* 1999; 130: 671–680.

24. Kahn BB and Flier JS. Obesity and insulin resistance. *J Clin Invest* 2000; 106: 473–481.
25. Öner Rİ and Karadağ AS. Evaluation of choroidal perfusion changes in obese patients: ocular effects of insulin resistance. *Arq Bras Oftalmol* 2018; 81: 461–465.
26. Collins S. Overview of clinical perspectives and mechanisms of obesity. *Birth Defects Res A Clin Mol Teratol* 2005; 73: 470–471.
27. Cui JZ, Hornan D, Potter MJ, *et al.* The role of leptin in choroidal neovascularization. *Am J Ophthalmol* 2001; 132: 792–794.
28. Gariano RF, Nath AK, D'Amico DJ, *et al.* Elevation of vitreous leptin in diabetic retinopathy and retinal detachment. *Invest Ophthalmol Vis Sci* 2000; 41: 3576–3581.
29. Boillot A, Zoungas S, Mitchell P, *et al.* Obesity and the microvasculature: a systematic review and meta-analysis. *PLoS ONE* 2013; 8: e52708.
30. Czernichow S, Greenfield JR, Galan P, *et al.* Macrovascular and microvascular dysfunction in the metabolic syndrome. *Hypertens Res* 2010; 33: 293–297.
31. De Jongh RT, Serné EH, IJzerman RG, *et al.* Impaired microvascular function in obesity: implications for obesity-associated microangiopathy, hypertension, and insulin resistance. *Circulation* 2004; 109: 2529–2535.
32. Shankar A, Sabanayagam C, Klein BE, *et al.* Retinal microvascular changes and the risk of developing obesity: population-based cohort study. *Microcirculation* 2011; 18: 655–662.
33. Wong TY, Duncan BB, Golden SH, *et al.* Associations between the metabolic syndrome and retinal microvascular signs: the Atherosclerosis Risk In Communities study. *Invest Ophthalmol Vis Sci* 2004; 45: 2949–2954.
34. Nickla DL and Wallman J. The multifunctional choroid. *Prog Retin Eye Res* 2010; 29: 144–168.
35. Grundy SM. Metabolic complications of obesity. *Endocr J* 2000; 13: 155–165.
36. Lam BCC, Koh GCH, Chen C, *et al.* Comparison of body mass index (BMI), body adiposity index (BAI), waist circumference (WC), waist-to-hip ratio (WHR) and waist-to-height ratio (WHtR) as predictors of cardiovascular disease risk factors in an adult population in Singapore. *PLoS ONE* 2015; 10: e0122985.
37. Tutunchi H, Ebrahimi-Mameghani M, Ostadrahimi A, *et al.* What are the optimal cut-off points of anthropometric indices for prediction of overweight and obesity? Predictive validity of waist circumference, waist-to-hip and waist-to-height ratios. *Health Promot Perspect* 2020; 10: 142–147.
38. Cheong KC, Ghazali SM, Hock LK, *et al.* The discriminative ability of waist circumference, body mass index and waist-to-hip ratio in identifying metabolic syndrome: variations by age, sex and race. *Diabetes Metab Syndr* 2015; 9: 74–78.
39. Taylor B, Rochtchina E, Wang JJ, *et al.* Body mass index and its effects on retinal vessel diameter in 6-year-old children. *Int J Obes* 2007; 31: 1527–1533.
40. Wong TY, Shankar A, Klein R, *et al.* Prospective cohort study of retinal vessel diameters and risk of hypertension. *BMJ* 2004; 329: 79.
41. McGeechan K, Liew G, Macaskill P, *et al.* Prediction of incident stroke events based on retinal vessel caliber: a systematic review and individual-participant meta-analysis. *Am J Epidemiol* 2009; 170: 1323–1332.
42. Kotsis VT, Stabouli SV, Papamichael CM, *et al.* Impact of obesity in intima media thickness of carotid arteries. *Obesity* 2006; 14: 1708–1715.
43. Berni A, Giuliani A, Tartaglia F, *et al.* Effect of vascular risk factors on increase in carotid and femoral intima-media thickness. Identification of a risk scale. *Atherosclerosis* 2011; 216: 109–114.
44. Cheung CY-I, Ikram MK, Sabanayagam C, *et al.* Retinal microvasculature as a model to study the manifestations of hypertension. *Hypertension* 2012; 60: 1094–1103.
45. Cheung CY, Ikram MK, Klein R, *et al.* The clinical implications of recent studies on the structure and function of the retinal microvasculature in diabetes. *Diabetologia* 2015; 58: 871–885.
46. Manna P and Jain SK. Obesity, oxidative stress, adipose tissue dysfunction, and the associated health risks: causes and therapeutic strategies. *Metab Syndr Relat Disord* 2015; 13: 423–444.
47. McMurray F, Patten DA and Harper ME. Reactive oxygen species and oxidative stress in obesity – recent findings and empirical approaches. *Obesity* 2016; 24: 2301–2310.
48. Trayhurn P. Hypoxia and adipose tissue function and dysfunction in obesity. *Physiol Rev* 2013; 93: 1–21.
49. Murata T, Nagai R, Ishibashi T, *et al.* The relationship between accumulation of advanced glycation end products and expression of vascular endothelial growth factor in human diabetic retinas. *Diabetologia* 1997; 40: 764–769.

50. Orasanu G and Plutzky J. The pathologic continuum of diabetic vascular disease. *J Am Coll Cardiol* 2009; 53(Suppl. 5): S35–S42.
51. Kurtul B, Çakmak A, Elbeyli A, *et al.* Association of childhood obesity with retinal microvasculature and corneal endothelial cell morphology. *J Pediatr Endocrinol Metab* 2021; 34: 171–176.
52. İçel E, Uçak T, Yilmaz H, *et al.* Optical coherence tomography angiography findings in obese women. *Türk Klin Oftalmol Derg* 2020; 29: 218–223.
53. Wang JJ, Taylor B, Wong TY, *et al.* Retinal vessel diameters and obesity: a population-based study in older persons. *Obesity* 2006; 14: 206–214.
54. Erşan I, Battal F, Aylanç H, *et al.* Noninvasive assessment of the retina and the choroid using enhanced-depth imaging optical coherence tomography shows microvascular impairments in childhood obesity. *J AAPOS* 2016; 20: 58–62.
55. Ahmad M, Kaszubski PA, Cobbs L, *et al.* Choroidal thickness in patients with coronary artery disease. *PLoS ONE* 2017; 12: e0175691.
56. Kase S, Endo H, Yokoi M, *et al.* Choroidal thickness in diabetic retinopathy in relation to long-term systemic treatments for diabetes mellitus. *Eur J Ophthalmol* 2016; 26: 158–162.
57. Chen FK, Yeoh J, Rahman W, *et al.* Topographic variation and interocular symmetry of macular choroidal thickness using enhanced depth imaging optical coherence tomography. *Invest Ophthalmol Vis Sci* 2012; 53: 975–985.
58. Xiong S, He X, Deng J, *et al.* Choroidal thickness in 3001 Chinese children aged 6 to 19 years using swept-source OCT. *Sci Rep* 2017; 7: 45059.
59. Mori T, Sugano Y, Maruko I, *et al.* Subfoveal choroidal thickness and axial length in preschool children with hyperopic anisometropic amblyopia. *Curr Eye Res* 2015; 40: 954–961.
60. Lee GY, Yu S, Kang HG, *et al.* Choroidal thickness variation according to refractive error measured by spectral domain-optical coherence tomography in Korean children. *Korean J Ophthalmol* 2017; 31: 151–158.
61. Heirani M, Shandiz JH, Shojaei A, *et al.* Choroidal thickness profile in normal Iranian eyes with different refractive status by spectral-domain optical coherence tomography. *J Curr Ophthalmol* 2020; 32: 58–68.
62. Karapetyan A, Ouyang P, Tang LS, *et al.* Choroidal thickness in relation to ethnicity measured using enhanced depth imaging optical coherence tomography. *Retina* 2016; 36: 82–90.
63. Regatieri CV, Branchini L, Carmody J, *et al.* Choroidal thickness in patients with diabetic retinopathy analyzed by spectral-domain optical coherence tomography. *Retina* 2012; 32: 563–568.
64. Koizumi H, Yamagishi T, Yamazaki T, *et al.* Subfoveal choroidal thickness in typical age-related macular degeneration and polypoidal choroidal vasculopathy. *Graefes Arch Clin Exp Ophthalmol* 2011; 249: 1123–1128.
65. Okamoto M, Matsuura T and Ogata N. Choroidal thickness and choroidal blood flow after intravitreal bevacizumab injection in eyes with central serous chorioretinopathy. *Ophthalmic Surg Lasers Imaging Retina* 2015; 46: 25–32.
66. Dhoot DS, Huo S, Yuan A, *et al.* Evaluation of choroidal thickness in retinitis pigmentosa using enhanced depth imaging optical coherence tomography. *Br J Ophthalmol* 2013; 97: 66–69.
67. Akay F, Gundogan FC, Yolcu U, *et al.* Choroidal thickness in systemic arterial hypertension. *Eur J Ophthalmol* 2016; 26: 152–157.
68. Vance SK, Imamura Y and Freund KB. The effects of sildenafil citrate on choroidal thickness as determined by enhanced depth imaging optical coherence tomography. *Retina* 2011; 31: 332–335.
69. Kim DY, Silverman RH, Chan RVP, *et al.* Measurement of choroidal perfusion and thickness following systemic sildenafil (Viagra®). *Acta Ophthalmol* 2013; 91: 183–188.
70. Sogawa K, Nagaoka T, Takahashi A, *et al.* Relationship between choroidal thickness and choroidal circulation in healthy young subjects. *Am J Ophthalmol* 2012; 153: 1129–1132.
71. Tanaka K, Mori R, Kawamura A, *et al.* Comparison of OCT angiography and indocyanine green angiographic findings with subtypes of polypoidal choroidal vasculopathy. *Br J Ophthalmol* 2017; 101: 51–55.
72. Singh R, Invernizzi A, Agarwal A, *et al.* Enhanced depth imaging spectral domain optical coherence tomography versus ultrasonography B-scan for measuring retinochoroidal thickness in normal eyes. *Retina* 2015; 35: 250–256.
73. Yamashita T, Yamashita T, Shirasawa M, *et al.* Repeatability and reproducibility of subfoveal choroidal thickness in normal eyes of Japanese using different SD-OCT devices. *Invest Ophthalmol Vis Sci* 2012; 53: 1102–1107.
74. Matsuo Y, Sakamoto T, Yamashita T, *et al.* Comparisons of choroidal thickness of normal eyes obtained by two different spectral-domain OCT instruments and one swept-source OCT

- instrument. *Invest Ophthalmol Vis Sci* 2013; 54: 7630–7636.
75. Nagasawa T, Mitamura Y, Katome T, *et al.* Macular choroidal thickness and volume in healthy pediatric individuals measured by swept-source optical coherence tomography. *Invest Ophthalmol Vis Sci* 2013; 54: 7068–7074.
76. Yilmaz I, Ozkaya A, Kocamaz M, *et al.* Correlation of choroidal thickness and body mass index. *Retina* 2015; 35: 2085–2090.
77. Dogan B, Kazim Erol M, Dogan U, *et al.* The retinal nerve fiber layer, choroidal thickness, and central macular thickness in morbid obesity: an evaluation using spectral-domain optical coherence tomography. *Eur Rev Med Pharmacol Sci* 2016; 20: 886–891.
78. Özen B, Öztürk H, Çatlı G, *et al.* Does obesity affect the ocular choroid tissue in children and adolescents? *Pediatr Res* 2018; 5: 112.
79. Öncül H, Çağlayan M, Fuat Alakus M, *et al.* Evaluation of the subfoveal choroidal and outer retinal layer thickness in obese women. *Clin Exp Optom* 2021; 104: 178–186.
80. Yumusak E, Ornek K, Durmaz SA, *et al.* Choroidal thickness in obese women. *BMC Ophthalmol* 2016; 16: 48.
81. Bulus AD, Can ME, Baytaroglu A, *et al.* Choroidal thickness in childhood obesity. *Ophthalmic Surg Lasers Imaging Retina* 2017; 48: 10–17.
82. Panon N, Luangsawang K, Rugaber C, *et al.* Correlation between body mass index and ocular parameters. *Clin Ophthalmol* 2019; 13: 763–769.
83. Nilsson SF. Nitric oxide as a mediator of parasympathetic vasodilation in ocular and extraocular tissues in the rabbit. *Invest Ophthalmol Vis Sci* 1996; 37: 2110–2119.
84. Kotsis V, Stabouli S, Papakatsika S, *et al.* Mechanisms of obesity-induced hypertension. *Hypertens Res* 2010; 33: 386–393.
85. Xin C, Wang J, Zhang W, *et al.* Retinal and choroidal thickness evaluation by SD-OCT in adults with obstructive sleep apnea-hypopnea syndrome (OSAS). *Eye* 2014; 28: 415–421.
86. Bouloumie A, Marumo T, Lafontan M, *et al.* Leptin induces oxidative stress in human endothelial cells. *FASEB J* 1999; 13: 1231–1238.
87. Parhami F, Tintut Y, Ballard A, *et al.* Leptin enhances the calcification of vascular cells: artery wall as a target of leptin. *Circ Res* 2001; 88: 954–960.
88. Dogan B, Dogan U, Erol MK, *et al.* Optical coherence tomography parameters in morbidly obese patients who underwent laparoscopic sleeve gastrectomy. *J Ophthalmol* 2016; 2016: 5302368.
89. Gonul S, Yilmaz H, Gedik S, *et al.* Evaluation of the choroidal thickness and retinal nerve fiber layer and visual fields in morbid obesity: does bariatric surgery affect retinal structure and function? *Indian J Ophthalmol* 2021; 69: 301–306.
90. Terra X, Auguet T, Guiu-Jurado E, *et al.* Long-term changes in leptin, chemerin and ghrelin levels following different bariatric surgery procedures: Roux-en-Y gastric bypass and sleeve gastrectomy. *Obes Surg* 2013; 23: 1790–1798.
91. Bidaut-Garnier M, Schwartz C, Puyraveau M, *et al.* Choroidal thickness measurement in children using optical coherence tomography. *Retina* 2014; 34: 768–774.
92. Kim JH, Kim JS, Lee KW, *et al.* The posterior choroidal profiles measured by spectral domain optical coherence tomography in healthy Korean children. *J Korean Ophthalmol Soc* 2013; 54: 1708–1714.