


## ARTICLE

# *Mycobacterium leprae* induces a tolerogenic profile in monocyte-derived dendritic cells via TLR2 induction of IDO

Jéssica A. P. Oliveira<sup>1</sup> | Mariana Gandini<sup>2</sup> | Jorgenilce S. Sales<sup>1</sup> | Sérgio K. Fujimori<sup>3</sup> |  
Mayara G. M. Barbosa<sup>4</sup> | Valber S. Frutuoso<sup>5</sup> | Milton O. Moraes<sup>1</sup> | Euzenir N. Sarno<sup>1</sup> |  
Maria C. V. Pessolani<sup>2</sup> | Roberta O. Pinheiro<sup>1</sup> 

<sup>1</sup>Leprosy Laboratory, Oswaldo Cruz Institute, Oswaldo Cruz Foundation, Rio de Janeiro, Brazil

<sup>2</sup>Laboratory of Cellular Microbiology, Oswaldo Cruz Institute, Oswaldo Cruz Foundation, Rio de Janeiro, Brazil

<sup>3</sup>Laboratory for Development and Analytical Validation, Oswaldo Cruz Foundation, Farmanguinhos, Rio de Janeiro, Brazil

<sup>4</sup>Cascalho-Platt Laboratory, Department of Surgery, University of Michigan, Ann Arbor, Michigan, USA

<sup>5</sup>Immunopharmacology Laboratory, Oswaldo Cruz Institute/Oswaldo Cruz Foundation, Rio de Janeiro, Brazil

## Correspondence

Roberta Olmo Pinheiro, Researcher in Public Health, Leprosy Laboratory, Oswaldo Cruz Foundation (Fiocruz), Rio de Janeiro, Brazil.  
Email: robertaolmo@gmail.com;  
rolmo@ioc.fiocruz.br

## Abstract

The enzyme IDO-1 is involved in the first stage of tryptophan catabolism and has been described in both microbicidal and tolerogenic microenvironments. Previous data from our group have shown that IDO-1 is differentially regulated in the distinctive clinical forms of leprosy. The present study aims to investigate the mechanisms associated with IDO-1 expression and activity in human monocyte-derived dendritic cells (mDCs) after stimulation with irradiated *Mycobacterium leprae* and its fractions. *M. leprae* and its fractions induced the expression and activity of IDO-1 in human mDCs. Among the stimuli studied, irradiated *M. leprae* and its membrane fraction (MLMA) induced the production of proinflammatory cytokines TNF and IL-6 whereas irradiated *M. leprae* and its cytosol fraction (MLSA) induced an increase in IL-10. We investigated if TLR2 activation was necessary for IDO-1 induction in mDCs. We observed that in cultures treated with a neutralizing anti-TLR2 antibody, there was a decrease in IDO-1 activity and expression induced by *M. leprae* and MLMA. The same effect was observed when we used a MyD88 inhibitor. Our data demonstrate that coculture of mDCs with autologous lymphocytes induced an increase in regulatory T (Treg) cell frequency in MLSA-stimulated cultures, showing that *M. leprae* constituents may play opposite roles that may possibly be related to the dubious effect of IDO-1 in the different clinical forms of disease. Our data show that *M. leprae* and its fractions are able to differentially modulate the activity and functionality of IDO-1 in mDCs by a pathway that involves TLR2, suggesting that this enzyme may play an important role in leprosy immunopathogenesis.

## KEYWORDS

dendritic cells, IDO, lepromatous leprosy, leprosy, *Mycobacterium leprae*, reversal reaction

## 1 | INTRODUCTION

Leprosy is a chronic infectious disease, which mainly affects the peripheral nerves and skin. The disease is caused by *Mycobacterium leprae*, an obligate intracellular parasite that predominantly infects macrophages, endothelial cells, and Schwann cells, being

the unique specie of mycobacteria capable of infecting peripheral nerves.<sup>1</sup>

Previous studies have demonstrated that *M. leprae* induces an increase in the gene and protein expression of the enzyme IDO-1 in human monocytes.<sup>2</sup> IDO-1 is an intracellular enzyme that catalyzes the early stage of tryptophan (trp) catabolism along the kynurenine (kyn)

Abbreviations: CMV, cytomegalovirus; HCV, Hepatitis C virus; HTLV, Human T cell lymphotropic virus; IDO, Indoleamine 2,3-dioxygenase; kyn, kynurenine; 1-MT, 1-methyl tryptophan; MLMA, *M. leprae* membrane fraction; MLSA, *M. leprae* cytosol fraction; mDCs, monocyte-derived dendritic cells; PBMCs, Peripheral blood mononuclear cells; TLR, Toll-like receptor; trp, tryptophan; Tregs, regulatory T cells; VDRL, Venereal Disease Research Laboratory test.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2020 The Authors. *Journal of Leukocyte Biology* published by Wiley Periodicals LLC on behalf of Society for Leukocyte Biology

pathway.<sup>3</sup> Several cell types such as macrophages, epithelial cells, and dendritic cells (DCs) express IDO-1 that can be induced by proinflammatory cytokines, such as IFN- $\gamma$ , TLR ligands, such as LPS, and interactions between immune cells through costimulatory molecules such as CD80 and CD86.<sup>3–5</sup> It is known that IDO-1 can affect immunity through two nonexclusive mechanisms: the establishment of a local response with “amino acid deprivation” that inhibits pathogen growth and the production of trp metabolites with immunomodulatory functions or cytotoxic agents that inhibit T-cell activation and modulate the differentiation of naïve T cells into regulatory T cells (Tregs).<sup>6,7</sup>

Our group has previously demonstrated a significant increase of IDO-1 in cells present in skin lesions of patients with multibacillary leprosy (lepromatous leprosy) compared to patients with the paucibacillary form (tuberculoid leprosy).<sup>8–10</sup> Lipoproteins (19 and 33 kDa) present in *M. leprae* plasma membrane are well known to activate monocytes and DCs through TLR2.<sup>11</sup> Analyses of skin lesions from leprosy patients show that TLR2 is strongly expressed in cells of paucibacillary patients, in contrast to poor expression in cells from multibacillary patient lesions.<sup>11</sup> A subsequent study showed that activation of TLR2/1 leads to rapid differentiation of human peripheral monocytes in CD1b<sup>+</sup> DCs in paucibacillary patients and in DC-SIGN<sup>+</sup> cells in multibacillary patients,<sup>12</sup> suggesting that TLR-induced monocyte differentiation in macrophages or DCs influences the host response to *M. leprae* infection. Here, we investigated the ability of *M. leprae* and subcellular fractions to modulate IDO-1 expression and activity as well as their capacity to induce a tolerogenic or microbicidal phenotype in human monocyte-derived dendritic cells (mDCs).

## 2 | MATERIALS AND METHODS

### 2.1 | Obtaining buffy coats

Buffy coats were obtained from healthy blood donors in the hemotherapy service of Clementino Fraga Filho University Hospital of the Federal University of Rio de Janeiro (UFRJ) through a technical-scientific partnership approved by the Research Ethics Committee of the Oswaldo Cruz Foundation (approval number: 1.538.467). Inclusion and exclusion criteria were the same as those used for screening in blood banks, and volunteers under 18 yr of age whose serologic screening was positive for hepatitis B (HbsAg and anti-HBc), hepatitis C (HCV), AIDS (HIV-1/2 Ag + Ab combined test), Chagas disease (anti-*Trypanosoma cruzi*), syphilis (Venereal Disease Research Laboratory)—nontreponemal, (Human T-cell lymphotropic virus)-I and HTLV-II, malaria, and CMV were excluded.

### 2.2 | Obtaining human dendritic cells differentiated from monocytes

Human PBMCs were obtained under endotoxin-free conditions by the Ficoll-Paque PLUS method (GE, Chicago, Illinois, USA). PBMCs were labeled with CD14 magnetic beads (Miltenyi Biotec, Bergisch Gladbach, Germany) microbeads for 15 min at 4°C and passed through a

positive separation column. After monocyte separation, CD14<sup>+</sup> cells were cultured in the presence of medium containing IL-4 (10 ng/ml) and GM-CSF (50 ng/ml) (PeproTech, Cranbury, NJ, USA) for 6 d at 37°C with 5% CO<sub>2</sub>. After this time the nonadherent cells (mDCs) were then counted and plated.

### 2.3 | Cell culture stimuli and infection

mDCs were stimulated with 0.1, 1, or 10  $\mu$ g/ml of irradiated *M. leprae*, and its membrane (MLMA) and cytosol fractions (MLSA) (BEI Resources [NIH/ATCC], Manassas, VA, USA, 200  $\mu$ M of the IDO inhibitor 1-methyl-D-tryptophan (1-MT, SigmaAldrich, Saint Louis, MO, USA), 500  $\mu$ g/ml of the anti-human TNF (PeproTech), 100  $\mu$ g/ml of the anti-human IL-10 (PeproTech), 10  $\mu$ g/ml of the TLR2 agonist Pam3Cys-Ser-(Lys)4 (SigmaAldrich), 100  $\mu$ M of the inhibitory peptide for MyD88 homodimerization (IMGEX, San Diego, CA, USA), or 1  $\mu$ g/ml of Mab-mTLR2 (InvivoGen, San Diego, CA, USA), and incubated for 24 h at 37°C in 5% CO<sub>2</sub>. *M. leprae* and its fractions were tested for purity and the absence of endotoxin. According to the limulus amebocyte lysate assay (Lonza, Basel, Switzerland), all stimuli used for in vitro cultures were shown to contain less than 0.1 U/ml endotoxin.

### 2.4 | Flow cytometry

Panels of antibodies used for phenotypic detection and intracellular cytokine detection are described in Table 1. Following stimulation,  $1 \times 10^6$  mDCs were transferred from the plate to cytometry minitubes. Cells were washed and then fixed (2% paraformaldehyde). Subsequently, mDCs were permeabilized (0.15% saponin in PBS) and incubated for 30 min at 4°C with their respective antibodies. At the end of the incubation, cells were washed, suspended, and cell phenotype was evaluated by flow cytometry (FACS Aria IIu, BD Biosciences, Franklin Lakes, NJ, USA). For each sample, a minimum of 10,000 events were acquired. The analysis was performed using the FlowJo software.

### 2.5 | ELISA

Supernatants were tested for the presence of TNF, IL-6, IL-10, IL-12, and IL-15 cytokines using commercially available ELISA (eBioscience, San Diego, CA, USA) following the protocols supplied by the manufacturer.

### 2.6 | HPLC

Determination of IDO-1 activity in mDC culture supernatants was performed by determination of kyn and trp levels<sup>13</sup> by HPLC. In an Eppendorf tube, 165  $\mu$ l of culture supernatant were homogenized with 5  $\mu$ l tyrosine and 25  $\mu$ l Trichloroacetic acid (TCA). The tubes were then centrifuged for 10 min at 15,800  $\times$ g. After centrifugation, the supernatant was collected and 50  $\mu$ l were injected into the reverse phase C18 column. Run detection was performed at 365 nm to detect kyn and 285 nm for trp and the internal normalizer tyrosine. IDO activity was assessed by the ratio of kyn to trp.

**TABLE 1** Antibodies used in flow cytometry

Antibody	Clone	Catalogue number	Company	Fluorochrome
CD14	61D3	120149-42	eBioscience	PE
HLA-DR	G46-6	555811	BD Biosciences	FITC
CD11c	3.9	301606	BioLegend	PE
CD209	eB-h209	11-209973	eBioscience	FITC
CD86	IT2.2	15-0869-71	eBioscience	PE-Cy5
CD83	HB1Se	305320	BioLegend	PerCP-Cy5.5
CD304	AD5-17F6	130-090-900	Miltenyi Biotec	APC
CD123	6H6	306012	BioLegend	APC
IDO	700838	IC6030P	R&D Systems (Minneapolis, MN, USA)	PE
TNF- $\alpha$	MAB11	561023	BD eBioscience	AlexaFluor700
IL-10	127107	IC2172A	R&D	APC

## 2.7 | Coculture of mDCs and lymphocytes

To assess the functional role of IDO-1, mDCs were obtained as described earlier and plated in 96-well plates ( $2 \times 10^4$ ). After the stimulus period, autologous lymphocytes were added at a ratio of 1 (mDC):20 (lymphocytes) for 5 d at 37°C 5% CO<sub>2</sub>. Subsequently, cells were harvested, washed, and labeled with anti-FoxP3 Alexa Fluor 488/CD4 PE-Cy5/CD25 PE antibodies according to the specifications of the True-Nuclear Human Treg Flow Kit (BioLegend, San Diego, CA, USA). Cells were analyzed by flow cytometry (FACS Aria IIu).

## 2.8 | Statistical analysis

Analyses of the experiments were performed by Kruskal-Wallis test or 1-way ANOVA. For all statistical analyses the value of  $P \leq 0.05$  was considered significant. Statistical analyses were performed using the Windows GraphPad Prism version 8.0 software (GraphPad Software, San Diego, CA, USA).

# 3 | RESULTS

## 3.1 | *M. leprae* increases IDO-1 expression and activity in mDCs


Previous data from our group have demonstrated that *M. leprae* induces the expression and activity of IDO-1 in human monocytes.<sup>8</sup> In order to investigate whether *M. leprae* and its fractions are capable of modulating IDO-1 protein expression in mDCs, cells were stimulated with *M. leprae*, MLMA, or MLSA for 24 h. DCs were analyzed by flow cytometry to evaluate IDO-1 expression (Fig. 1A) and a panel of specific markers was used to confirm their differentiation (Supporting Information Fig. S1A–R). *M. leprae* and MLMA fraction were efficient in inducing IDO-1 expression at 10  $\mu$ g/ml, but not MLSA (Fig. 1B, C). The kyn/try ratio in the supernatants reflects IDO-1 activity. To confirm if the enzymatic activity of IDO-1 was also modulated by different mycobacterial stimuli, the kyn/try ratio in the supernatants was ana-

lyzed by HPLC. As observed in Fig. 1D, *M. leprae* ( $0.3 \pm 0.02$  in ML vs.  $0.05 \pm 0.01$  in NS,  $P = 0.01$ ) and MLMA ( $0.24 \pm 0.01$  in MLMA vs.  $0.05 \pm 0.01$  in NS,  $P = 0.02$ ) were able to significantly increase the enzymatic activity of IDO-1, but the same was not observed with the MLSA fraction. These data show that antigens present in membrane of the bacilli are able to induce IDO-1 activity.

## 3.2 | TNF is important in the induction of IDO-1 activity by *M. leprae* and its fractions in mDCs

Analysis of TNF<sup>+</sup> cells by flow cytometry revealed that the MLMA fraction was able to induce an increase in the frequency of double positive IDO-1<sup>+</sup> TNF<sup>+</sup> cells when compared with nonstimulated cells ( $8.35 \pm 2.17$  in MLMA vs.  $0.92 \pm 0.24$  in NS,  $P = 0.022$ ) (Fig. 2A–E). However, neither *M. leprae* nor its fractions were able to induce an increase in the frequency of IDO-1<sup>+</sup> IL-10<sup>+</sup> cells (Supporting Information Fig. S2). Cytokine concentrations in the culture supernatants of mDCs stimulated with *M. leprae*, MLMA and MLMA for 24 h were evaluated by ELISA. As observed in Figure 3, *M. leprae* was not able to significantly modulate the production of the cytokines tested. However, MLMA was able to induce a significant increase in TNF, IL-6, and IL-12 concentrations (Fig. 3A, C, ) when compared to nonstimulated cells. In contrast, MLSA induced an increase in IL-10 levels (Fig. 3B). There was no change in IL-15 production by mDCs after the stimuli with *M. leprae* and its fractions (not shown).

TNF, which is normally present in infectious environments, synergistically enhances IFN- $\gamma$ -induced IDO-1 activity.<sup>14</sup> Because the stimulation with MLMA induced an increase in TNF and stimulation with MLSA-induced IL-10, we investigated the role of both IL-10 and TNF in the induction of IDO-1 expression and activity by *M. leprae* and its fractions on mDCs. As shown in Figure 4, the blockade of TNF in cultures stimulated with *M. leprae* or its fractions led to a significant decrease in IDO-1 activity in mDCs cultures, suggesting an important role of this cytokine in the induction of IDO-1 by *M. leprae* and its fractions. The same effect was not observed after IL-10 blockade (Fig. 4A–C).



**FIGURE 1** IDO-1 activity and expression in human monocyte-derived dendritic cells (mDCs). The differentiated mDCs were plated ( $1 \times 10^6$  cells per well). (A) Gate strategy for flow cytometry analysis. (B) mDCs were stimulated with *M. leprae* (ML), membrane fraction of *M. leprae* (MLMA), or soluble fraction of *M. leprae* (MLSA) at 0.1, 1, 10, or 50  $\mu\text{g/mL}$  for 24 h. After this time, cells were labeled with anti-IDO-1 intracellular antibody. Acquisition was performed in a BD FACS Aria IIu flow cytometer and analysis performed using FlowJo software. NS—unstimulated cell. The graph shows the percentage of IDO-1-positive mDC of three independent experiments performed in triplicate. (C) Representative histograms of mDCs stimulated with ML or its fractions. (D) Supernatants from cell cultures were collected and IDO-1 activity was assessed by the kynurenine/tryptophan ratio (kyn/trp) by HPLC. The plot represents the mean  $\pm$  SD of three independently performed experiments. \* $P < 0.05$ ; \*\* $P < 0.01$ ; and \*\*\* $P < 0.001$

### 3.3 | TLR2 is important for the induction of IDO-1 activity in mDCs


*M. leprae* membrane contains lipoproteins that can activate TLR2 signaling triggering an inflammatory response<sup>15,16</sup> and therefore a role for this signaling pathway on IDO-1 induction in stimulated mDCs was investigated. As observed in Figure 5A, the blockade of TLR2 decreased IDO-1 activity induced by *M. leprae* and its fractions, demonstrating that IDO-1 activity and expression in *M. leprae*-stimulated mDCs is TLR2 dependent. The TLR2 agonist Pam3Cys was used as positive control, suggesting that TLR2 pathway is involved in IDO-1 induction in mDCs ( $0.25 \pm 0.01$  in ML vs.  $0.03 \pm 0.01$  in ML + TLR2,  $P = 0.20$ ;  $0.20 \pm 0.02$  in MLMA vs.  $0.01 \pm 0.002$  in MLMA + TLR2,  $P = 0.01$ ;  $0.13 \pm 0.03$  in MLSA vs.  $0.02 \pm 0.006$  in MLSA + TLR2,  $P = 0.01$ ). In addition, we used a peptide inhibitor of the MyD88 pathway in parallel to a peptide control of the same pathway. A decrease in IDO-1 activity was observed when the inhibitory peptide was added to the cultures

( $0.22 \pm 0.03$  in ML vs.  $0.02 \pm 0.005$  in ML +  $\alpha\text{MyD88}$ ,  $P = 0.01$ ;  $0.18 \pm 0.02$  in MLMA vs.  $0.05 \pm 0.005$  in MLMA +  $\alpha\text{MyD88}$ ,  $P = 0.007$ ;  $0.08 \pm 0.02$  in MLSA vs.  $0.01 \pm 0.007$  in MLSA +  $\alpha\text{MyD88}$ ,  $P = 0.03$ ) (Fig. 5B).

### 3.4 | *M. leprae* and its fractions influence the tolerogenic profile of lymphocytes

In order to verify whether the increase in IDO-1 expression and activity influenced the lymphocyte phenotype after priming by the mDCs, a coculture assay was performed in which the mDCs stimulated with *M. leprae* and its fractions were cultivated with autologous lymphocytes in a proportion of 20 lymphocytes:1 mDC.

Through flow cytometric analysis (Supporting Information Fig. S3; Fig. 6A–H), an overall increase in Treg frequencies in MLSA or ML-stimulated mDCs cocultures was observed. Inhibition of IDO-1



**FIGURE 2** MLMA is able to increase the frequency of IDO-1<sup>+</sup>TNF<sup>+</sup> human monocyte-derived dendritic cells (mDCs). DCs differentiated from monocytes were plated ( $1 \times 10^6$  cells per well) and stimulated with *M. leprae* (ML), membrane fraction of *M. leprae* (MLMA), and soluble fraction of *M. leprae* (MLSA) at  $10 \mu\text{g/ml}$ , for 24 h, or not (NS—unstimulated cell). Cells were labeled with anti-IDO-1 PE and Alexa Fluor 700 TNF intracellular antibodies. Acquisition was performed in a BD FACS Aria IIu flow cytometer and analysis performed using FlowJo software. Analysis is representative of cells from a donor from four experiments. For each sample, a minimum of 10,000 events were analyzed. \* $P < 0.05$  (Mann-Whitney)

activity significantly abrogated Treg differentiation for those stimuli, suggesting that acquisition of the regulatory profile during cocultures may be IDO-1 dependent ( $9.19 \pm 2.20$  in ML vs.  $0.24 \pm 0.10$  in ML + 1MT,  $P = 0.007$  and  $7.32 \pm 0.73$  in MLSA vs.  $0.24 \pm 0.10$  in MLSA + 1MT,  $P = 0.007$ ) (Fig. 6I).

## 4 | DISCUSSION


The characterization of infectious environments is important to define how innate immunity influences the subsequent development of adaptive immunity. Regulatory mechanisms that operate at both stages of immunity are also critical in determining how the immune response will dictate the severity of an infectious process. In this context, we have explored the regulatory mechanisms exerted by the enzyme IDO-1 during *M. leprae* infection. At the lepromatous leprosy pole, high IDO-1 expression along with high levels of IL-10 and TGF- $\beta$  could be responsible for the pronounced reduction of the antigen-specific cellular immune response observed in these patients.<sup>2,8</sup> At the tuberculoid pole of the disease, where the expression of IFN- $\gamma$  predominates, IDO-1 would be induced mainly by this cytokine<sup>10</sup>

and could be associated with microbicidal activity, suggesting that in leprosy IDO-1 may have a dual role, being tolerogenic or microbicidal depending on the environment and the cytokines involved in its induction.


In skin lesions of leprosy patients, in addition to intact bacilli, some mycobacterial fractions from the membrane and the *M. leprae* cytosol exposed after bacterial death, which may be a result of both the host immune response and the treatment, should induce immune responses.<sup>1</sup> Thus, in addition to the use of irradiated *M. leprae*, mDCs were stimulated by the MLMA and MLSA, mimicking the tissue microenvironment.

We are the first to demonstrate that *M. leprae* is able to induce IDO-1 activity in mDCs when compared to unstimulated cells. In parallel, the intracellular expression of IDO-1 was analyzed by flow cytometry and we observed that *M. leprae* and its fractions were able to increase IDO-1 expression, with a higher increase after MLMA stimulation, but not MLSA. It has already been shown that *M. tuberculosis* infection leads to up-regulation of IDO-1 expression in murine macrophages.<sup>17</sup> In addition, elevated IDO-1 activity was described in many chronic inflammatory syndromes, including cancer, infections, allergies, and pregnancy.<sup>18</sup>






**FIGURE 3** Differential cytokine production in cell supernatants from MLMA or MLSA-stimulated human monocyte-derived dendritic cells (mDCs). mDCs were plated ( $1 \times 10^6$  cells per well) and stimulated with *M. leprae* (ML), membrane fraction of *M. leprae* (MLMA), and soluble fraction of *M. leprae* (MLSA) at  $10 \mu\text{g/ml}$ , for 24 h, or not (NS—unstimulated cell). (A–D) Supernatants from these cultures were then collected and TNF, IL-6, IL-10, and IL-12 concentrations were evaluated by ELISA. Graphs represent the mean  $\pm$  SD of four independently performed experiments. \* $P < 0.05$  (1-way ANOVA)



**FIGURE 4** Importance of TNF in the induction of IDO-1 activity in *M. leprae*-stimulated human monocyte-derived dendritic cells (mDCs). mDCs were plated ( $1 \times 10^6$  cells per well) and 1 h prior to challenge, anti-human IL-10 ( $100 \mu\text{g/ml}$ ) or anti-human TNF ( $500 \mu\text{g/ml}$ ) neutralizing antibodies were added at cultures followed by stimuli: *M. leprae* (ML), membrane fraction of *M. leprae* (MLMA), or soluble fraction of *M. leprae* (MLSA) were added  $10 \mu\text{g/ml}$  or not (NS—nonstimulated cell). After 24 h, supernatants were collected and IDO-1 activity was assessed by the kynurenine/tryptophan ratio (kyn/tryp) by HPLC. The plot represents the mean  $\pm$  SD of four independently performed experiments. \* $P < 0.05$  (Kruskal-Wallis)



**FIGURE 5** TLR2 is important in inducing IDO-1 activity in human monocyte-derived dendritic cells (mDCs). mDCs were plated ( $1 \times 10^6$  cells per well) and 1 h prior to challenge, anti-human TLR2 (1  $\mu$ g/ml) neutralizing antibody or MyD88 peptide inhibitor (MyD88i; IMGEX) (100  $\mu$ M) was added, followed by stimuli: *M. leprae* (ML), membrane fraction (MLMA), or soluble fraction (MLSA) was added at 10  $\mu$ g/ml, TLR2 agonist Pam3Cys was used as positive control (10  $\mu$ g/ml) for 24 h at 37°C 5%CO<sub>2</sub>. Controls are nonstimulated cultures. Supernatants from these cultures were collected and IDO-1 activity was assessed by the kynurenine/tryptophan ratio (kyn/trp) by HPLC. The plot represents the mean  $\pm$  SD of five independently performed experiments. \* $P < 0.05$  (Mann-Whitney)

The production of cytokines in these environments was evaluated. It should be noted that among the stimuli studied, MLMA significantly induced an increase in the production of inflammatory cytokines, such as TNF, IL-6, and IL-12. It was also observed that the MLMA fraction induced an increase in the frequency of IDO<sup>+</sup>TNF<sup>+</sup> cells. TNF has been suggested to be one of the most important cytokines for the induction of IDO-1. The contribution of TNF to the activation of IDO-1 induced by LPS has already been observed.<sup>19</sup> In addition, TNF mediates stress-induced depression by positively regulating the IDO-1 enzyme.<sup>20</sup> As TNF modulates the response to cytotoxicity and the production of inflammatory cytokines,<sup>21</sup> our data suggest that the MLMA fraction may be involved in a microbicidal effect.


Our results showed that *M. leprae* and MLSA led to an increase in IL-10 secretion, which could suggest a regulatory effect, because an earlier study<sup>2</sup> demonstrated that *M. leprae* is able to induce the gene and protein expression of IDO-1 in human monocytes by an IL-10-dependent mechanism. In leprosy, IL-15 induces the vitamin D-dependent antimicrobial pathway and this ability is consistent with the expression of this cytokine in self-limited tuberculoid lesions.<sup>22</sup> Here, we investigated the secretion of this cytokine in cell supernatants and observed that there was no change between the stimuli used (not shown).

Our data also show that blocking TNF decreases IDO-1 activity in mDCs by *M. leprae* and its fractions, unlike IL-10 blockade. These results led us to hypothesize that TNF is induced at different levels by different antigens present in cytosol and on bacilli cell membrane and that this cytokine is important for IDO-1 expression and activity in human mDCs. In contrast, IL-10 is not as efficient in inducing IDO-1 expression and activity in mDCs as observed previously in human monocytes stimulated with *M. leprae*.<sup>2</sup>

Recognition of components of the mycobacterial cell wall, such as sugars, lipids, and peptides can induce macrophages to secrete TNF and other cytokines like IL-12 that are essential for the development of an inflammatory response.<sup>23</sup> TLR2, activated by lipoarabinomannan, is required for the induction of TNF in macrophages by *M. tuberculosis*. Previous studies have further shown that the expression of IDO-1 in DCs can be induced by inflammatory cytokines and TLR ligands.<sup>6</sup>

Lipoproteins from *M. tuberculosis*, such as 19 kD and mannose-capped lipoarabinomannan, are TLR2 ligands that trigger the development of regulatory immune responses.<sup>24–26</sup> Previous studies have shown that lysophosphatidylcholine subverts TLR-mediated signaling in DCs, which drives cell differentiation to a tolerogenic phenotype.<sup>27</sup> In addition, it was demonstrated that tolerogenic effects induced by IDO-1 depend on TLR2/6-induced JNK signaling by *Faecalibacterium prausnitzii* and ectonucleotidase activity.<sup>28</sup> Our data demonstrated that *M. leprae* induces an increase in IDO-1 expression and activity in mDCs, and that the membrane components of the bacilli play an important role in this context via TLR2 signaling associated with the production of inflammatory cytokines. We cannot exclude the involvement of TLR4 on IDO-1 activation induced by *M. leprae*. Although *M. leprae* is an acid-fast intracellular gram-positive bacillus, there are evidences that it can activate TLR4.<sup>29</sup> Mycobacterial glycolipids such as the lipooligosaccharides contain the lipid A portion, which is responsible for the LPS interaction with TLR4.

IDO-1 may require TLR-MyD88-NF- $\kappa$ B signaling to promote the development of colitis.<sup>30</sup> Because NF- $\kappa$ B activation depends on MyD88 we performed assays using a MyD88 inhibitor, and we observed that the absence of MyD88 in cultures of mDCs leads to a decrease in the activity of IDO-1 also suggesting it might be the case.



**FIGURE 6** Increased regulatory T (Treg) cell expression by *M. leprae* and MLSA. Human monocyte-derived dendritic cells (mDCs) were plated ( $1 \times 10^4$  cells per well) and stimulated with *M. leprae* (ML), membrane fraction of *M. leprae* (MLMA) and soluble fraction of *M. leprae* (MLSA) at  $10 \mu\text{g/ml}$ , for 24 h, or 1MT (IDO inhibitor—1 h before other stimuli) or not (NS—unstimulated cell). Cells were washed and the autologous lymphocytes (20 lymphocytes:1 mDC) were cultured for 5 d. Cells were then labeled with intracellular anti-CD4, CD25, and FoxP3 or isotype antibodies. Acquisition was performed by BD FACS Aria IIu flow cytometry and analysis performed using FlowJo software. Analysis is representative of cells from a donor from four experiments. For each sample, a minimum of 10,000 events were analyzed. \* $P < 0.05$  (Mann-Whitney)

A key mechanism involved in mDCs-mediated immunosuppression is the expression of IDO-1.<sup>31</sup> The interaction of DCs with TLR2 and TLR4 treated with ES L1 (excretory-secretory muscle larvae) promotes the expansion of CD4<sup>+</sup> CD25<sup>high</sup>Foxp3<sup>high</sup> producing IL-10 and TGF- $\beta$  cells in an IDO-1-dependent form.<sup>32</sup> The 6-formylindolo[3,2-b]carbazole-treated DCs are able to induce differentiation of naive T lymphocytes into regulatory-like T cells.<sup>33</sup> Our coculture data showed that *M. leprae* is capable of enhancing the expression of Treg CD4<sup>+</sup> CD25<sup>+</sup>FOXP3<sup>+</sup> lymphocytes by an IDO-1-dependent form. MLSA, which increased IL-10 secretion, is capable of enhancing the expression of Treg by an IDO-1-independent form. It has also been shown that lepromatous patients increase the expression of Treg

cells, one factor that may explain the large number of bacilli in these individuals.<sup>34</sup> These findings provide a new insight into the mechanisms of TLR-induced tolerogenic phenotype in DCs, which may help to better understand the processes involved in the induction and resolution of chronic inflammation and tolerance.

Together, *M. leprae*-induced IDO-1 axis may play an important role in controlling the immunity and severity of leprosy. In clinical terms, our hypothesis is that in multibacillary patients, the high flow of living and dead bacilli induces IDO-1 by an IL-10-dependent mechanism that in turn leads to a tolerogenic effect, decreasing the cellular immune response. In paucibacillary patients and in the reaction episodes, the exposure of bacilli membrane antigens favors a context similar to that



described by mDCs stimulation with MLMA, that is, the increase of IDO-1 due to the activation of TNF and IL-12, which consequently leads to an increase in the pro-inflammatory response and could contribute to a decrease in bacillary load.

## AUTHORS

ROP, MOM, VSF, MCVP and ENS contributed to the design and implementation of the research. JAPO, MG and ROP to the writing of the manuscript. JAPO, MG, JSS, SKF, MGMB and ROP processed the experimental data and performed the analysis. All authors provided critical feedback and helped shape the research, analysis and manuscript. All authors contributed to the article and approved the submitted version.

## ACKNOWLEDGMENTS

We thank the Hemotherapy service of Clementino Fraga Filho University Hospital of the Federal University of Rio de Janeiro (HUCFF/UFRJ/Brazil) for the buffy coats. The following reagents were kindly obtained from the NIH Biodefense and Emerging Infections Research Resources Repository, NIAID, NIH: *M. leprae*, MLSA, and MLMA. We also thank Dr. John Spencer and Dr. Márcia Berredo for the MLMA antigen; Dr. Elzinandes Leal for the MyD88 inhibitor peptide; the Laboratory for Development and Analytical Validation (LDVA) for their support in HPLC; and the flow cytometry platform of Oswaldo Cruz Institute (IOC/FIOCRUZ) for acquisition and analysis of samples. Finally we thank CAPES, FAPERJ, and CNPq funding institutions for all their financial support.

## ORCID

Roberta O. Pinheiro  <https://orcid.org/0000-0001-8471-4227>

## REFERENCES

- Scollard DM, Adams LB, Gillis TP, Krahenbuhl JL, Truman RW, Williams DL. The continuing challenges of leprosy. *Clin Microbiol Rev*. 2006;19:338-381.
- Moura DF, de Mattos KA, Amadeu TP, et al. CD163 favors *Mycobacterium leprae* survival and persistence by promoting anti-inflammatory pathways in lepromatous macrophages. *Eur J Immunol*. 2012;42:2925-2936.
- Mellor AL, Munn DH. IDO expression by dendritic cells: tolerance and tryptophan catabolism. *Nat Rev Immunol*. 2004;4:762-774.
- Mbongue JC, Nicholas DA, Torrez TW, Kim NS, Firek AF, Langridge WH. The role of Indoleamine 2, 3-dioxygenase in immune suppression and autoimmunity. *Vaccine*. 2015;3:703-729.
- Heitger A. Regulation of expression and function of IDO in human dendritic cells. *Curr Med Chem*. 2011;18:2222-2233.
- Harden JL, Elgimez NK. Indoleamine 2,3 dioxygenase and dendritic cell tolerogenicity. *Immunol Invest*. 2012;41:738-764.
- Schmidt SV, Schultze JL. New insights into IDO biology in bacterial and viral infections. *Front Immunol*. 2014;5:1-12.
- De Souza Sales J, Lara FA, Amadeu TP, et al. The role of Indoleamine 2, 3-dioxygenase in lepromatous leprosy immunosuppression. *Clin Exp Immunol*. 2011;165:251-263.
- De Mattos Barbosa MG, da Silva Prata RB, Andrade PR, et al. Indoleamine 2,3-dioxygenase and iron are required for *Mycobacterium leprae* survival. *Microbe Infect*. 2017;19:505-514.
- Andrade PR, Pinheiro RO, Sales AM, et al. Type 1 reaction in leprosy: a model for a better understanding of tissue immunity under an immunopathological condition. *Exp Rev Clin Immunol*. 2015;11:391-407.
- Krutzik SR, Ochoa MT, Sieling PA, et al. Activation and regulation of Toll-like receptors 2 and 1 in human leprosy. *Nat Med*. 2003;9:525-532.
- Krutzik SR, Tan B, Li H, et al. TLR activation triggers the rapid differentiation of monocytes into macrophages and dendritic cells. *Nature*. 2005;11:653-660.
- Maneglier B, Rogez-Kreuz C, Cordonnier P, et al. Simultaneous measurement of kynurenine and tryptophan in human plasma and supernatants of cultured human cells by HPLC with coulometric detection. *Clin Chem*. 2004;50:2166-2168.
- Robinson CM, Shirey KA, Carlin JM. Synergistic Transcriptional Activation of IDO by IFN- $\gamma$  and TNG- $\alpha$ . *J Interferon Cytokine Res*. 2006;23:413-421.
- Modlin R. The innate immune response in leprosy. *Curr Opin Immunol*. 2010;22(1):48-54.
- Pinheiro RO, Schmitz V, Silva BJA, et al. Innate immune responses in leprosy. *Front Immunol*. 2018;9:518.
- Blumenthal A, Nagalingam G, Huch JH, et al. M. tuberculosis induces potent activation of IDO-1, but this is not essential for the immunological control of infection. *PLoS ONE*. 2012;7:37314.
- Ye Z, Yue L, Shi J, Shao M, Wu T. Role of IDO and TDO in cancers and related diseases and the therapeutic implications. *J Cancer*. 2019;10:2771-2782.
- Fujigaki S, Saito K, Sekikawa K, et al. Lipopolysaccharide induction of Indoleamine 2,3-dioxygenase is mediated dominantly by an IFN-gamma-independent mechanism. *Eur J Immunol*. 2001;31:2313-2318.
- Liu YN, Peng YL, Lei-Liu TY, et al. TNF $\alpha$  mediates stress-induced depression by upregulating Indoleamine 2,3-dioxygenase in a mouse model of unpredictable chronic mild stress. *Eur Cytokine Netw*. 2015;26:15-25.
- Chu W. Tumor necrosis factor. *Cancer Lett*. 2013;328:222-225.
- Montoya D, Modlin RL. Learning from leprosy. Insight into the human innate immune response. *Adv Immunol*. 2010;105:1-24.
- Underhill DM, Ozinsky A, Smith KD, Aderem A. Toll-like receptor-2 mediates mycobacteria-induced proinflammatory signaling in macrophages. *Proc Natl Acad Sci U S A*. 1999;96:14459-14463.
- Gehring AJ, Rojas RE, Canaday DH, Lakey DL, Harding CV, Boom WH. The *Mycobacterium tuberculosis* 19-kilodalton lipoprotein inhibits gamma interferon-regulated HLA-DR and Fc $\gamma$ R1 on human macrophages through Toll-like receptor 2. *Infect Immun*. 2003;71:4487-4497.
- Liu H, Komai-Koma M, Xu D, Liew FY. Toll-like receptor 2 signaling modulates the functions of CD4 +CD25+ regulatory T cells. *Proc Natl Acad Sci U S A*. 2006;103:7048-7053.
- Garg A, Barnes PF, Roy S, et al. Prostaglandin E2-dependent expansion of regulatory T cells in human *Mycobacterium tuberculosis* infection. *Eur J Immunol*. 2008;38:459-469.
- Tounsi N, Meghari S, Moser M, Djerdjouri B. Iyosphosphatidylcholine exacerbates *Leishmania* major-dendritic cell infection through interleukin-10 and a burst in arginase1 and Indoleamine 2,3-dioxygenase activities. *Int Immunopharmacol*. 2015;25:1-9.
- Alameddine J, Godefroy E, Papargyris L, et al. *Faecalibacterium prausnitzii* skews human dC to prime IL10-producing T cells through TLR2/6/JNK signaling and IL-10, IL-27, CD39, and IDO-1 induction. *Front Immunol*. 2019;10:1-11.

29. Polycarpou A, Holland MJ, Karageorgiou I, et al. *Mycobacterium leprae* activates Toll-like receptor-4 signaling and expression on macrophages depending on previous Bacillus Calmette-Guerin vaccination. *Front Cell Infect Microbiol.* 2016;6:72.
30. Shon WJ, Lee YK, Shin JH, Choi EY, Shin DM. Severity of DSS-induced colitis is reduced in Ido1-deficient mice with down-regulation of TLR-MyD88-NF-kB transcriptional networks. *Sci Rep.* 2015;5: 1-12.
31. Mellor AL, Lemos H, Huang L. Indoleamine 2,3 dioxxygenase and tolerance: where are we now? *Front Immunol.* 2017;8:1360.
32. Ilic N, Gruden-Movsesijan A, Cvetkovic J, et al. *Trichinella spiralis* excretory-secretory products induce tolerogenic properties in human dendritic cells via Toll-like receptors 2 and 4. *Front Immunol.* 2018; 9:11.
33. Jurado-Manzano BB, Zavala-Reyes D, Turrubiarres-Martínez EA, Portales-Pérez DP, González-Amaro R, Layseca-Espinosa E. FICZ generates human tDCs that induce CD4<sup>+</sup> CD25<sup>high</sup> Foxp3<sup>+</sup> Treg-like cell differentiation. *Immunol Lett.* 2017;190: 84-92.
34. Palermo ML, Pagliari C, Trindade MAB, et al. Increased expression of regulatory T cells and down-regulatory molecules in lepromatous leprosy. *Am J Trop Med Hyg.* 2012;86:878-883.

## SUPPORTING INFORMATION

Additional information may be found online in the Supporting Information section at the end of the article.

**How to cite this article:** Oliveira JAP, Gandini M, Sales JS, et al. *Mycobacterium leprae* induces a tolerogenic profile in monocyte-derived dendritic cells via TLR2 induction of IDO 2,3-dioxygenase. *J Leukoc Biol.* 2021;110:167–176. <https://doi.org/10.1002/JLB.4A0320-188R>