

Original paper

A new species of the genus *Cryptocanthon* from Peru (Coleoptera: Scarabaeidae: Scarabaeinae)

Nueva especie del género *Cryptocanthon* de Perú (Coleoptera: Scarabaeidae: Scarabaeinae)

 ALFREDO EDGARDO GIRALDO-MENDOZA

Universidad Nacional Agraria La Molina – Museo de Entomología Klaus Raven Büller. Av. La Molina s/n, Lima 12, Lima, Perú.

Responsible editor: Magdalena Cruz Rosales

ABSTRACT. A new brachypterous species of Scarabaeinae, *Cryptocanthon iskaypachak* **sp. nov.** is described, discussed and illustrated based on specimens collected in Cajamarca department. The new species is the third of the genus *Cryptocanthon* recorded from Peru.

Key words: Demarziellini; dung beetles; Neotropics; South America; Systematics

RESUMEN. Se describe, discute e ilustra una nueva especie braquíptera de Scarabaeinae, *Cryptocanthon iskaypachak* **sp. nov.** con base en especímenes recolectados en el departamento de Cajamarca. La nueva especie es la tercera registrada para el género *Cryptocanthon* en Perú.

Palabras clave: Demarziellini; escarabajos coprófagos; Neotrópico; Sudamérica; Sistemática

INTRODUCTION

The genus *Cryptocanthon* was described by Balthasar (1942) to place the dung beetle species *Cryptocanthon paradoxus* Balthasar, 1942 based on a single male specimen from Loja province (Villonaco) in southern Ecuador. During the eight subsequent decades, 41 species were added to the genus, eight



OPEN ACCESS

***Corresponding author:**

 Alfredo Edgardo Giraldo-Mendoza
aegmendoza@gmail.com

Cite:

Giraldo-Mendoza, A. E. (2022) A new species of the genus *Cryptocanthon* from Peru (Coleoptera: Scarabaeidae: Scarabaeinae). *Acta Zoológica Mexicana (nueva serie)*, 38, 1–11. [10.21829/azm.2022.3812443](https://doi.org/10.21829/azm.2022.3812443)
elocation-id: e3812443

Received: 20 August 2021

Accepted: 28 January 2022

Published: 10 February 2022



species from Brazil, Colombia, Mexico and Venezuela (Howden, 1973), two species from Colombia and Mexico (Howden, 1976), two species from Costa Rica and Panama (Howden & Gill, 1987), 22 species from Colombia, Costa Rica, Ecuador, French Guiana, Mexico, Panama and Venezuela (Cook, 2002), three species from Colombia (Arias & Medina, 2014), one species from Mexico (Mora-Aguilar & Delgado, 2018) and three species from Colombia (Martínez-Revelo *et al.*, 2020). A complete list of *Cryptocanthon* species is presented in Table 1.

The systematic position of *Cryptocanthon* is still doubtful, although traditionally it has been placed in the tribe Deltochilini (= Canthonini), it should be included in the tribe Demarziellini according to available morphological evidence (Vaz-de Mello, 2007; Martínez-Revelo *et al.*, 2020). While the highest species richness for this genus is in Colombia with 15 species (Martínez-Revelo *et al.*, 2020), Mexico with ten species (Mora-Aguilar & Delgado, 2018) and Ecuador with six species (Chamorro *et al.*, 2019), only two species have been recorded from Peru (Ratcliffe *et al.*, 2015). The two species previously recorded from Peru are the following ones: *Cryptocanthon campbellorum* Howden, 1973, known from Brasília and Pará in Brazil, Caquetá, Meta and Vaupes in Colombia and Madre de Dios in Peru (Cook, 2002; Arias & Medina, 2014; Martínez-Revelo *et al.*, 2020), and *C. paradoxus*, known from Ecuador (Loja, Villonaco) and Peru (Piura, Ayabaca) (Chamorro *et al.*, 2019; Juárez-Noe & Gonzalez-Coronado, 2021).

The objective of this paper is to describe a new species of *Cryptocanthon* from Cerro Coymolache, Hualgayoc, in northern Peru (Cajamarca).

MATERIALS AND METHODS

The author reviewed Scarabaeinae specimens housed at Museo de Entomología Klaus Raven Büller (MERKB), during which specimens attributed to a new species of the genus *Cryptocanthon* were found. Morphological terms were applied to make a consensus between basic terminology for Scarabaeinae genera (Vaz de Mello *et al.*, 2011), with the most comprehensive review of the genus (Cook, 2002) and most recent species descriptions (Martínez-Revelo *et al.*, 2020). The phylogenetic species concept (Wheeler & Platnick, 2000) is followed in this work, which defines a species as the smallest aggregation of populations diagnosable by a unique combination of character states.

Type specimens of the new species were deposited in Museo de Entomología Klaus Raven Büller (MEKRB) and Museo de Historia Natural Javier Prado (MUSM). Label data for the new species is quoted verbatim; different lines of the label are separated by a diagonal slash (/), and different labels are indicated by square brackets ([]).

When it was necessary, specimens were dissected for better examination. Parameres were extracted, treated for 10 minutes in 20 % KOH, washed with distilled water and adhered to a small piece of cardboard. The first approach to male specimens took into account the extraction of endophallites (Génier, 2019), since these sclerites have proved to be distinctive between *Cryptocanthon* species (Arias & Medina, 2014; Martínez-Revelo *et al.*, 2020). However, the detailed dissections that are required were not performed, due to a combination of lack of skills and fine

instruments, both essential for the proper extraction and manipulation of structures measuring less than 0.5 mm; these techniques were detailed by Medina *et al.* (2013).

Specimens were photographed with a Canon EOS Rebel T5i DSLR, equipped with Macro lens and rail. Each image includes a series of photos taken in different planes, which were stacked with the Combine ZP software (Hadley, 2006). Drawings were done by the prints of photographs, observations with a stereomicroscope and digital improving with graphic design software. Distribution map was prepared using SimpleMapp (Shorthouse, 2010).

Table 1. List of known species of genus *Cryptocanthon*, sorted according to their description date.

	Species	Type locality	Country records
1	<i>C. paradoxus</i> Balthasar, 1942	Loja (Ecuador)	Ecuador, Peru
2	<i>C. lobatus</i> Howden, 1973	Veracruz (Mexico)	Mexico
3	<i>C. brevisetosus</i> Howden, 1973	Chiapas (Mexico)	Mexico, Guatemala
4	<i>C. cristobalensis</i> Howden, 1973	Chiapas (Mexico)	Mexico
5	<i>C. parvus</i> Howden, 1973	Valle (Colombia)	Colombia
6	<i>C. campbellorum</i> Howden, 1973	Brasilia (Brazil)	Brazil, Colombia, Peru
7	<i>C. nebulinus</i> Howden, 1973	Aragua (Venezuela)	Venezuela
8	<i>C. humidus</i> Howden, 1973	Valle (Colombia)	Colombia, Panama
9	<i>C. peckorum</i> Howden, 1973	Amazonas (Colombia)	Colombia, Brazil
10	<i>C. altus</i> Howden, 1976	Santander (Colombia)	Colombia
11	<i>C. newtoni</i> Howden, 1976	Chiapas (Mexico)	Mexico
12	<i>C. chiriquinus</i> Howden & Gill, 1987	Chiriqui (Panama)	Panama
13	<i>C. lindemanae</i> Howden & Gill, 1987	Puntarenas (Costa Rica)	Costa Rica
14	<i>C. andersoni</i> Cook, 2002	Merida (Venezuela)	Venezuela
15	<i>C. gilli</i> Cook, 2002	Bolivar (Venezuela)	Venezuela
16	<i>C. galbao</i> Cook, 2002	Saül (French Guiana)	French Guiana
17	<i>C. pumilus</i> Cook, 2002	Vaupes (Colombia)	Colombia
18	<i>C. bochilae</i> Cook, 2002	Chiapas (Mexico)	Mexico
19	<i>C. denticulum</i> Cook, 2002	Chiriqui (Panama)	Panama
20	<i>C. howdeni</i> Cook, 2002	Chiapas (Mexico)	Mexico
21	<i>C. montebello</i> Cook, 2002	Chiapas (Mexico)	Mexico
22	<i>C. ocosingo</i> Cook, 2002	Chiapas (Mexico)	Mexico
23	<i>C. osaensis</i> Cook, 2002	Puntarenas (Costa Rica)	Costa Rica
24	<i>C. rayonensis</i> Cook, 2002	Chiapas (Mexico)	Mexico
25	<i>C. solisi</i> Cook, 2002	Heredia (Costa Rica)	Costa Rica
26	<i>C. curticrinis</i> Cook, 2002	Napo (Ecuador)	Ecuador
27	<i>C. escobari</i> Cook, 2002	Boyacá (Colombia)	Colombia
28	<i>C. foveatus</i> Cook, 2002	Cundinamarca (Colombia)	Colombia
29	<i>C. genieri</i> Cook, 2002	Napo (Ecuador)	Ecuador

	Species	Type locality	Country records
30	<i>C. lobipygus</i> Cook, 2002	Darien (Panama)	Panama
31	<i>C. medinae</i> Cook, 2002	Valle (Colombia)	Colombia
32	<i>C. napoensis</i> Cook, 2002	Napo (Ecuador)	Ecuador
33	<i>C. otonga</i> Cook, 2002	Cotopaxi (Ecuador)	Ecuador
34	<i>C. punctatus</i> Cook, 2002	Trujillo (Venezuela)	Venezuela
35	<i>C. urguensis</i> Cook, 2002	Napo (Ecuador)	Ecuador
36	<i>C. andradei</i> Arias & Medina, 2014	Risaralda (Colombia)	Colombia
37	<i>C. buriticaorum</i> Arias & Medina, 2014	Boyacá (Colombia)	Colombia
38	<i>C. pulidoae</i> Arias & Medina, 2014	Risaralda (Colombia)	Colombia
39	<i>C. chimalapensis</i> Mora-Aguilar & Delgado, 2018	Oaxaca (Mexico)	Mexico
40	<i>C. cephalopunctatus</i> Martínez-Revelo, Torres & Neita, 2020	Santander (Colombia)	Colombia
41	<i>C. encenillensis</i> Martínez-Revelo, Torres & Neita, 2020	Cundinamarca (Colombia)	Colombia
42	<i>C. mailinae</i> Martínez-Revelo, Torres & Neita, 2020	Antioquia (Colombia)	Colombia
43	<i>C. iskaypachak</i> sp. nov.	Cajamarca (Peru)	Peru

RESULTS AND DISCUSSION

Cryptocanthon iskaypachak Giraldo, new species (Figs. 1–3)

<http://zoobank.org/urn:lsid:zoobank.org:act:16176B7A-3E3B-457B-B1D4-747A6E6E49AF>

Type locality: Cerro Coymolache (06° 46' 45.52" S, 78° 36' 57.97" W), Provincia de Hualgayoc, Cajamarca, Peru.

Deposit of types: Museo de Entomología Klaus Raven Büller, Universidad Nacional Agraria La Molina, Lima, Peru (MERKB) and Museo de Historia Natural Javier Prado, Universidad Nacional Mayor de San Marcos, Lima, Peru (MUSM).

Type material. Holotype (male): [PERU, Cajamarca, Hualgayoc/ Hualgayoc, Cerro Coymolache/06° 46' 45.52" S, 78° 36' 57.97" W/XI-2014, R. Sánchez coll.] [Puna grassland, 3,500 m, pitfall traps], with red holotype label (MEKRB). Paratypes (five males and eight females) with the same label data, with yellow paratype labels, four males and six females (MEKRB), one male and two females (MUSM).

Diagnosis. The following combination of characters characterizes this species: vertical surface of clypeus not foveate; posterior pronotal angles obtuse and weakly incised, pronotal hypomeron not carinate, pseudopleuron lacking basal fovea; metathoracic wings greatly reduced, meso-metasternal suture straight; male protibiae weakly notched, male metatibiae with internal margin crenulated; apical abdominal sternite not lobed posteriorly; pygidium without median depression at base and parameres with apices distally widened and domed, not setose.

Description. Holotype. Male (Fig. 1A) (pinned and dissected). Total length: 5.2 mm; humeral width: 3.1 mm. Body shiny, dark brown, reddish on anterior margins of clypeus and pronotum, and on apices of femora and tibiae; covered with yellowish arched setae on pronotal and elytral margins, more conspicuous on posterior fourth of pronotum and posterior third of elytra.

Head (Fig. 2A). Clypeus not strongly produced anteriorly. Clypeal teeth moderately developed, widely separated, weakly reflexed. Clypeal emargination between teeth evenly rounded; vertical surface moderately broad, not foveate. Apical margin of clypeus lateral to teeth convexly sinuated. Disc of clypeus concave, with coarse punctures separated by a distance less than its diameter, punctures finer and sparser towards the anterior margin. Clypeogenal suture vaguely carinate. Gena slightly produced, forming a slight incision at junction with anterior margin of clypeus; coarsely, densely punctate such as the clypeus. Vertex with punctures similar to the clypeus and gena, without posteromedial depression.

Pronotum (Fig. 2B). Convex medially, laterally explanate, without shallow lateral depressions on posterior half. Anterior angles approximately right-angled. Lateral margin nearly straight both anterior and posterior to lateral angles. Posterior angles obtuse, each one preceded by a weak incision. Disc of pronotum with fine punctures separated by a distance equal to their diameter, punctures coarser and denser towards lateral margins. Pronotal hypomeron convex, with coarse and dense punctures, with evident microsculpture between punctures; not carinate posteriorly.

Elytra (Fig. 2C). Transversely and longitudinally strongly convex; lateral carinae not strongly elevated, without obvious setae on edge. Each elytron dorsally with seven striae, each stria indicated by two wavy lines. Elytral intervals slightly convex; with rows of punctures adjacent to striae. Pseudepipleuron broad, rather abruptly narrowed apically, lacking basal fovea, with impressed microsculpture; with scattered, minute punctures and erratic transverse wrinkles; striae 8 and 9 incomplete, faintly indicated by obscure wavy lines near middle; stria 10 incomplete, indicated by a row of punctures adjacent to epipleuron near middle. Epipleuron with erratically spaced, coarse punctures.

Wings. Metathoracic wings completely reduced. Venter (Fig. 2D). Prosternum surface with coarse punctures on lateral edge next similar to hypomeron. Mesosternum surface without evident microsculpture, with coarse punctures separated by less than their diameter. Meso-metasternal suture straight. Mesepisternum with similar surface than lateral margins of mesosternum. Metasternum surface without evident microsculpture, with coarse punctures separated by less than their diameter. Metepisternum with similar surface than lateral margins of metasternum.

Legs (Fig. 2E, G). Profemur with coarse and dense punctures on ventral surface. Protibia expanded medially in apical half, with three apical teeth on external margin, two distal teeth closest to each other and distant from proximal tooth, with smaller teeth between them; with a weakly pre-apical notch on inner margin. Protibial ventral surface smooth, with well-marked central carina, glabrous. Mesofemur and metafemur with coarse and dense punctuation on ventral surface. Mesotibia apically dilated, with two lateral rows of yellowish setae on lateral margins, from middle

to apex. Metatibia strongly curved; dorsal surface flattened with two rows of setae from middle to apex; internal lateral margin slightly crenulated from middle to apex, external lateral margin apically carinated.

Abdomen (Fig. 2I). Apical sternite with coarse and dense punctures, not lobed posteriorly. Pygidium convex, glabrous, with coarse punctures, without median depression at the base. Male genitalia (Fig. 2J, K). In lateral view, parameres with dome-shaped apices, distally directed downwards. In caudal view, parameres are wider basally, narrowed at middle and widened towards apices; apices are distally domed, not setose.




Figure 1. *Cryptocanthon iskaypachak* sp. nov. habitus: A) Male holotype, B) Female paratype. Scale = 1 mm.

Variation. Male. Total length: 4.8–5.2 mm; humeral width: 2.5– 3.0 mm. Female (Fig. 1B). Total length: 4.8–5.3 mm; humeral width: 2.7–3.0. Similar characters to male, except that clypeus and protibia bear prominent teeth. Also, tibiae unmodified, protibia without pre-apical notch on inner margin (Fig. 2F) and internal margin of metatibia smooth (Fig. 2H).

Etymology. The specific name alludes to the Quechua language word “Iskay Pachak” that means two hundred, because this species was described during the commemoration of bicentennial of Peruvian Republic (1821–2021).

Differential diagnosis. The two *Cryptocanthon* species previously recorded from Peru are clearly distinguishable from *C. iskaypachak*, because *C. campbellorum* has well-developed wings, vertical

surface of clypeus foveate and pseudopleuron with basal fovea and *C. paradoxus* has posterior pronotal angles strongly incised and pygidium with median depression at base. Also, parameres of *C. campbellorum* (Howden, 1973, Plate III, Fig. 22; Cook, 2002, Fig. 29) and *C. paradoxus* (Cook, 2002, Fig. 174) are distinct. According to its combination of morphological characters, *C. iskaypachak* is related to the species included in the so-called Andean species group (Cook, 2002), which have as synapomorphy a non-umbilicate type of head punctuation.




Figure 2. *Cryptocanthon iskaypachak* sp. nov. body parts: A) head, B) pronotum, C) inflexed portion of elytron, D) meso-metasternum, E) male protibia, F) female protibia, G) male metatibia, H) female metatibia, I) pygidium, J) aedeagus lateral view, K) aedeagus caudal view. Scale = 1mm.

The *C. iskaypachak* specimens are keyed to different species applying the available keys to South American *Cryptocanthon* species. Following the key proposed by Cook (2002), these are keyed to *Cryptocanthon napoensis* Cook, 2002, but this species has male protibia strongly notched near apex and pygidium with median depression at base. According to the key by Arias and Medina (2014), these are keyed to *Cryptocanthon pulidoae* Arias & Medina, 2014, but this species has posterior pronotal angles obtusely curved, not incised and male metatibia has internal margin smooth. Applying the key by Martínez-Revelo *et al.* (2020), these are keyed to *Cryptocanthon mailinae* Martínez-Revelo, Torres & Neita, 2020, but this species has pseudepipleuron with stria eight complete and pygidium with median depression at base. Also, parameres of *C. napoensis* (Cook, 2002, Fig. 164), *C. pulidoae* (Arias & Medina, 2014, Fig. 5f) and *C. mailinae* (Martínez-Revelo *et al.*, 2020, Fig. 11i) are distinct.

Although *C. iskaypachak* is clearly distinguishable from its relatives by the combination of characters presented, its endophallites have not been described. According to growing evidence, the endophallus characters of the Scarabaeinae are important for species delimitation (Zunino, 2012; Medina *et al.*, 2013) and could provide crucial information for to resolve phylogenies (Tarasov & Solodovnikov, 2011; Tarasov & Génier, 2015). The revision performed by Cook (2002) is still the most comprehensive work about the genus *Cryptocanthon*; however, species description, species groups and cladistic analysis presented by this author were supported largely on external morphological characters. After the male genitalia of the Scarabaeinae was studied for a worldwide sample of tribes (Medina *et al.*, 2013), the endophallites were incorporated as diagnostic elements in descriptions and determination of Colombian species of the genus *Cryptocanthon* (Arias & Medina, 2014; Martínez-Revelo *et al.*, 2020). Currently, internal sclerites of endophallus have been drawn for 14 *Cryptocanthon* species, that is, these are still unknown for around 67 % of known species, which with the exception of *Cryptocanthon peckorum* Howden, 1973, are recorded outside of Colombia.

Distribution and habitat (Fig. 3). Known only from type locality. According to label data, vegetation is Puna grassland, at 3,500 m of altitude, and specimens were collected using unbaited pitfall traps. Due to the presence of cattle in the area, herbivore dung could be a potential food source for this species. Most *Cryptocanthon* species inhabit litter of lowland and montane wet forests (Cook, 2002), as has been widely documented for *C. peckorum*, a species collected in tropical forests of Brazil and Colombia using dung-baited traps (Andresen, 2003; Ratcliffe, 2013; Cajaiba *et al.*, 2017; Noriega *et al.*, 2021). However, some species have been recorded from Paramo province, high cordilleras above 3,000 m of altitude in Colombia (*Cryptocanthon altus* Howden, 1973, *Cryptocanthon buriticaorum* Arias & Medina, 2014, *Cryptocanthon escobari* Cook, 2002) and Venezuela (*Cryptocanthon andersoni* Cook, 2002, *Cryptocanthon punctatus* Cook, 2002). As far as we known, it is the first *Cryptocanthon* recorded in these ecological conditions. As can be seen in the distribution map, *C. iskaypachak* is spatially segregated from *C. campbellorum* and *C. paradoxus*, species that inhabit lowland and montane wet forests in Peru, respectively.




Figure 3. Distributional map of *Cryptocanthon* species recorded in Peru, including South American neighboring countries: *C. iskaypackak* (red circle), *C. paradoxus* (white squares) and *C. campbellorum* (black triangles).

ACKNOWLEDGMENTS. The author thanks to Clorinda Vergara for her constant support at MEKRB and Raquel Sánchez for kindly donation of specimens. Also, to two anonymous reviewers for their valuable suggestions that improved a previous version of the manuscript.

LITERATURE CITED

- Andresen, E.** (2003) Effect of forest fragmentation on dung beetle communities and functional consequences for plant regeneration. *Ecography*, 26, 87–97.
- Arias, J. A., Medina, C. A.** (2014) Three new species of *Cryptocanthon* Balthasar, 1942 (Coleoptera: Scarabaeidae: Scarabaeinae) from Colombia. *Caldasia*, 36 (1), 165–180.
<https://doi.org/10.15446/caldasia.v36n1.43898>
- Balthasar, V.** (1942) Neue Scarabaeiden aus Süd-Amerika. 72. Beitrag zur Kenntnis der Scarabaeiden. (Col.). *Casopis Ceskoslovenske Spolecnosti Entomologicke*, 39, 36–44.
- Cajaiba, R. L., Perico, E., Dalzochio, M. S., da Silva, W. B., Bastos, R., Cabral, J. A., Santos, M.** (2017) Does the composition of Scarabaeidae (Coleoptera) communities reflect the extent of land use changes in the Brazilian Amazon? *Ecological Indicators*, 74, 285–294.
<https://doi.org/10.1016/j.ecolind.2016.11.018>
- Chamorro, W., Marin-Armijos, D., Asenjo, A., Vaz-De-Mello, F. Z.** (2019) Scarabaeinae dung beetles from Ecuador: a catalog, nomenclatural acts, and distribution records. *ZooKeys*, 826, 1–343.
<https://doi.org/10.3897/zookeys.826.26488>

- Cook, J.** (2002) A revision of the Neotropical genus *Cryptocanthon* Balthasar (Coleoptera: Scarabaeidae: Scarabaeinae). *Coleopterists Society Monograph*, 1, 1–96.
[https://doi.org/10.1649/0010-065x\(2002\)56\[3:arotng\]2.0.co;2](https://doi.org/10.1649/0010-065x(2002)56[3:arotng]2.0.co;2)
- Génier, F.** (2019) Endophallites: a proposed neologism for naming the sclerotized elements of the insect endophallus (Arthropoda: Insecta). *Annales de la Société entomologique de France, New Series*, 55, 482–484.
<https://doi.org/10.1080/00379271.2019.1685907>
- Hadley, A.** (2006) CombineZM public domain image processing software. Available at: <https://combinezp.software.informer.com> (accessed on 20 July 2021).
- Howden, H. F.** (1973) Revision of the New World genus *Cryptocanthon* Balthasar (Coleoptera: Scarabaeidae). *Canadian Journal of Zoology*, 51, 39–48.
<https://doi.org/10.1139/z73-007>
- Howden, H. F.** (1976) New species in the genera *Bdelyroopsis*, *Cryptocanthon* and *Drepanocerus* (Coleoptera: Scarabaeidae: Scarabaeinae). *Proceedings of the Entomological Society of Washington*, 78, 95–103.
- Howden, H. F., Gill, B. D.** (1987) New species and new records of Panamanian and Costa Rican Scarabaeinae (Coleoptera: Scarabaeidae). *Coleopterists Bulletin*, 41, 201–224.
- Juárez-Noé, G., González-Coronado, U.** (2021) Actualización a la lista de coleópteros (Insecta: Coleoptera) del bosque de neblina de Cuyas, Ayabaca-Región Piura, Perú. *Graellsia*, 77 (1), e126.
<https://doi.org/10.3989/graellsia.2021.v77.278>
- Martínez-Revelo, D. E., Torres, E., Neita-Moreno, J. C.** (2020) El género *Cryptocanthon* (Coleoptera: Scarabaeidae) en Colombia: descripción de especies nuevas, distribución geográfica y conservación. *Revista Mexicana de Biodiversidad*, 91, e913156.
<https://doi.org/10.22201/ib.20078706e.2020.91.3156>
- Medina, C. A., Molano, F., Scholtz, C.** (2013) Morphology and terminology of dung beetles (Coleoptera: Scarabaeidae: Scarabaeinae) male genitalia. *Zootaxa*, 3626 (4), 455–476.
<https://doi.org/10.11646/zootaxa.3626.4.3>
- Mora-Aguilar, E. F., Delgado, L.** (2018) A new species of *Cryptocanthon* Balthasar (Coleoptera: Scarabaeidae: Scarabaeinae) from the Region of Chimalapas, Oaxaca, Mexico. *The Coleopterists Bulletin*, 72, 792–796.
<https://doi.org/10.1649/0010-065x-72.4.792>
- Noriega, J. A., Santos, A. M. C., Calatayud, J., Chozas, S., Hortal, J.** (2021) Short- and long-term temporal changes in the assemblage structure of Amazonian dung beetles. *Oecologia*, 195, 719–736.
<https://doi.org/10.1007/s00442-020-04831-5>
- Ratcliffe, B. C.** (2013) The dung- and carrion-feeding scarabs (Coleoptera: Scarabaeoidea) of an Amazonian Blackwater rainforest: results of a continuous, 56-week, baited-pitfall trap study. *The Coleopterist Bulletin*, 67 (4), 481–520.
<https://doi.org/10.1649/0010-065X-67.4.481>
- Ratcliffe, B. C., Jameson, M. L., Figueroa, L., Cave, R. D., Paulsen, M. J., Cano, E., Beza-Beza, C., Jimenez-Ferbans, L., Reyes-Castillo, P.** (2015) Beetles (Coleoptera) of Peru: A Survey of the Families. Scarabaeoidea. *Journal of the Kansas Entomological Society*, 88 (2), 186–207.

- Shorthouse, D. P.** (2010) SimpleMappr, an online tool to produce publication-quality point maps. Available at: <http://www.simplemappr.net> (accessed on 20 July 2021).
- Tarasov, S., Génier, F.** (2015) Innovative Bayesian and Parsimony phylogeny of dung beetles (Coleoptera, Scarabaeidae, Scarabaeinae) enhanced by ontology-based partitioning of morphological characters. *PLoS ONE*, 10 (3), e0116671.
<https://doi.org/10.1371/journal.pone.0116671>
- Tarasov, S., Solodovnikov, A.** (2011) Phylogenetic analyses reveal reliable morphological markers to classify mega-diversity in Onthophagini dung beetles (Coleoptera: Scarabaeidae: Scarabaeinae). *Cladistics*, 27, 1–39.
<https://doi.org/10.1111/j.1096-0031.2011.00351.x>
- Vaz-de Mello, F.** (2007) *Revisión taxonómica y análisis filogenético de la tribu Ateuchini (Coleoptera: Scarabaeidae: Scarabaeinae)*. Tesis doctoral, Instituto de Ecología, A.C. Xalapa, Veracruz.
- Vaz de Mello, F., Edmonds, W. D., Ocampo, F. C., Schoolmeesters, P.** (2011) A multilingual key to the genera and subgenera of the subfamily Scarabaeinae of the New World (Coleoptera: Scarabaeidae). *Zootaxa*, 2854, 1–73.
<https://doi.org/10.11646/zootaxa.2854.1.1>
- Wheeler, Q. D., Platnick, N. I.** (2000) The phylogenetic species concept (sensu Wheeler and Platnick). Pp. 55–69. In: Wheeler, Q. D., Meier, R. (Eds.). *Species Concepts and Phylogenetic Theory: A Debate*. Columbia University Press, New York, NY.
- Zunino, M.** (2012) Cuarenta años de anatomía de las piezas genitales en la taxonomía de los escarabajos Coleoptera: Scarabaeoidea: el estado del arte. *Dugesiana*, 18 (2), 197–206.
<https://doi.org/10.32870/dugesiana.v18i2.4027>