

RESEARCH

Open Access



# Eco-friendly management of *Spodoptera litura* (Lepidoptera: Noctuidae) in tomato under polyhouse and field conditions using *Heterorhabditis bacteriophora* Poinar, their associated bacteria (*Photorhabdus luminescens*), and *Bacillus thuringiensis* var. *kurstaki*

Neelam Thakur<sup>1</sup>, Preety Tomar<sup>1\*</sup> , Jaspreet Kaur<sup>1</sup>, Simranjeet Kaur<sup>1</sup>, Anuja Sharma<sup>1</sup>, Samiksha Jhamta<sup>1</sup>, Ajar Nath Yadav<sup>2</sup>, Harcharan Singh Dhaliwal<sup>2</sup>, Rajesh Thakur<sup>3</sup> and Seema Thakur<sup>3</sup>

## Abstract

**Background** Insect pests cause substantial agriculture losses annually, and their regulation through chemical pesticides instigates a prolonged detrimental impact on the environment and human health. The upsurge in difficulties such as pesticide residue, soil degradation and pest resistance acted as the pacesetter for research on biological control with a prime focus on entomopathogens. To aid in knowing the biocontrol potential of these creatures, the present work deals with the applications of entomopathogenic nematode *Heterorhabditis bacteriophora* EUPT-SD, and entomopathogenic bacteria (EPBs) *Photorhabdus luminescens* and *Bacillus thuringiensis* var. *kurstaki* against the 4th larval instar of *Spodoptera litura* (Fabricius) (Lepidoptera: Noctuidae) in the laboratory experiment, polyhouse as well as field condition.

**Results** The results demonstrated that EPB, *P. luminescens*, caused the highest 100% larval mortality at the highest concentration of  $5 \times 10^2$  CFU/ml after 96 h of infection, followed by *B. thuringiensis* var. *kurstaki* which resulted in 98% at a concentration of  $5 \times 10^2$  CFU/ml and 92% mortality at the treatment with *H. bacteriophora* EUPT-SD (140IJs/ml) under laboratory bioassay study. In the polyhouse and field evaluation tests, again the *P. luminescens* was recorded as the most effective, followed by *B. thuringiensis* and *H. bacteriophora*, respectively. It was found that the treated plots experienced lesser damage when compared to non-treated plots.

**Conclusion** Applications of these pathogens are nature friendly and are a proficient alternative to synthetic chemical insecticides. It is suggested from the present investigation that the use of *P. luminescens* was the best biocontrol agent to overcome the local pest problems of this region because it is safe for animals, humans, non-target insect pests, plants, as well as for the environment.

**Keywords** *Spodoptera litura*, Bioassay, Entomopathogens, Management, Tomato

\*Correspondence:

Preety Tomar  
preetytomar9@gmail.com

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## Background

Occurrences of abiotic and biotic stresses are the key factors in the crops' losses (Pandey et al. 2017). The main obstacles to crop productivity are the losses due to insect pests. The prominent management strategies against these pests are the dependence on chemicals spray (Brodeur et al. 2017). Enormous applications of pesticides for minimizing losses caused by these organisms showed the prejudicial effects on the beneficial living organisms. It also added pesticide residues in the food and promotes environmental pollution.

Environment sustainability and better human health are the crucial components that can be achieved through biological control methods to replace harmful chemical pesticides. Biopesticides have a wide range of benefits with less side effect (Thakur et al. 2020). In recent agriculture practices, sustainability is the prime aim which is achieved by utilizing organisms present in the environment in such a way that they can help in improving crop health, increasing yields and reducing pollution (Thakur et al. 2022). Researchers nowadays are interested in the usage of entomopathogenic nematodes (EPNs) as a bio-control agent due to their wide host range, eco-friendly nature, active host finding strategy, easy mass culturing and compatibility with standard irrigation equipment along with the ability to recycle them in the soil environment (Thakur et al. 2021; Tomar et al. 2022a). The nematodes live in a symbiotic association with the bacteria *Xenorhabdus* and *Photorhabdus* belonging to the family Enterobacteriaceae, which are gram-negative with non-fermentative rodlike appearance (Koppenhöfer 2007). These bacteria are virulent against Lepidoptera, Diptera, Coleoptera and Hemiptera orders of class Insecta (Vitta et al. 2018) as they produce one to many groups of bioactive secondary metabolites. *Bacillus thuringiensis* (*Bt*) along with several microbial species proves to be successful industrial and academic efforts. It is the spore-forming bacteria that is omnipresent and forms crystal proteins. These crystal proteins are also known as  $\delta$ -endotoxin and are toxic to insects.

*Spodoptera litura* (Fabricius, 1775) (Lepidoptera: Noctuidae), a lesser armyworm, beet armyworm, tobacco caterpillar and cutworm, is a serious pest in India, Pakistan, China, Japan (Ghaffar et al. 2002) and South Asia (Qin et al. 2004). These polyphagous insect pests are very destructive and cause damage to various host crops such as cotton, potato, capsicum, clover, tomato, okra, soybean and onion (Saleem et al. 2016). In the field conditions, the pest caused 26–100% losses in yield among approximately 150 species of plants (Rao et al. 1993). The young larvae of the pest enter deep into the fruits and contaminate the fruits, which affects the yield and causes losses economically (Czepak et al. 2013). This tobacco

caterpillar has developed resistance against different synthetic pesticides. The present study was planned to evaluate the potential of EPNs and their associated bacteria, against *S. litura* larvae under the laboratory, polyhouse and field conditions.

## Methods

### Bioassay experiment

EPNs were isolated from rhizospheric soils of field crops in Solan district of Himachal Pradesh (H.P.), India, through the soil baiting technique (Orozco et al. 2014). The extracted nematodes were identified as *Heterorhabditis bacteriophora* Poinar, 1976 morphologically and molecularly and denoted as *H. bacteriophora* EUPT-SD. Nematodes collected by the white traps were lifted in the disinfected double distilled water at a temperature of  $15 \pm 1$  °C for 6–14 days before usage (Shapiro-Ilan et al. 2002). To attain the viable pathogenic infective juveniles, these identified nematodes were then replicated by in vivo culturing using the last instar of *Corcyra cephalonica* Stainton, 1865 (Malan et al. 2014). The collected nematode suspension was poured over the container that had small pieces of the sterilized synthetic sponges at a temperature of 10–15 °C (Ramakuwela et al. 2015).

Nutrient agar–bromothymol blue–tetrazolium chloride (NBT) microbiological media was used for the isolation of EPNs-associated bacteria (*P. luminescens* EUPT-SD). Bromothymol blue (25 mg/ml) and triphenyl tetrazolium chloride (TTC) 40 mg/ml were added at the rate of 100  $\mu$ l to 100 ml media. Nutrient broth, bromothymol blue (BTB) and agar were mixed in distilled water to make a volume of 1000 ml. Initially, the pH of the medium was adjusted to 7.0–7.2 and the medium in 50–100 ml aliquots was then steam-autoclaved at the 15 psi of steam for about 20 min. The stock of TTC was filtered, sterilized and added @100  $\mu$ l in 100 ml to the media. Ampicillin 100  $\mu$ g/ml was added to the media at the level of 100  $\mu$ l of ampicillin stock solution, i.e., 100 mg ampicillin sodium salt/ml with sterilized Milli-Q water. Cholesterol 10  $\mu$ g/ml was added to the medium for the proper growth and reciprocation of the EPNs. Hundred infective juveniles of nematodes were picked with the help of a sharp needle. The collected juveniles were centrifuged (5000 rpm for 5 min) for the pellet formation. This pellet was then dissolved in 1% sodium hypochlorite solution for surface sterilization for about 5 min. After that 1 ml distilled water was applied for washing and centrifuged for 5 min at 5000 rpm. Autoclaved sterile mortar and pestle were used, and crushed IJs were used to make a loop. Streaking was done on the NBT agar media in the petri plates with sterile needles. Petri plates were incubated ( $28 \pm 1$  °C) for 48 h. Single bacterial colonies obtained were purified by re-streaking on the same medium and then incubation

again ( $28 \pm 1$  °C) for 48 h (Akhurst 1980). The cultures of bacteria isolated from the individual colonies were streak purified on NBT agar medium and incubated ( $28 \pm 1$  °C). The plates with the bacterial growth were maintained at 4 °C after 48 h and these plates were mentioned as the master plates. The fully grown bacterial colonies were preserved in nutrient broth (NB) as well as in 25% glycerol stock for future use. Commercially available formulations of *B. thuringiensis* var. *kurstaki* Mahastra (Strain: DOR BT-1, 0.5% W.P.) were bought from the marketplace and used for the experiment.

*Solanum lycopersicum* Linnaeus, 1753 (Heemsohna) seeds were procured from the market and grown as a nursery in the Departmental biological control polyhouse with temperature 19–24 °C (avg.), 70–75% humidity and 11 h light/13 h dark light regime. Post-germinated tomato seedlings (28–30 days) were shifted into the plastic pot (20 × 20 × 14 cm) filled with rice husk and soil. Irrigation was applied twice daily and the seedlings were nourished with the Agrobium's® (NPK-8:25:25, 3–4 g/plant) at 15 days intervals. This was done to establish the insect breeding stalk as more numbers of insects were required for the bioassay experiments.

Adult and the larvae of *S. litura* were gathered from the University Agricultural Farms of Eternal University. As a feed, castor leaves were used and kept at  $28 \pm 1$  °C temperature for the development (Patil et al. 2014). The adults were shifted to the chimney for egg laying that already contained sucrose syrup for the feeding (Santaram 1985). The egg masses were found in clusters usually after 3–5 days of mating. The breeding stock of *S. litura* was retained by relocating the larvae of *S. litura* on the tomato seedlings under the plastic pots

with 05 larvae/plant. The pot was placed in entomological cages (40 cm<sup>3</sup>) for confinement. The *S. litura* life cycle lasted approximately 33–39 days under polyhouse conditions.

#### Bioassay experiment

##### *Heterorhabditis bacteriophora* EUPT-SD, *P. luminescence* and *B. thuringiensis* var. *kurstaki* applications

Bioefficacy of different treatments of EPNs (*H. bacteriophora* EUPT-SD), *P. luminescens* and *B. thuringiensis* var. *kurstaki* were evaluated against *S. litura* (4th instar), alone and also in combination (Table 1) in the laboratory. Bioefficacy tests were performed using 3 treatments at the 5 different concentrations along with the absolute control in the Petri plate. The nematode concentrations were 20, 40, 60, 80 and 100 IJs/ml. Laboratory-reared active 4th instar larvae were used to check the bioefficacy in the laboratory. These larvae were kept under Petri dishes (10 larvae/plate) lined with Whatman filter paper and nematode suspension 1–2 ml was sprinkled into the plates that contained 05 concentrations, namely 20, 40, 60, 80 and 100 IJs/ml. Diet (natural/artificial) was provided to the larvae for feeding, and after that the Petri dish was incubated at  $25 \pm 1$  °C. Mortality among the target insect pest was examined after 24, 48, 72 and 96 h of inoculation. At the same time, colonies of *P. luminescens* and *B. thuringiensis* were grown on nutrient agar (NA) media and nutrient broth (NB) was prepared. From the prepared broth, *P. luminescens* concentration was adjusted to  $1 \times 10^2$ ,  $2 \times 10^2$ ,  $3 \times 10^2$ ,  $4 \times 10^2$  and  $5 \times 10^2$  CFU/ml and used against 4th instar larvae. The *B. thuringiensis* var. *kurstaki* concentration was also adjusted to  $1 \times 10^2$ ,  $2 \times 10^2$ ,  $3 \times 10^2$ ,  $4 \times 10^2$  and

**Table 1** Biocontrol agent used against *Spodoptera litura* larvae (4th instar) in bioassay experiment under laboratory, under polyhouse and field conditions

| <i>Heterorhabditis bacteriophora</i> (IJs/ml)  | <i>Photorhabdus luminescence</i> (CFU/ml) | <i>Bacillus thuringiensis</i> var. <i>kurstaki</i> (CFU/ml) |
|--|---|---|
| <i>Different concentrations used against 4th instar larvae of Spodoptera litura in bioassay experiment under laboratory conditions</i> |   |   |
| 20   | $1 \times 10^2$                           | $1 \times 10^2$   |
| 40   | $2 \times 10^2$                           | $2 \times 10^2$   |
| 60   | $3 \times 10^2$                           | $3 \times 10^2$   |
| 80   | $4 \times 10^2$                           | $4 \times 10^2$   |
| 100  | $5 \times 10^2$                           | $5 \times 10^2$   |
| Absolute control   | Absolute control                          | Absolute control  |
| <i>Different concentrations used against S. litura larvae (4th instar) in polyhouse and field conditions</i>                           |   |   |
| 80   | $2 \times 10^2$                           | $2 \times 10^2$   |
| 100  | $3 \times 10^2$                           | $3 \times 10^2$   |
| 120  | $4 \times 10^2$                           | $4 \times 10^2$   |
| 140  | $5 \times 10^2$                           | $5 \times 10^2$   |
| Absolute control   | Absolute control                          | Absolute control  |

$5 \times 10^2$  CFU/ml and was used against the 10 larvae of target insect. In control, only distilled water (1–2 ml) was used. The diet was provided to feed them and the Petri dish was again incubated at  $25 \pm 1^\circ\text{C}$ . Insect mortality was tested after 24, 48, 72 and 96 h of inoculation. Every treatment was replicated 05 times and insect larval mortality was measured after every 24 h of application. The experiment was performed twice, and data were pooled for statistical analysis.

#### Polyhouse phase

##### *Heterorhabditis bacteriophora* EUPT-SD, *P. luminescens* and *B. thuringiensis* var. *kurstaki* applications

*Spodoptera litura* vulnerability against nematodes *H. bacteriophora* EUPT-SD was examined by accomplishing an experiment on the tomato at its vegetative stages. This experiment was performed in a departmental polyhouse ( $44 \times 36$  m) with an area of  $22 \times 11$  m that was divided into two different blocks that contained 26 plots of  $2 \text{ m}^2$ . The pre-grown tomato seedlings (20 per plot) were transplanted into these plots with an average distance of 20 cm. The cultured larvae of *S. litura* (4th instar) were transferred to each plant at the rate of 5 larvae per plant and 280 adults were also exposed to each block 24 h prior to the applications of a biocontrol agent. After 24 h of insect inoculation, the different suspensions of infective juveniles (combined with 0.3% polysorbate 20) were applied using a sprayer on abaxial and adaxial surfaces. Each treatment was replicated five times and the percentage data on the leaf damage and larval mortality was examined every 24 h for 15 days. The level of damage was calculated by adopting a graphical method, earlier described by Tomar et al. (2022d). The trial was executed under a randomized block design and the statistical analysis was performed by calculating the analysis of variance (ANOVA).

#### Field phase

##### *Heterorhabditis bacteriophora* EUPT-SD, *P. luminescens* EUPT-SD and *B. thuringiensis* var. *kurstaki* applications

A field experiment was conducted in the agricultural field, located at ( $30.76^\circ$  N and  $77.29^\circ$  E at 1027 m altitude), with 14 to  $25^\circ\text{C}$  mean temperature and 50 to 64% RH. In transplanting,  $100 \text{ m}^2$  areas were prefabricated first and then the soil was allowed to recuperate before plowing. Agrobium's NPK compost was applied to this selected area. The selected region was divided into two blocks, each with  $50 \text{ m}^2$  area. These two blocks were further distributed into 50 plots of  $4 \text{ m}^2$  area. Each of these plots was transplanted with tomato seedlings 20 seedlings per plot with a plant-to-plant distance of 20 cm. After 28 days of uprooting, the pupae (800) and adult (280) were inoculated in the field, each block containing 400 pupae and 140 adults. At the phonological growth stage

of the plant, 3 applications of EPNs and EPB after mixing with 0.3% polysorbate 20, were used in accordance with the variation in population density of *S. litura*. The 1st foliar application of these pathogens was performed after 1–2 weeks of pest inoculation. The second application was performed at the vegetative stage, 1–2 weeks after the 1st application. The 3rd spray was applied at the reproductive stage (1–2 weeks after the 2nd application). These applications of these pathogens were performed in the evening time during the complete cropping period until harvesting as per the fluctuations in the *S. litura* population. The experiment was executed under a randomized block design, and the statistical variables such as damage and mortality were assessed by the method prescribed by Tomar et al. (2022a).


#### Statistical analysis

The analysis was done by calculating the analysis of variance (ANOVA). The experiment was performed twice and the statistical analysis was performed on OPSTAT software. The treatment of bioagents included 5 applications of *H. bacteriophora* EUPT-SD, *P. luminescens* and *B. thuringiensis* var. *kurstaki* 5 along with control. The statistical relationship among all the treatments was also calculated using Pearson's correlation analysis.

## Results

### Bioassay experiment

A laboratory bioassay experiment was performed in order to evaluate the insecticidal potential of *H. bacteriophora* EUPT-SD, *P. luminescens* and *B. thuringiensis* var. *kurstaki* against *S. litura* larvae (4th instar). Among the three treatments, the most effective treatments were the applications of *P. luminescens* that caused the highest mortality (100%) in the highest concentration of  $5 \times 10^2$  CFU/ml among *S. litura* larvae after 96 h of infection, although the highest concentrations of *B. thuringiensis* var. *kurstaki* ( $5 \times 10^2$  CFU/ml) and *H. bacteriophora* EUPT-SD (140IJs/ml) resulted in 98 and 92% larval mortality after 96 h of exposure (Fig. 1). Insect mortality represented considerable variations among the different concentrations of *P. luminescens* ( $F = 34.16$ ,  $DF = 5$ ,  $P < 0.05$ ) with the mortality percentages of 42, 64, 88, 100 and 100% 96 h post-application. Among different concentrations of *B. thuringiensis* var. *kurstaki*, the mortality percentages were 34, 56, 76, 84 and 98% 96 h post-application with  $F = 20.22$ ,  $DF = 5$ ,  $P < 0.05$ . The different concentrations of *H. bacteriophora* EUPT-SD showed the mortality percentages 34, 44, 60, 76 and 92% after 96 h of nematode application with  $F = 43.97$ ,  $DF = 5$ ,  $P < 0.05$ . The data demonstrated that all the treatments were highly significant and caused considerable mortality among insects.



**Fig. 1** Larval mortality by (A) *Heterorhabditis bacteriophora* strain EUPT-SD; (B) *Photorhabdus luminescens* and (C) *Bacillus thuringiensis* against *Spodoptera litura* larvae under the laboratory conditions

### Polyhouse phase

Foliar applications of *H. bacteriophora* EUPT-SD in tomato plants resulted in lower damage percentage when compared to absolute control ( $F=60.48$ ,  $DF=4$ ,  $P<0.05$ ). The plot treated with the highest nematode concentration showed minimum damage (Fig. 2). The damage percentage was also increased to 32.4 and 37.2%, after spray 1st, which further declined after spray 2nd, and reached minimum after spray 3rd. The data collected over the larvae mortality demonstrated that considerable variation was observed among the IJs concentration with percent mortality 48, 78, 85 and 90% ( $F=70.52$ ,  $DF=4$ ,  $P<0.05$ ) after 5 days of nematodes application (after the third spray) (Fig. 2). Initially, it was observed that under the polyhouse conditions *H. bacteriophora* EUPT-SD concentration of 140IJs/ml provides high protection to the tomato plants.

Applications of *P. luminescens* resulted in reduced percent damage when compared to absolute control ( $F=67.10$ ,  $DF=4$ ,  $P<0.05$ ). The highest bacterial inoculum showed minimum damage (Fig. 2). Although in the lowest inoculum concentrations, the damage percentage was also increased to 30 and 31%, after 1st spray that further declined after 2nd spray and minimum after 3rd spray. The data collected over the larvae mortality demonstrated that application of different concentrations of *P. luminescens* caused a high mortality that ranged from 58, 80, 98 and 100% ( $F=127.15$ ,  $DF=4$ ,  $P<0.05$ ) after the third spray (Fig. 2). It was observed that the highest concentration of *P. luminescens* provides high protection to the plants.

Various concentrations of *B. thuringiensis* var. *kurstaki* were also applied against *S. litura*, resulting in a lower damage percentage in comparison with absolute control ( $F=62.39$ ,  $DF=4$ ,  $P<0.05$ ). The damage percentage in *Bt*-treated plots was also increased to 36 and 38%, after spray 1st, which further declined after spray 2nd, and reached minimum after spray 3rd. The data collected over the larvae mortality demonstrated that significant variation was observed among the percent mortality 49, 78, 86 and 98% ( $F=106.23$ ,  $DF=4$ ,  $P<0.05$ ) after five days of nematodes application (after the third spray) (Fig. 2). It was found that under the polyhouse conditions *B. thuringiensis* var. *kurstaki* concentration is highly effective. Maximum damage with no mortality was observed in the plots treated with absolute control.

### Field phase


The insecticidal potential of all the entomopathogens was also evaluated under the field conditions. The foliar applications of *H. bacteriophora* EUPT-SD were applied at different concentrations against the larvae and pupae of *S.*

*litura* under field conditions that showed significant variations in the damage ( $F=42.84$ ,  $DF=4$ ,  $P<0.05$ ) (Fig. 3). The highest damage (100%) was recorded in the control. In the EPNs concentrations of 120IJs/ml and 140IJs/ml, the damage percentage ranged from 40 and 35% after 3rd nematode application. Initially, much low larval mortality was recorded that increased with the increase in the nematode application. Maximum larvae were killed after 3rd application of EPNs. A damage percentage of 35% remained constant up to harvesting, signifying the biocontrol potential of EPNs.

The foliar application of *P. luminescens* showed the least damage percentage of 30% after 1st spray, increased to 31% after 2nd spray and remained constant after 3rd spray ( $F=79.44$ ,  $DF=4$ ,  $P<0.05$ ) (Fig. 3). The highest damage was recorded in the control (100%). The maximum larval mortality was recorded among the highest concentration that resulted in 70% larval mortality after the 1st spray and 100% mortality after the 2nd and 3rd spray, respectively. The different concentrations of *Bt* were also applied against *S. litura* under field conditions. The data recorded demonstrated the significant variations in the damage  $F=61.34$ ,  $DF=4$ ,  $P<0.05$  (Fig. 3). The damage percentage ranged from 38 and 39% after the 3rd spray of the highest bacterial concentration. The highest damage (100%) was recorded in the control. Initially, 66% larval mortality was recorded after 1st spray that increased to 91% after 2nd spray and 96% after 3rd spray. Damage percentage of 39% remained constant up to harvesting, establishing the biological pest control.

### Discussion

Solanaceous crops such as tomato are widely cultivated in India and are the cash crops of many farmers which demonstrate their economic importance in their life. Huge losses to the economy were caused by armyworm larvae (*S. litura*) in the tomato field. Although applications of chemical pesticides might be reduced the larval population but also resulted in resistance among larvae. This resulted in the rapid outburst of the *Spodoptera* population in the field. Furthermore, relocation among the *Spodoptera* moths also increased their survivability and population in the field. Owing to the great losses by *Spodoptera* larvae, interest in adopting an eco-friendly method has developed. The application of a biocontrol agent reflected a better opportunity in the management of armyworm population, although there are some biotic factors (bacteria, fungi, virus, weeds, animal interference) and abiotic stresses (temperature, humidity, application time) that limit the effectiveness of these pathogens against susceptible pests. In the present investigation, applications of three biocontrol agents (*H. bacteriophora* EUPT-SD, *P. luminescens* and *B. thuringiensis*) were



**Fig. 2** Polyhouse evaluations; (A, C & E) damage; (B, D & F) larval mortality by *Heterorhabditis bacteriophora* strain EUPT-SD, *Photorhabdus luminescence* and *Bacillus thuringiensis* against *Spodoptera litura*





Fig. 2 continued



**Fig. 3** Field evaluations; (A, C & E) damage; (B, D & F) larval mortality, by *Heterorhabditis bacteriophora* strain EUPT-SD, *Photorhabdus luminescence* and *Bacillus thuringiensis* against *Spodoptera litura*




Fig. 3 continued

evaluated against *S. litura* larvae (4th instar) in the laboratory, polyhouse and field conditions. Among the three treatments, the most effective treatments were the applications of *P. luminescens* that caused the highest mortality (100%) in the highest concentration of  $5 \times 10^2$  CFU/ml among *S. litura* larvae after 96 h of infection.

Earlier, Salazar-Gutiérrez et al. (2017) applied *Photobacterium luminescens* strain SL0708 and its cell-free filtrate against *Spodoptera frugiperda* Smith, 1797 and *Galleria mellonella* Linnaeus, 1758 larvae in a bioassay experiment. They reported 100% larval mortality after 48 h of bacterial application at the rate of  $1 \times 10^3 - 1 \times 10^4$  CFU/larva. It was recorded that *H. bacteriophora* EUPT-SD (140IJs/ml) resulted in 92% larval mortality after 96 h of exposure. The insecticidal potential of *Photobacterium* and *Xenorhabdus* also was demonstrated by Adithya et al. (2020). Earlier, Yan et al. (2020) also reported the insecticidal effect of EPNs against tobacco cutworm. Sun et al. (2021) evaluated the biocontrol potential of *Heterorhabditis* and *Steinernema* nematodes against *S. litura* larvae (5th instar) and recorded 90% mortality in *S. litura* after 72 h. Similar observations were recorded by (Burana et al. 2022). Tomar et al. (2022b) reported that application of EPNs is an effective strategy in managing the 3rd and 4th larval instars of *S. litura* in laboratory conditions. Tomar et al. (2022c) reported that the native strain of EPNs was highly effective in managing the lepidopteran insect population including *S. litura*.

The highest concentration of *B. thuringiensis* var. *kurstaki* ( $5 \times 10^2$  CFU/ml) resulted in 98% larval mortality after 96 h of exposure. The study was also supported by the earlier investigation of Sajid et al. (2020) who recorded 100% larval mortality among *S. litura* (2nd instar larvae) after being treated with *B. thuringiensis*. Similarly, the effectiveness of *B. thuringiensis* granules (water dispersible) along with starch and adjuvant Tween20 against *Spodoptera* larvae was also described by Vimala Devi et al. (2021). Virulence of *Bt* cry proteins was also explained against lepidopteran insect pests by Singh et al. (2021).

Under polyhouse evaluation test, foliar applications of *H. bacteriophora* EUPT-SD at the rate of 140 IJs/ml in tomato plants resulted in lower damage percentage and high larval mortality (90%) than with absolute control. Earlier, Andaló et al. (2010) evaluated the efficacy of *Steinernema arenarium* Artyukhovsky, 1967 and *Heterorhabditis* sp. RSC02 against *S. frugiperda* under the laboratory and polyhouse conditions towards maize crop. They recorded 100 and 97.6% insect mortality under laboratory conditions and 77.5 and 87.5% mortality under greenhouse conditions, respectively. Tomar et al. (2022b) reported that applications of EPNs are highly effective in

controlling the insect population under polyhouse and also in field conditions. Applications of *P. luminescens* and *B. thuringiensis* var. *kurstaki* resulted in 100 and 98% larval mortality after the third spray under polyhouse conditions. The present study also supported by Thakur et al. (2022c) evaluated the individual and combined impact of entomopathogens *H. bacteriophora*, *B. bassiana* and *B. thuringiensis* against *Spodoptera* larvae and reported that applications of these entomopathogens were highly effective in managing the insect population under the laboratory as well as in greenhouse conditions.

Under field conditions also significant control in the damage and high larval mortality was recorded after treatment with *H. bacteriophora* EUPT-SD, *P. luminescens* and *B. thuringiensis* var. *kurstaki* against *Spodoptera* larvae. Earlier, Mohan et al. (2003) reported that 100 percent cabbage butterfly larvae were killed upon treatment with foliar applications of *Photobacterium* bacteria. Similarly, Uma et al. (2010) applied cellular and cell-free filtrate against aphids and reported a significant reduction in aphid nymph population. Navon (2000) also reported the application of *B. thuringiensis* in controlling the agricultural insect pests.

## Conclusions

Entomopathogens have been known to manage insect populations for years. During the present study, entomopathogenic nematode *H. bacteriophora* EUPT-SD, and entomopathogenic bacteria *P. luminescens* EUPT-SD and *B. thuringiensis* var. *kurstaki* were applied against the 4th instar larvae of *S. litura* under the laboratory, polyhouse and field conditions. The obtained results demonstrated that entomopathogenic bacteria *P. luminescens* caused a significant mortality rate, followed by *B. thuringiensis* var. *kurstaki* and then *H. bacteriophora* EUPT-SD under laboratory bioassay study. In addition, they were in the polyhouse experiments and in field conditions. So the treated plots experienced lesser damage when compared to non-treated plots. The applications of these pathogens are nature-friendly and are a proficient alternative to synthetic chemical insecticides.

## Abbreviations

|                         |                                      |
|-------------------------|--------------------------------------|
| <i>H. bacteriophora</i> | <i>Heterorhabditis bacteriophora</i> |
| IJs                     | Infective juveniles                  |
| EPNs                    | Entomopathogenic nematodes           |
| CFU                     | Colony-forming unit                  |
| %                       | Percent                              |
| Hrs                     | Hours                                |
| EPA                     | Environment protection agency        |

## Acknowledgements

The study was carried out in the Zoology laboratory at Eternal University Baru Sahib, Himachal Pradesh. The authors acknowledge the financial assistance provided by the Department of Science and Technology, Govt. of India (SP/

YO/506/2018-G). The authors are also thankful to Vice Chancellor, Eternal University, Baru Sahib, for providing necessary laboratory facilities.

#### Author contributions

NT gave the concept. NT and PT conducted the experiment and wrote the manuscript with statistical analysis. JK, SK, AS and SJ helped in conducting the survey to collect the insects from various fields. ANY, HSD, RT and ST gave their valuable suggestions during the field experiments. All authors have read and reviewed the manuscript.

#### Funding

The financial assistance is provided by the Department of Science and Technology, Govt. of India (SP/YO/506/2018-G).

#### Declarations

##### Ethics approval and consent to participate

All procedures performed in studies are in accordance with the ethical standards of the institutional and/or national research committee. We further declare that no animal was harmed during this study.

##### Consent for publication

Informed consent was obtained from all individual participants included in the study.

##### Competing interests

The authors declare that there is no conflict of interest.

##### Author details

<sup>1</sup>Department of Zoology, Akal College of Basic Sciences, Eternal University, Baru Sahib, Himachal Pradesh, India. <sup>2</sup>Department of Biotechnology, Dr. KSG Akal College of Agriculture, Eternal University, Baru Sahib, Himachal Pradesh, India. <sup>3</sup>Krishi Vigyan Kendra Kandaghat, Solan 173215, India.

Received: 21 October 2022 Accepted: 12 January 2023

Published: 26 January 2023

#### References

- Adithya S, Shivaprakash M, Sowmya E (2020) Evaluation of insecticidal activity of entomopathogenic bacteria *Photorhabdus* and *Xenorhabdus* against shoot and fruit borer *Earias vittella* (Lepidoptera: Noctuidae) of vegetable crops. *J Entomol Zoöl Stud* 8:2343–2348
- Akhurst R (1980) Morphological and functional dimorphism in *Xenorhabdus* spp., bacteria symbiotically associated with the insect pathogenic nematodes *Neoalectana* and *Heterorhabditis*. *Microbiology* 121:303–309
- Andaló V, Santos V, Moreira GF, Moreira CC, Moino Junior A (2010) Evaluation of entomopathogenic nematodes under laboratory and greenhouse conditions for the control of *Spodoptera frugiperda*. *Ciência Rural* 40:1860–1866
- Brodeur J, Hajek AE, Heimpel GE, Sloggett JJ, Mackauer M, Pell JK, Völkl W (2017) Predators, parasitoids and pathogens. In: Emden HF, van Harrington R (eds) *Aphids as crop pests*. CABI, Wallingford, pp 225–261
- Burana K, Ehlers RU, Nimkingrat P (2022) Entomopathogenic nematodes for control of the cotton cutworm *Spodoptera litura* in marigolds. *J App Entomol* 146:415–423
- Czepak C, Albernaz KC, Vivan LM, Guimarães HO, Carvalhais T (2013) Primeiro registro de ocorrência de *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae) no Brasil. *Pesqui Agropecu Trop* 43:110–113
- Ghaffar A, Attique MR, Naveed MR, Jan MT (2002) Host range and population dynamics of beet armyworm, *Spodoptera exigua* (Hubner) (Noctuidae: Lepidoptera) in cotton agroecosystem of Punjab, Pakistan. *Pak J Zool* 34:209–213
- Koppenhöfer AM (2007) Nematodes. In: Lacey LA, Kaya HK (eds) *Field manual of techniques in invertebrate pathology*. Springer, Dordrecht, pp 249–264
- Malan AP, Knoetze R, Tiedt L (2014) *Heterorhabditis noenieputensis* n. sp. (Rhabditida: Heterorhabditidae), a new entomopathogenic nematode from South Africa. *J Helminthol* 88:139–151
- Mohan S, Raman R, Gaur H (2003) Foliar application of *Photorhabdus luminescens*, symbiotic bacteria from entomopathogenic nematode *Heterorhabditis indica*, to kill cabbage butterfly *Pieris brassicae*. *Curr Sci* 84:1397–1397
- Navon A (2000) *Bacillus thuringiensis* insecticides in crop protection-reality and prospects. *Crop Prot* 19:669–676
- Orozco RA, Lee MM, Stock SP (2014) Soil sampling and isolation of entomopathogenic nematodes (Steinernematidae, Heterorhabditidae). *JoVE* 89:e52083
- Pandey P, Irulappan V, Bagavathiannan MV, Senthil-Kumar M (2017) Impact of combined abiotic and biotic stresses on plant growth and avenues for crop improvement by exploiting physio-morphological traits. *Front Plant Sci* 8:537
- Patil RA, Mehta DM, Jat BL (2014) Studies on life fecundity tables of *Spodoptera litura* Fabricius on tobacco *Nicotiana tabacum* Linnaeus. *Entomol Ornithol Herpetol* 3:1000118
- Qin H, Ye Z, Huang S, Ding J, Luo R (2004) The correlations of the different host plants with preference level, life duration and survival rate of *Spodoptera litura* Fabricius. *Chin J Eco-Agric* 12:40–42
- Ramakuwela T, Hatting J, Laing MD, Hazir S, Thiebaut N (2015) Effect of storage temperature and duration on survival and infectivity of *Steinernema innovationi* (Rhabditida: Steinernematidae). *J Nematol* 47:332
- Rao GR, Wightman J, Rao DR (1993) World review of the natural enemies and diseases of *Spodoptera litura* (F.) (Lepidoptera: Noctuidae). *Int J Trop Insect Sci* 14:273–284
- Sajid Z, Ali A, Usman M, Mujahid A, Jafar B, Kashif A, Bashir HS, Abbas Q, Tariq MU, Shakeel MD, Sultan Y (2020) Toxicity of *Bacillus thuringiensis* against second instar larvae of *Spodoptera litura* on different host plants. *J Sci Agric* 4:93–95
- Salazar-Gutiérrez JD, Castelblanco A, Rodríguez-Bocanegra MX, Teran W, Sáenz-Aponte A (2017) *Photorhabdus luminescens* subsp. *akhurstii* SL0708 pathogenicity in *Spodoptera frugiperda* (Lepidoptera: Noctuidae) and *Galleria mellonella* (Lepidoptera: Pyralidae). *J Asia Pac Entomol* 20:1112–1121
- Saleem M, Hussain D, Ghouse G, Abbas M, Fisher SW (2016) Monitoring of insecticide resistance in *Spodoptera litura* (Lepidoptera: Noctuidae) from four districts of Punjab, Pakistan to conventional and new chemistry insecticides. *Crop Protec* 79:177–184
- Shapiro-Ilan D, Gaugler R (2002) Production technology for entomopathogenic nematodes and their bacterial symbionts. *J Ind Microbiol Biotechnol* 28:137–146
- Singh D, Thayil SM, Sohal SK, Kesavan AK (2021) Exploration of insecticidal potential of Cry protein purified from *Bacillus thuringiensis* VIID1. *Int J Biol Macromol* 174:362–369
- Sun B, Zhang X, Song L, Zheng L, Wei X, Gu X, Cui Y, Hu B, Yoshika T, Abd-Elgawad MM (2021) Evaluation of indigenous entomopathogenic nematodes in Southwest China as potential biocontrol agents against (Lepidoptera: Noctuidae). *J Nematol* 53:1–17
- Santharam G (1985) Studies on the nuclear polyhedrosis virus of the tobacco cutworm, *Spodoptera litura* (Fabricius) (Noctuidae: Lepidoptera). Ph. D Thesis, Tamil Nadu Agricultural University, Coimbatore
- Thakur N, Tomar P, Sharma S, Kaur S, Sharma S, Yadav AN, Hesham AE (2022) Synergistic effect of entomopathogens against *Spodoptera litura* (Fabricius) under laboratory and greenhouse conditions. *Egypt J Biol Pest Control* 32:1–10
- Tomar P, Thakur N (2022) Biocidal potential of indigenous isolates of Entomopathogenic Nematodes (EPNs) against tobacco cutworm, *Spodoptera litura* Fabricius (Lepidoptera: Noctuidae). *Egypt J Biol Pest Control* 32:1–10
- Thakur N, Kaur S, Tomar P, Thakur S, Yadav AN (2020) Microbial biopesticides: current status and advancement for sustainable agriculture and environment. In: *New and future developments in microbial biotechnology and bioengineering*. Elsevier, pp 243–282
- Thakur N, Tomar P, Kaur S, Jhamta S, Thakur R, Yadav AN (2021) Entomopathogenic soil microbes for sustainable crop protection. In: *Soil microbiomes for sustainable agriculture*. Springer, pp 529–571
- Tomar P, Thakur N, Yadav AN (2022a) Indigenous entomopathogenic nematode as biocontrol agents for insect pest management in hilly regions. *Plant Sci Today* 8:51–59
- Tomar P, Thakur N, Yadav AN (2022b) Endosymbiotic microbes from entomopathogenic nematode (EPNs) and their applications as biocontrol

- agents for agro-environmental sustainability. *Egypt J Biol Pest Control* 32:1–19
- Tomar P, Thakur N, Sharma A (2022c) Infectivity of entomopathogenic nematode against the cabbage butterfly (*Pieris brassicae* L.) in polyhouse and in field condition. *Egypt J Biol Pest Control* 32(1):1–7
- Uma G, Prabhuraj A (2010) Effect of the entomopathogenic bacterium *Photorhabdus luminescens* to *Aphis gossypii* Glover. *J Biopestic* 3:100
- Vimala Devi PS, Duraimurugan P, Chandrika K, Vineela V (2021) Development of a water dispersible granule (WDG) formulation of *Bacillus thuringiensis* for the management of *Spodoptera litura* (Fab.). *Biocontrol Sci Technol* 31:850–864
- Vitta A, Thimpoo P, Meesil W, Yimthin T, Fukruksa C, Polseela R, Mangkit B, Tandhavanant S, Thanwisai A (2018) Larvicidal activity of *Xenorhabdus* and *Photorhabdus* bacteria against *Aedes aegypti* and *Aedes albopictus*. *Asian Pac J Trop Biomed* 8:31
- Yan F, Huang C, Wang X, Tan J, Cheng S, Wan M, Wang Z, Wang S, Luo S, Li A, Guo X (2020) Threonine ADP-ribosylation of ubiquitin by a bacterial effector family blocks host ubiquitination. *Mol Cell* 78:641–652

### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

---

Submit your next manuscript at ▶ [springeropen.com](https://www.springeropen.com)

---