



Heavy silicone oil tamponade: a multicentre experience

Max Davidson ,¹ Samir Dowlut,² Jufen Zhang,³ Khayam Naderi,² Teresa Sandinha,⁴ Megan Kay Wood ,⁵ Matthew Schneiders,⁶ Shohista Saidkasimova,⁷ Shamfa Peart,⁸ Ray Chaudhuri,⁸ Manish Gunda,⁹ Manzar Saeed,⁹ Florian Heussen,¹⁰ Johannes Keller,¹⁰ Sonali Tarafdar,¹¹ Aman Chandra²

To cite: Davidson M, Dowlut S, Zhang J, *et al*. Heavy silicone oil tamponade: a multicentre experience. *BMJ Open Ophthalmology* 2022;**7**:e001018. doi:10.1136/bmjophth-2022-001018

Received 18 March 2022
Accepted 13 November 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Sheffield Teaching Hospitals NHS Foundation Trust, Sheffield, UK

²Southend University Hospital, Southend, UK

³Anglia Ruskin University, Cambridge, UK

⁴St Paul's Eye Hospital, Liverpool, UK

⁵Alder Hey Children's NHS Foundation Trust, Liverpool, UK

⁶Norfolk & Norwich University Hospitals NHS Foundation Trust, Norfolk, UK

⁷Gartnavel General Hospital, Glasgow, UK

⁸Leicester Royal Infirmary, Leicester, UK

⁹Queen Elizabeth Hospital, King's Lynn, UK

¹⁰University Hospitals Bristol and Weston NHS Foundation Trust, Bristol, UK

¹¹Ninewells Hospital, Dundee, UK

Correspondence to

Dr Max Davidson; max.davidson@hotmail.co.uk

ABSTRACT

Objective To report multicentred use of the heavy silicone oil Densiron 68 for anatomical reattachment following rhegmatogenous retinal detachment (RRD) repair and its associated complications.

Methods and analysis Patients from seven vitreoretinal units within the UK that underwent RRD repair with Densiron 68 between January 2015 and December 2019 were identified. Primary outcome measures were primary and final reattachment rate, retained Densiron and failure rate. Secondary outcome measures were duration of tamponade, final visual acuity (VA) and complications of heavy silicone oil.

Results 134 eyes of 134 patients were involved in the study. Primary surgical success was achieved in 48.5%, while a final reattachment rate of 73.4% was observed. The mean duration of Densiron 68 tamponade was 139.5 days. Mean final VA was 1.01 (range 0–2.9). 8 eyes (6.0%) required long-term topical steroids for anterior uveitis, whereas none of the eyes required long-term pressure-lowering treatment. Emulsification rate was 10.7% (6 eyes).

Conclusion This is the largest real-world study on Densiron 68 in the UK. Densiron 68 facilitates tamponade of inferior retinal pathology and may be considered as an option for tamponade of inferior retinal pathologies.

INTRODUCTION

Rhegmatogenous retinal detachment (RRD) can cause significant vision loss, particularly if not treated in a timely fashion. For uncomplicated RRD, current surgical techniques carry a high anatomical success rate (>90%).^{1 2} Recognised risk factors for failure include choroidal detachment, hypotony, grade C proliferative vitreoretinopathy (PVR).^{3–5} RRD associated with inferior retinal breaks located at a position between 4 and 8 o'clock hours, particularly with concomitant PVR, remains a surgical challenge. Pars plana vitrectomy (PPV) with intraocular gases and standard silicone oil tamponades are unlikely to provide adequate support for such inferior peripheral retinal pathology.^{6 7} Recurrent RRD secondary to PVR is estimated to occur

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Densiron 68 and other heavy silicone oils have been used as tamponade for inferior retinal detachment (RD) with varying success rates and complication profiles.

WHAT THIS STUDY ADDS

⇒ This is the largest real-world data series on Densiron 68 in the UK, showing that Densiron 68 can achieve similar long-term reattachment rates and complications compared with conventional silicone oil.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ These data will better inform vitreoretinal colleagues about heavy silicone oil tamponade as a management strategy for inferior RD.

in about 5%–10% of all patients with failed primary RD repair.⁸

Retinal redetachment after surgery is likely to arise by several mechanisms: continued vitreoretinal traction, passage and propagation of fluid through an untreated retinal break, possibly due to intraocular fluid currents. In tamponade the gas or oil bubble prevents flow of fluid through tears in the retina.⁹ This allows retinal reattachment to be temporarily maintained while a permanent chorioretinal adhesion forms in response to retinopexy. Although inferior RD breaks without adequate tamponade have been reported to reattach,⁹ this is not replicated consistently in clinical practice and the search for adequate tamponade in these cases continues.

Endotamponades with a higher specific gravity within the vitreous cavity than aqueous are theorised to be useful for treating inferior retinal pathology.^{6 7 10} Examples include heavy oils and perfluorocarbon liquids. The former are used for longer periods. First generation heavy silicone oils (HSO) such as fluorosilicone and perfluorocarbons (introduced in

1990s) were marked with complications such as fibrinous reactions, cataract or even retinal necrosis.^{11–13} A number of newer generations of HSO of different compositions are available for use: Densiron 68 (Geuder), Oxane HD (Bausch & Lomb, mixture of Oxane 5700, mixed fluorinated and hydrocarbonated olefin (RMN3)) and Alaheavy (Alamedics).^{6,7}

Densiron 68 (Geuder AG, Neu-Ulm, Germany) is a mixture of perfluorohexyloctane (F_6H_8) and siluron 5000, giving a mid-range viscosity of 1400 mPas and gravity of 1.06 g/cm³.¹⁴ It is licensed for the treatment of inferior or posterior RD in cases of PVR, proliferative diabetic retinopathy and giant retinal tears.¹⁴ It was chosen as one of the endotamponades in the HSO study.¹⁵ Anatomical success rates with Densiron 68 varies from 33.3% to 81.3%.^{7,15,16}

Here, we present the largest evaluation of patterns of use, clinical outcomes and complications associated with the use of Densiron 68 as an internal tamponade in the treatment of RRDs among multiple vitreoretinal units within the UK.

MATERIALS AND METHODS

This was a retrospective, non-randomised study involving seven National Health Service vitreoretinal units within the UK; Bristol Eye Hospital (University Hospitals Bristol and Weston), Dundee eye clinic (Ninewells hospital, Dundee), Queen Elizabeth Hospital (King's Lynn, Norfolk), Leicester Royal Infirmary, St Paul's Eye Hospital (Royal Liverpool and Broadgreen University Hospitals), Norfolk & Norwich University Hospitals and Southend University Hospital. Surgical teams reported their outcomes using a standard anonymised spreadsheet. The tenets of the Declaration of Helsinki were followed in the study.

Inclusion criteria were patients receiving Densiron 68 during primary or recurrent RRD surgery during the period of January 2015 to December 2019 with a minimum follow-up of 3 months post-Densiron 68 removal. Cases receiving other heavy oil such as Densiron Xtra, Alaheavy or Oxane HD were excluded. Cases were collected via a combination of EPR, personal and theatre logs. All surgical procedures were performed by either vitreoretinal fellows or consultants.

Surgeons used their conventional vitreoretinal management to allow intraoperative reattachment of the retina, including use of retinotomies, drainage through breaks, retinectomies or use of perfluorooctane/perfluorodecalin. Patients were not instructed to undertake any particular posture.

Patients who required Densiron 68 tamponade were identified by the vitreoretinal units and their anonymised outcomes were pooled and analysed with Microsoft Excel 2020/GraphPad 2020. The primary outcome measures were primary success, final reattachment, qualified success and failure rates. Primary success was defined as achieving a flat retina at 3 months after one Densiron surgery and no subsequent retinal reintervention other

than Densiron removal. Final reattachment was defined as achieving a flat retina at 3 months after repeated surgery following the introduction of Densiron. Qualified success referred to the proportion of patients with a flat retina and Densiron retained rather than removed at 3 months. Failure was defined as inability to achieve a flat retina despite all efforts.

The secondary outcome measures were median duration of tamponade, postoperative visual acuity (VA), initial postoperative uveitis, long-term anterior uveitis, ocular hypertension (OHT) and emulsification of oil. Adequate follow-up was defined as at least 3 months after Densiron 68 removal.

Best-corrected VA (BCVA) was assessed using LogMAR. If Snellen VA had been used at any of the participating centres, this was converted to a LogMAR equivalent value. Very poor VA groups were assigned the following LogMAR equivalent: counting fingers (1.85), hand movement (2.3), light perception (2.6) and no light perception (2.9).¹⁷

The outcomes were analysed using Excel for mac 2020 and STATA statistical software V.14. A 5% significance level was used (two tailed). Continuous data were presented as mean (SD) or median (IQR) depending on the distribution of data and categorical variables were expressed as counts and percentages. χ^2 test or Fisher's exact test was used to compare proportions between groups. The independent t-test and paired sample t-test were used to compare means from parametric datasets, and the Mann-Whitney and Wilcoxon tests were used to compare non-parametric data.

No patients nor the public were involved in the design, conduct or reporting of this study.

RESULTS

One hundred and thirty four eyes of 134 patients were included in the study. During the study period, eleven patients (8.0%) had long-term Densiron 68 retention. The median age of patients was 64 years old (range 25–91 years), with 91 males (67.9%) and 43 females (32.1%).

The median (range) number of operations prior to Densiron 68 tamponade was 1 (0–4). All patients had RRD, with the most common indication being inferior retinal pathology (N=128; 93.4%) as displayed in [table 1](#). Eighty-six cases (64.2%) had failed previous RD surgery secondary to inferior pathology. The median (range) number of inferior retinal tears (between 4 and 8 o'clock associated with subretinal fluid) was 1 (0–7). Densiron 68 was used in cases presenting with all grades of PVR, including no PVR as shown in [table 2](#). All PVR C cases involved the inferior retina.

Seventy-two eyes (53.7%) were phakic, 63 (47.0%) were pseudophakic and 2 (1.5%) were aphakic at time of Densiron 68 insertion.

The median (IQR) duration for Densiron 68 tamponade was 107 days (70–156). Densiron 68 was used most commonly during recurrent RRD repair (87 eyes;

Table 1 Indications for Densiron 68 tamponade

Indications	N
Inferior RD	116
Inferior GRT	4
Inferior schisis RD	4
Inferior RD under light silicone oil	2
Dense VH due to wet AMD and inferior retinal tears	2
Multiple RD	1
Primary macular hole RD	1
PDR and TRD	2
Total PVR RD	1
Dropped nucleus with inferior RD	1
	134

AMD, age-related macular degeneration; GRT, giant retinal tear; PDR, proliferative diabetic retinopathy; PVR, proliferative vitreoretinopathy; RD, retinal detachment; TRD, tractional retinal detachment; VH, vitreous haemorrhage.

64.7%) with the remainder using it during primary RRD surgery.

Primary outcomes

Sixty-five patients (48.5%) achieved a flat retina with Densiron surgery without need for further retinal reoperation. Final reattachment was observed in 99 patients (73.9%). Eleven patients (8.2%) had long-term Densiron 68 retention, while failure was seen in 24 patients (17.9%). There was a trend for higher final reattachment rate after Densiron 68 removal for recurrent RRD (68 eyes (79.1%)) compared with those who had primary RRD (31 eyes (64.6%)) although this did not quite reach statistical significance (table 3; $p=0.067$). Table 4 shows subgroups of the 48 patients that underwent primary RRD surgery with Densiron. There were no significant differences in final reattachment when grouped by age, diabetic status or number of inferior retinal tears. Males ($p=0.024$) and pseudophakic patients ($p=0.007$) showed a higher rate of final reattachment with recurrent surgery compared with primary surgery. Reattachment rates for primary and recurrent RRD surgery showed a general decrease with increasing severity of PVR, however, differences between

grades did not differ significantly ($p=0.162$). Patients in our cohort achieving final reattachment following Densiron removal underwent a mean of 2.01 (range 1–5) surgeries.

Secondary outcomes

Complete BCVA data were available for 85 eyes in both baseline and 3 months post-Densiron 68 removal. The mean VA improved slightly from baseline to 3 months (mean (SD): 1.19 (0.10) at baseline, to 0.97 (0.08) at 3 months; a slight increase of 0.22 (95% CI: 0.001 to 0.44, $t(84)=1.996$, $p=0.049$). Forty-nine eyes (36.6%) developed anterior uveitis during the initial (first 2 weeks) postoperative period, including one patient who developed an inflammatory hypopyon (0.75%). Long-term anterior uveitis which required treatment was present in 8 patients (6.0%), of whom two had retained long-term Densiron 68 tamponade. There was no association between inflammation and duration of Densiron 68 tamponade ($p=0.779$).

Forty-nine eyes (36.6%) required topical intraocular pressure (IOP) control postoperatively. Long-term raised IOP persisted in 24 eyes (17.9%). Eight patients (6.0%) required surgical intervention to lower IOP. Interestingly, none of the patients with long-term Densiron 68 retention required long-term antihypertensive drops, although two required surgical intervention. There was no correlation between long-term IOP issues and duration of tamponade ($p=0.358$).

Six eyes (4.5%) had evidence of Densiron 68 emulsification. Those patients with long-term tamponade were not recorded to have emulsification. There was no correlation between emulsification rate and duration of tamponade ($p=0.984$).

Ten eyes (7.5%) remained phakic by the 3 months postoperative point. There was no change to pseudophakic or aphakic patients' lens status.

DISCUSSION

The total final reattachment rate of 73.9% from this study of heavy oil tamponade for RRD compare favourably with previous published reports of 33%–92%.^{7 15 18 19} This total comprised 64.6% of primary surgeries and 79.1% of surgeries for recurrent RRD. Although this difference did not quite reach statistical significance, it does

Table 2 Primary outcomes by grade of proliferative vitreoretinopathy (PVR)

PVR	N (%)	Primary success	Final reattachment	Qualified success
0	74 (55.2)	48 (64.9)	60 (81.1)	3 (4.1)
A	7 (5.2)	4 (57.1)	5 (71.4)	0 (0.0)
B	14 (10.4)	7 (50)	10 (71.4)	1 (7.1)
C	39 (29.1)	18 (46.2)	24 (61.5)	7 (17.9)
	P value	0.259	0.162	0.088
Total	134	65 (48.5)	99 (73.9)	11 (8.2)

P values are shown in bold.

**Table 3** Primary outcomes by indication for surgery—primary versus recurrent RD

	N (%)	Primary success	Final reattachment	Qualified success
Primary	48 (35.8)	21 (43.8)	31 (64.6)	3 (6.3)
Recurrent	86 (64.2)	45 (52.3)	68 (79.1)	8 (9.3)
	P value	0.341	0.067	0.537
Total	134	65 (48.5)	99 (73.9)	11 (8.2)

P values are shown in bold.
RD, retinal detachment.

suggest that those cases having Densiron 68 for primary RRD may have a higher failure rate. This may reflect the challenging nature of the cases with primary RRD that had Densiron 68 chosen as a tamponade. A similar reported primary success rate of 71.4% has previously been published in one large series.¹⁸ Primary surgery success for inferior PVR with light silicone oil has been reported to be as high as 82%,²⁰ however, this rate was detected in a small single-centre retrospective subgroup of 22 patients and larger scale data are lacking. Our study is multicentred and larger than all previous reports and is therefore encouraging. The major indication was inferior PVR C. Pseudophakic patients achieved better results than phakic patients undergoing recurrent surgery in our cohort, while among males, those undergoing recurrent surgery had better outcomes than those undergoing primary surgery. As discussed above, this may reflect the nature of primary RRD cases and Densiron use, or the early use of Densiron 68 in failed cases. No

other significant differences were seen between pooled or subgroup data.

Herbrig *et al* provided a large previously published series (99 eyes) on Densiron 68 for the treatment of complicated RRD. 74% of their cases had previously failed RRD surgery, which is in keeping with our indications and suggestive of the most common approach for use of HSO. Seventy-eight of 89 eyes (87.6%) that underwent PPV and Densiron 68 endotamponade showed retinal reattachment at 9 months, with 34 eyes (36%) requiring reoperation to achieve reattachment. These further operations included light silicone oil, second heavy oil or gas tamponade. In all eyes with a second Densiron 68 tamponade, the retina was completely reattached. This series compared 21 prospective patients with no previous history of retinal surgery before receiving Densiron 68, with a retrospective similar control group of 21 patients that had received light silicone oil tamponade. Fifteen eyes with Densiron 68 (71%) showed reattachment at 3

Table 4 Primary outcomes between different subgroups divided by baseline characteristics

Baseline characteristics	Subgroup	N		Final reattachment (%)		
		P	R	P	R	
Age	10–64	25	30	17 (68)	24 (80)	P=0.309
	65–91	23	56	14 (61)	44 (79)	P=0.106
				P=0.606	P=0.877	
Gender	Female	15	31	12 (80)	24 (77)	P=0.842
	Male	33	55	19 (58)	44 (80)	P=0.024*
				P=0.132	P=0.778	
Lens status	Phakic	32	38	22 (69)	26 (68)	P=0.976
	Pseudophakic	16	48	9 (56)	42 (88)	P=0.007*
				P=0.393	P=0.031*	
Diabetic status	Diabetic	6	9	3 (50)	8 (89)	P=0.095
	Non-diabetic	42	77	28 (67)	60 (78)	P=0.181
				P=0.425	P=0.444	
No of inferior retinal tears	0	6	16	5 (83)	15 (94)	P=0.116
	1	16	29	11 (69)	21 (72)	P=0.915
	2	14	14	7 (50)	12 (86)	P=0.373
	3+ or GRT	8	11	5 (63)	7 (64)	P=0.981
	Not documented	4	17			
				P=0.509	P=0.193	

Asterisk (*) denotes a statistical significance between subgroups.
GRT, giant retinal tear; P, primary surgery; R, recurrent surgery.

months while the heavy oil was still in situ, compared with only nine eyes (43%) with light silicone oil. While it failed to reach statistical significance ($p=0.059$), this difference may suggest greater efficacy of Densiron 68 in such cases.¹⁸ In contrast the HSO study, a multicentre randomised masked controlled trial by Jousseaume *et al*, suggested lower reattachment rates with HSO (28.3%) compared with standard silicone oil (1000 or 5000 centistokes; 40.4%) in the treatment of complicated inferior RRD.¹⁵ This trial also failed to show statistical difference between the two different oil tamponades in terms of reattachment rates or VA at 12 months follow-up. However, the trial was stopped early due to slow recruitment and may not have been adequately powered as initially planned.¹⁵ In our series we had a higher anatomical success rate (61.5%) in inferior PVR C cases with Densiron 68 compared with the HSO study, accepting that our data is retrospective. Furthermore, we do not compare to light oil so any comparisons with historical data must be tentative. Our findings from the largest dataset on this topic (a decade later) provide consideration to perhaps review this subject prospectively. The causes for redetachments were not investigated as part of this study.

The HSO study did not find any difference in vision comparing HSO against standard silicone oil, but there was an improvement from baseline in both groups.¹⁵ While our study showed a VA improvement from baseline to post-oil removal, it failed to show statistical significance, supporting the findings of Kocak and Koc.²¹ This may reflect the complex nature of these eyes in which retinal reattachment may be a primary goal. However, other HSO studies did show statistically significant improvement in vision and anatomical reattachment.^{18 19 22}

The reported adverse effects of HSO tamponade include raised IOP, inflammatory response, cystoid macular oedema, oil dispersion/emulsification and cataract formation.^{23 24}

In our series, 49 patients (36.6%) required topical treatment for IOP control. Of those, long-term raised IOP persisted in 24 eyes (17.9%). In their smaller series, Caporossi *et al* found a rate of 34.7% of patients requiring topical treatment, reducing to 26.5% after Densiron 68 removal.²⁵ Other published series showed OHT rates of 12%–27%.^{15 18 23 26} Eight patients in our cohort (6.0%) required surgical intervention for IOP management, whereas Stappeler *et al*'s prospective cohort of 122 patients did not require any surgical IOP-lowering treatment, describing a steady decline in OHT rate over 3 months post-HSO removal.⁷

Nguyen *et al* found that 42% of patients required removal of oil and/or glaucoma surgery to lower their IOP during the early phase, and 32% required medication for chronic OHT.²⁷ This study had a much higher reported incidence of OHT (48%) and this could be attributed to having a small cohort of 47 patients. Our rates were lower and may represent a closer reflection of the true rate, considering the size of our cohort. Whether Densiron 68 has a greater effect on IOP than light silicone oil is unclear; but our data suggest similar rates, although we acknowledge we have no direct control group.

HSOs are suggested to promote an inflammatory response.^{6 28} In our series, 49 patients (35.7%) developed anterior uveitis within the postoperative period. Following Densiron 68 removal, 8% of patients required long-term topical anti-inflammatory treatment. Auriol *et al* reported a 40.7% rate of anterior chamber inflammation with fibrin accumulation and a mean time to removal of Densiron 68 of 14 weeks.²⁹ However, Kocak and Koc found a much lower rate (3%) of anterior inflammation with fibrin.²¹ Russo *et al*'s review of HSO concluded that inflammation may depend on tamponade duration.²⁸ However, data analysis of our cohort did not reveal any association between inflammation and duration of Densiron 68 tamponade. Our data must be interpreted with caution, as prospective objective analysis of inflammation was not undertaken. Data on long-term inflammation rates with light silicone oil is limited.

Emulsification is also recognised as a complication of heavy and light silicone oil.⁶ It has been reported to occur in 8.3%–16% of patients with HSO.^{7 18 19 30} In our series, the emulsification rate was 10.1%, in keeping with reported emulsification rates of Densiron 68. Densiron Xtra (a newer tamponade) is promoted as being more resistant to emulsification due to its higher viscosity and longer silicone oil molecular chain.³¹ In vivo confirmation is still awaited.³² Emulsification of light silicone has been reported to occur in up to 40% of cases with long chain silicone oil (5700 cSt)³³ and 63.4% in shorter chain silicone oil (1300 cSt). Other series have reported similar disparities over a mean period of 7.3 months (SD 4.2 months, range 1–17 months).³⁴ Our rates of emulsification with Densiron 68 are therefore reassuring.

Cataract is a recognised consequence of PPV.²⁶ While our study did not specifically look at progression of cataract, 85.7% of patients required lens extraction. This could be secondary to multiple factors: pre-existing lens status, previous tamponades, multiple surgeries or use of steroids.⁶ Our series has similar lens findings to a series of 122 patients by Stappeler *et al*,⁷ however, Jousseaume *et al* did not show conclusive evidence of cataract progression in their randomised control trial.¹⁵

There are several limitations to this study. Its retrospective nature limited more detailed interrogation of the outcomes and complications of Densiron 68. Unlike previous published case series, our data offer a high proportion of patients receiving Densiron 68 as a primary endotamponade. We did not collect data on superior retinal breaks, therefore while it is unlikely that any patient with superior pathology received Densiron as primary endotamponade, we cannot exclude this as a possibility that may have contributed to our relatively high redetachment rate. Therefore, this patient group is significantly different from others in published literature, so we recommend cautious interpretation of our results when compared with other studies. Subgroups for analysis were retrospective and therefore limited in interpretation. It is possible that not all cases of Densiron 68 use were identified across all centres, which may represent a source of bias in the study. However, this process was optimised using a

combination of personal logbooks, theatre records and electronic patient records. We were unable to compare Densiron to light silicone oil in this study, nor were we able to investigate causes for retinal redetachment.

However, this is the largest series presenting real-world, long-term, multicentre UK data on the use and outcomes of surgery with Densiron 68. Densiron 68 is perhaps not widely used due to lack of familiarity with the injection and removal process. The data presented here should reassure surgeons that Densiron 68 is a useful tamponade alternative for the management of complex inferior RRD, with similar morbidity to standard silicone oil.

Correction notice This article has been corrected since it first published. Affiliation 9 has been updated.

Contributors SD wrote the initial draft and analysed data. MD analysed data, made major reworkings to the text and co-ordinated author responses before producing the final manuscript. TS and JK made invaluable suggestions during the editing process. JZ provided valuable statistical input. AC was supervising author and guarantor. Remaining authors all helped with data collection.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Max Davidson <http://orcid.org/0000-0002-3895-3953>

Megan Kay Wood <http://orcid.org/0000-0003-1975-477X>

REFERENCES

- Schwartz SG, Flynn HW, Wang X, *et al.* Tamponade in surgery for retinal detachment associated with proliferative vitreoretinopathy. *Cochrane Database Syst Rev* 2020;5:CD006126.
- Fleissig E, Barak A, Goldstein M, *et al.* Massive subretinal and subretinal pigment epithelial hemorrhage displacement with perfluorocarbon liquid using a two-step vitrectomy technique. *Graefes Arch Clin Exp Ophthalmol* 2017;255:1341–7.
- Wickham L, Ho-Yen GO, Bunce C, *et al.* Surgical failure following primary retinal detachment surgery by vitrectomy: risk factors and functional outcomes. *Br J Ophthalmol* 2011;95:1234–8.
- Wickham L, Bunce C, Wong D, *et al.* Retinal detachment repair by vitrectomy: simplified formulae to estimate the risk of failure. *Br J Ophthalmol* 2011;95:1239–44.
- Adelman RA, Parnes AJ, Michalewska Z, *et al.* Clinical variables associated with failure of retinal detachment repair: the European vitreo-retinal Society retinal detachment study report number 4. *Ophthalmology* 2014;121:1715–9.
- Heimann H, Stappler T, Wong D. Heavy tamponade 1: a review of indications, use, and complications. *Eye* 2008;22:1342–59.
- Stappler T, Heimann H, Wong D, *et al.* Heavy tamponade 2 Densiron 68 in routine clinical practice: anatomical and functional outcomes of a consecutive case series. *Eye* 2008;22:1360–5.
- Charteris DG, Sethi CS, Lewis GP, *et al.* Proliferative vitreoretinopathy-developments in adjunctive treatment and retinal pathology. *Eye* 2002;16:369–74.
- Angunawela RI, Azarbadegan A, Aylward GW, *et al.* Intraocular fluid dynamics and retinal shear stress after vitrectomy and gas tamponade. *Invest Ophthalmol Vis Sci* 2011;52:7046–51.
- Wetterqvist C, Wong D, Williams R, *et al.* Tamponade efficiency of perfluorohexyloctane and silicone oil solutions in a model eye chamber. *Br J Ophthalmol* 2004;88:692–6.
- Roeder J, Hoerauf H, Kobuch K, *et al.* Clinical findings on the use of long-term heavy tamponades (semifluorinated alkanes and their oligomers) in complicated retinal detachment surgery. *Graefes Arch Clin Exp Ophthalmol* 2002;240:965–71.
- Schatz B, El-Shabrawi Y, Haas A, *et al.* Adverse side effects with perfluorohexyloctane as a long-term tamponade agent in complicated vitreoretinal surgery. *Retina* 2004;24:567–73.
- Gremillion CM, Peyman GA, Liu KR, *et al.* Fluorosilicone oil in the treatment of retinal detachment. *Br J Ophthalmol* 1990;74:643–6.
- Long-Term Tamponades: Geuder AG. Available: <https://www.geuder.de/en/products/biomaterials/long-term-tamponades/> [Accessed 30 Jun 2020].
- Joussen AM, Rizzo S, Kirchhof B, *et al.* Heavy silicone oil versus standard silicone oil in as vitreous tamponade in inferior PVR (HSO study): interim analysis. *Acta Ophthalmol* 2011;89:e483–9.
- Keilani C, Augstburger E, Robin M, *et al.* Comparative biochemical outcomes, effectiveness and tolerance of Densiron 68 and Oxane HD for the management of complicated retinal detachment. *Turk J Ophthalmol* 2019;49:334–41.
- Holladay JT. Visual acuity measurements. *J Cataract Refract Surg* 2004;30:287–90.
- Herbrig E, Sandner D, Engelmann K. Anatomical and functional results of endotamponade with heavy silicone oil - Densiron 68 - in complicated retinal detachment. *Ophthalmic Res* 2007;39:198–206.
- Wong D, Van Meurs JC, Stappler T, *et al.* A pilot study on the use of a perfluorohexyloctane/silicone oil solution as a heavier than water internal tamponade agent. *Br J Ophthalmol* 2005;89:662–5.
- Sheng Y, Sun W, Mo B, *et al.* Non-buckled vitrectomy for retinal detachment with inferior breaks and proliferative vitreoretinopathy. *Int J Ophthalmol* 2012;5:591–5.
- Kocak I, Koc H. Comparison of Densiron 68 and 1 000 cSt silicone oil in the management of rhegmatogenous retinal detachment with inferior breaks. *Int J Ophthalmol* 2013;6:81–4.
- Sandner D, Engelmann K. First experiences with high-density silicone oil (Densiron) as an intraocular tamponade in complex retinal detachment. *Graefes Arch Clin Exp Ophthalmol* 2006;244:609–19.
- Duan A, She H, Qi Y. Complications after heavy silicone oil tamponade in complicated retinal detachment. *Retina* 2011;31:547–52.
- Hostovsky A, Yap J, Mandelcorn MS, *et al.* DENSIRON 68 heavy silicone oil as a short-term intraocular tamponade for MACULA-ON inferior retinal DETACHMENTS-A case series. *Retin Cases Brief Rep* 2022;16:581–585.
- Caporossi T, Franco F, Finocchio L, *et al.* Densiron 68 heavy silicone oil in the management of inferior retinal detachment recurrence: analysis on functional and anatomical outcomes and complications. *Int J Ophthalmol* 2019;12:615–20.
- Li W, Zheng J, Zheng Q, *et al.* Clinical complications of Densiron 68 intraocular tamponade for complicated retinal detachment. *Eye* 2010;24:21–8.
- Nguyen QH, Lloyd MA, Heuer DK, *et al.* Incidence and management of glaucoma after intravitreal silicone oil injection for complicated retinal detachments. *Ophthalmology* 1992;99:1520–6.
- Russo A, Morescalchi F, Donati S, *et al.* Heavy and standard silicone oil: intraocular inflammation. *Int Ophthalmol* 2018;38:855–67.
- Auriol S, Pagot-Mathis V, Mahieu L, *et al.* Efficacy and safety of heavy silicone oil Densiron 68 in the treatment of complicated retinal detachment with large inferior retinectomy. *Graefes Arch Clin Exp Ophthalmol* 2008;246:1383–9.
- Dooley IJ, Duignan ES, Kilmartin DJ. Long-term heavy silicone oil intraocular tamponade. *Int Ophthalmol* 2016;36:3–7.
- Densiron® Xtra - new heavy silicone oil for intraoperative complication management: Geuder AG. Available: https://www.geuder.de/en/news-events/press-center/press-releases/?tx_news_pi1%5Baction%5D=detail&tx_news_pi1%5Bcontroller%5D=News&tx_news_pi1%5Bnews%5D=22&chash=a9088ae09904af514731a5bace08aed [Accessed 10 Jul 2020].
- Caramoy A, Schröder S, Fauser S, *et al.* In vitro emulsification assessment of new silicone oils. *Br J Ophthalmol* 2010;94:509–12.
- Ratanapakorn T, Thongmee W, Meethongkam K, *et al.* Emulsification of different viscosity silicone oil in complicated retinal detachment surgery: a randomized double-blinded clinical trial. *Clin Ophthalmol* 2020;14:359–67.
- Zhao X-J, Tang N-N, Lian Y, *et al.* Analysis of the rates of emulsification in intraocular silicone oil tamponades of differing viscosities. *Int J Ophthalmol* 2020;13:761–5.